

type and paratypes collected at Gialiskari — Ikaria island, Greece, on 15.VII.1981. Leg. S. Drosopoulos, in collection S. Drosopoulos of the Benaki Phytopathological Institute.

The new species seems to be ecologically and geographically separated from its close relative *F. leptosoma*. Ecologically, it can be considered a coastal species being separated from *F. leptosoma* which is not found in such habitats in Greece. Geographically, it is an insular species probably endemic to Ikaria, because investigations in other islands close to Ikaria in similar biotopes neither *F. leptosoma* nor *F. mourikisi* was found. However, Ikaria is a unique island because many other delphacids (e.g. *Iubsoda stigmatica* (Melichar, 1897), *Kelisia melanops* Fieber, 1878, *Alatades trilineatus* Dlabola, 1957, *Ditropis pteridis* (Spinola, 1839)), which are present on Ikaria, were not found on Naxos or Paros.

The new species is named after the director of the Benaki Phytopathological Institute Dr. P. Mourikis, who has contributed greatly to the development of entomology in Greece.

ACKNOWLEDGEMENTS

I thank Drs. M. Asche (Marburg, W. Germany), J. Dlabola (Praha, Czechoslovakia) for their comments on the material of the new *Florodelphax* species, Prof. F. E. Strong (F.A.O. project, Greece) for editing the manuscript; Drs. F. M. Muller (Wageningen, The Netherlands) and M. Damanakis for identifying some of the grass species; Dr. C. J. H. Booij (Wageningen, The Netherlands) for supplying material of *M. extrusa*; Miss V. Kapothanasi for valuable assistance of mounting the material and typing the manuscript, and the authorities of the Benaki Phytopathological Institute for supporting my hemipterological studies in Greece.

LITERATURE

- Booij, C. J. H., 1981. Biosystematics of the Muellerianella complex (Homoptera, Delphacidae), Taxonomy, morphology and distribution. — *Neth. J. Zool.* 31 (3): 572-595.
- Drosopoulos, S., 1977. Biosystematic studies on the Muellerianella complex (Delphacidae, Homoptera — Auchenorrhyncha). — *Meded. Landbouwhogeschool Wageningen* 77: 1-133.
- , 1982. Hemipterological studies in Greece. Part II Homoptera-Auchenorrhyncha. On the family Delphacidae. — *Marburger ent. Publ.* 1 (6): 35-88.
- Morris, M. G., 1974. Auchenorrhyncha (Hemiptera) of the Burren, with special reference to species associations of the grasslands. — *Proc. R. Ir. Acad.* 74 (B): 7-30.
- Nast, J., 1972. *Palaeartic Auchenorrhyncha (Homoptera), an annotated checklist*: 1-550. Polish Scientific Publishers Warsaw.
- Ossiannilsson, F., 1978. The Auchenorrhyncha (Homoptera) of Fennoscandia and Denmark. — *Fauna ent. Scand.* 7 (1): 1-222.
- Vilbaste, J., 1971. *Die Zikaden Estlands I*. Tallinn: 1-284 (In Estonian).

LUCANUS CERVUS (LINNAEUS) IN NEDERLAND (COL.: LUCANIDAE). Enige jaren geleden werd in dit tijdschrift (1980, deel 40: 24) door ons het verzoek gedaan informatie te verstrekken over het vliegend hert. Inmiddels is over het voorkomen van *Lucanus cervus* in Nederland een rapport verschenen in de Nieuwsbrief EIS-Nederland 12 (1982): 35-43, waarin de verspreiding, biologie, kenmerken en natuurbeschermingsaspecten worden behandeld. *Lucanus* is in totaal (ca. 1830-1980) op minstens 69 plaatsen in Nederland waargenomen, op minstens 16 plaatsen nog na 1960. De waarnemingen zijn vooral geconcentreerd in de omgeving van Apeldoorn, van Nijmegen, en in Zuid-Limburg.

Overigens is zojuist over Lucanidae in het algemeen een aardig boekje verschenen in de serie „Die Neue Brehmbücherei”, no. 551. Het is van B. Klausnitzer, 1982, Hirschkäfer oder Schröter (Lucanidae), 83 pp., 55 fig., 1 pl., A. Ziemsen Verlag (Wittenberg Lutherstadt), ISSN 0138-1423. Wij hebben er bij onze boekhandel f 13,25 voor betaald.

J. Krikken & H. C. Pijpers, Rijksmuseum van Natuurlijke Historie, Postbus 9517, 2300 RA Leiden.