

The variability in wing neuration of *Arenipses sabella* Hampson, 1901 (Lepidoptera, Pyralidae)

by

R. H. KLEINPASTE

Department of zoology, Agricultural University, Wageningen

ABSTRACT. — It has been shown that in *Arenipses sabella* Hampson there is a considerable variability in the pattern of wing venation. This variability appears to be greater in females than in males. It is concluded that this intraspecific variation creates difficulties for the classification of the Galleriinae (the subfamily of the Pyralidae to which *sabella* belongs), because the latter is mainly based on the patterns of wing venation.

Thus far the study of the variability of wing venation in Lepidoptera has been hardly developed. In my opinion there is a necessity to pay special attention to this variability in those taxonomic groups in which the classification is primarily based on configuration of the wing venation e.g. the subfamily Galleriinae of the Pyralidae (Hampson, 1917).

During a stay at the British Museum (Natural History) in London last year, I had the opportunity to study the intraspecific variability in wing venation in *Arenipses sabella* Hampson. This species occurs in the southern Mediterranean area and the Middle East. It has been described by Hampson (1901: 501) from Arabia. From literature it is known that the larvae mainly feed on dates both in the palmtrees and in the stored product (Buxton, 1920). Transport of dried dates all through the world may therefore spread *sabella* everywhere.

Venation of the males. Examined: 20 specimens.

A. Forewing. The general pattern is shown in fig. 1a. The large cell, which is clothed with fine silky hair on the underside, is produced to a point at vein 5. All twelve veins were always present but vein 5 was sometimes inconspicuous. Vein 10 can rise from nearly any point in between vein 11 and the stalk of the complex 7, 8 and 9. The two extremes are shown in fig. 1b by dashed lines.

B. Hindwing. In the hindwings of the males of *sabella* vein 5 is absent. In these wings the neuration shows more variation than the neuration in the forewings. 16 specimens (80%) showed a configuration of vein 6 as presented in fig. 1c, while 4 specimens (20%) showed a configuration as drawn in fig. 1d. Moreover, vein 3 has a variable length, as it varies in point of rising from vein 4. Fig. 1c and 1d also show the two extremes in length that have been found.

Venation of the females. Examined: 20 specimens.

A. Forewing. The forewings of the females of *sabella* show all twelve veins. The pattern of neuration appeared to be extremely variable. Taking vein 6, two extremes and one "intermediate situation" have been indicated. Vein 6 can be stalked with the veins 7, 8 and 9 and consequently does not rise from the cell (fig. 2a). This configuration was found in 12 females (60%). Vein 6 can also rise from the cell at the same point from which the stalk of veins 7, 8 and 9 rises (fig. 2b). The other extreme consists of vein 6 rising from the cell separately from the stalk of veins 7, 8 and 9 (fig. 2c); the latter configuration occurred in 7 specimens (35%). Only one "intermediate form" was found. I never observed the cell entirely closed. Veins 4 and 5 both rise from the cell either from one point (fig. 2d) or from two separate points (fig. 2e), the former configuration being found as often as the latter. One of the females showed a configuration in which vein 11 does not rise from the cell, but is stalked with vein 10 (fig. 2f).

B. Hindwing. The hindwing venation of the females of *sabella* is drawn in fig. 2g. The neuration shows hardly any variability, apart from very small differences in length of vein 3. The variability of the length of the stalk, indicated in fig. 2g as (p-q), deserves closer attention. Its varying length is mainly due to the convergence of point p on point q, without actually touching point q. Apparently the venation of the females of *sabella* shows more variation when compared to the males.




Fig. 1. *Arenipses sabella* Hampson. ♂. Explanation: see text. a-b. Patterns of forewing venation. c-d. Patterns of hindwing venation.




Fig. 2. *Arenipses sabella* Hampson. ♀. Explanation: see text. a-f. Patterns of forewing venation. g. Pattern of hindwing venation.

The variability in neuration as described in this paper is a general feature among the North West European species belonging to the subfamily of the Galleriinae (Kuchlein & Kleinpaste, 1977, in press). This makes it extremely difficult to maintain many of the distinguishing marks of venation as significant features on generic or specific level. Apparently the existing keys in the handbooks do not hold for a considerable fraction of the populations!

I am indebted to the Agricultural University (Wageningen) for supporting the research in London and Dr. P. E. S. Whalley and Mr. M. Shaffer for their assistance in the British Museum (Natural History). I would also like to thank Drs. J. H. Kuchlein for his kind assistance and, together with Miss C. M. A. de Goey, for correcting this manuscript thoroughly.

REFERENCES

- Buxton, P. A., 1920. Insect pests of dates and the date palm in Mesopotamea and elsewhere. - *Bull. ent. Res.* 11: 287-303 + pls.
- Hampson, G. F., 1917. A classification of the Pyralidae, subfamily Gallerianae. - *Novit. zool.* 24: 17-58.
- Kuchlein, J. H. & R. H. Kleinpaste, 1977. *Lamoria anella* (Denis & Schiffermüller), new to the Netherlands (Lepidoptera, Pyralidae). - *Ent. Ber., Amst.* in press.
- Hampson, G. F. in E. L. Ragonot & G. F. Hampson, 1901. Monographie des Phycitinae et des Galleriinae. - Pars II, *Romanoff Mém.* 8, 1-602.
- Ritzema Bosweg 32a, 6703 AZ Wageningen, the Netherlands.

NIEUWE AANWINSTEN VOOR DE BIBLIOTHEEK

- ANDREWS, M., 1976, The life that lives on man.
- BUTLER, C. G., 1974, The world of the honeybee.
- CHORINE, V., 1931, Contribution à l'étude de l'immunité chez les insectes. Diss. Paris.
- CODREANU, R., 1939, Recherches biologique sur un chironomide *Symbiocladius rhitrogenae* (Zav.), ectoparasite cancérigène des éphémères torrenticoles. Diss. Paris.
- CORNWELL, P. B., 1976, The Cockroach vol. II. Insecticides and cockroach control.
- EVANS, L. K. & K. G. W. Evans, 1973, A survey of the Macro-Lepidoptera of Croydon and North-East Surrey.
- FRANZ, H., 1975, Die Bodenfauna der Erde in biozönotischer Betrachtung. vol. I—II.
- GOMEZ BUSTILLO, M. R. & F. Fernandez Rubio, 1976, Mariposas de la Peninsula Ibérica. Vol. III. Heterocera.
- HINTON, H. E. & A. S. Corbet, 1972, Common insect pestst of stored food products. A guide to their identification, 5th ed.
- INSECTS and plants of the Alver estuary, 1976 (D. Appleton a.o.).
- IWATA, K., 1976, Evolution of instinct. Comparative ethology of Hymenoptera.
- JOHNSON, W. T. & H. L. Lyon, 1976, Insects that feed on trees and shrubs.
- KLOET, G. S. & W. D. Hincks, 1977, A check list of British Insects, 3, Coleoptera and Strepsiptera, 2nd ed. (Handbook Ident. brit. Ins. 11(3)).
- KULLMANN, E. & H. Stern, 1975, Leven aan een zijden draad; fascinerende wereld van de spinnen.
- KUZNETSOVA, N. Ya., 1972 (1916), A review of the family Sphingidae of the Palaearctic and Chinese-Himalayan faunas. Translation.
- LIST of serial publications in the British Museum (Natural History) 2nd. ed., 1975.
- PERSPECTIVES in forest entomology, 1976 (J. F. Anderson & H. K. Kaya, eds.).
- SPENCER, K. A., 1976, The Agromyzidae of Fennoscandia and Denmark. (Fauna ent. scand. 5(1/2)).
- ZEUNER, F. E. & F. J. Manning, 1976, A monograph of fossil bees (Hymenoptera: Apoidea), (Bull. brit. Mus. (nat. Hist.) Geology 27(3)).

BIBLIOGRAFIE VAN TABELLEN TOT DE EUROPESE INSEKTEN. Als vervolg op Göllner-Scheidung's „Bibliographie der Bestimmungstabellen europäischer Insekten, 1880—1963” (*Beitr. Ent.*, 17(1957): 697—858; *Mitt. zool. Mus. Berl.*, 45(1969): 3—156; *Dtsch. ent. Z.*, (N.F.), 17(1970): 33—118, 433—476) is er van de hand van Reinhard Gaedike, onder dezelfde titel, in *Beiträge zur Entomologie* 26(1976): 49—166, een overzicht verschenen van de tabellen, gepubliceerd in 1964—1973. Deze bibliografie refereert aan 1570 publicaties, alle daarin behandelde hogere taxa zijn in de index (pag. 156—166) per orde gerangschikt. Alhoewel zeer uitvoerig en waardevol, bestaat de indruk dat alleen artikelen zijn opgenomen die in reguliere tijdschriften zijn gepubliceerd; de tabellen, verschenen als wetenschappelijke mededeling der KNNV zijn bijvoorbeeld niet opgenomen. — P. Oosterbroek.