

## Two new Mordellidae (Coleoptera) from Southern Europe, and a key to the *Mordellistena micans* group

by

R. BATTEN

**ABSTRACT.** — Two species of Mordellidae from southern Europe are described: *Variimorda* (*Sulcatimorda*) *krikkeni* n.sp. and *Mordellistena wiebesi* n.sp. A key to the West Palaearctic species of the *Mordellistena micans* group is given, with illustrations of each species. A lectotype for *Mordellistena stenidea* is selected.

### INTRODUCTION

During a foray in Macedonia I caught some Mordellidae which represent two new species, of *Variimorda* and *Mordellistena* respectively, which are described in this paper. I have inserted the new *Mordellistena* in a key to the species of the *micans* group.

The following abbreviations are used:

AC — author's collection.

MA — Zoölogisch Museum (Instituut voor Taxonomische Zoölogie) van de Universiteit van Amsterdam.

ML — Rijksmuseum van Natuurlijke Historie, Leiden.

MP — Muséum National d'Histoire Naturelle, Paris.

PT — collection Méquignon, property of Mr. P. Teocchi, Sérignan near Orange (France).

SMD — Staatliches Museum für Tierkunde, Dresden.

*Variimorda* (*Sulcatimorda*) *krikkeni* n.sp. figs. 3, 6, 7.

This Mordellid has the usual features of the genus *Variimorda*, and belongs to the subgenus *Sulcatimorda* because the pygidium is clearly depressed at both sides above the lateral grooves. This species however lacks the patches of white or yellow pubescence characteristic of the other species in the genus.

**Description.** — Dimensions: ♂: length of the body 4.5-5.5 mm (without pygidium); length of elytra 3.0-3.6 mm, width 1.6-1.9 mm; pygidium 1.5-1.8 mm; fifth sternite 1.0-1.1 mm. ♀: length of body 4.6-6.6 mm; length of elytra 3.1-4.2 mm; width 1.65-2.3 mm; pygidium 1.5-2 mm; fifth sternite 0.8-1.1 mm.

Head transverse, width/length ratio ♂ ca. 1.2-1.5, ♀ ca. 1.1-1.3, finely punctured, hind margin convex, eye minutely granulated and hairy. Galea long and narrow, without appendages, and ending in a very fine point. Antenna segments of the ♂ 1-4 and partly 5 reddish brown, ♀ darker. Maxillary palp of the ♂ reddish brown, apex of segment 4 black, ♀ brown black, segment 4 black.

Pronotum slightly broader than elytra at shoulders, width/length ratio ♂ ca. 1.3, ♀ 1.2-1.4, punctures slightly larger than on head, anterior angles convex, basal angles obtuse and rounded at edge; scutellum approximately semicircular.

Elytra black, densely punctate, punctation stronger than on pronotum, pubescence black, length/width ratio 1.8-1.9 (1 ♂ 1.94), sides from shoulders strongly attenuated posteriorly (fig. 3).

Integument of underside black, pubescence black.

Pygidium clearly depressed at both sides above the lateral grooves, length 1.35-1.8 times that of fifth sternite, about 3 times as long as broad.

Legs black, ♂ anterior tibiae reddish brown to brownish black. Spurs of hind leg black with dark reddish point.


Parameres: ventral branch of right paramere with rounded and reflexed apex (figs. 6, 7).

Material examined. — Greece (Mak.) 15 km W. of Édhessa, 6.VII.1976, ca. 100 m, leg.

R. Batten (1 ♂ holotype, 1 ♀ allotype ML, 3 ♂ and 3 ♀ paratypes AC), roadside on flowering *Daucus* and other Umbelliferae, slopes with *Quercus coccifera*, shrubs, *Euphorbia*.

Discussion. — The galeae of *Variimorda fagniezi*, *theryi* and *krikkeni* are similar; *V. fagniezi* has its elytra two times as long as their combined width at base; *theryi* is similar in this character (according to the type-specimens I studied), though Méquignon (1946: 60) writes they are 1.5 times as long as broad. The new species, however, has strongly attenuated elytra 1.8-1.9 as long as broad; the pygidium is a bit longer in proportion to the fifth sternite, while the parameres differ as well (figs. 6, 7).

I dedicate this new species to Mr. J. Krikken of the ML, for his kind advice.



Figs. 1—6. - 1, 4, *Variimorda fagniezi* (Spain, Barcelona, Avinyo); 2, 5, *V. theryi* (cotype Maroc la Mimouna); 3, 6, *V. krikkeni* (holotype). - 1—3 elytra and pygidium; 4—6 combined parameres, dextrolateral view.



*Variimorda theryi* (Méquignon) figs. 2, 5.

*theryi* Méquignon, 1946: 73-75, (*Mordella* subgenus *Sulcatimorda*); Ermisch 1969: 166 (in *Variimorda*).

Discussion. — The types of *V. theryi* differ from *fagniezi* in the elytra, which in *theryi* are a bit more parallel posteriorly (figs. 1, 2), and in the colour of the middle tibia, being reddish in the male. Méquignon (1946: 75) writes: „Extérieurement les différences sont en effet minimes”. The difference in the parameres is that in *theryi* the ventral branch of the right paramere is abruptly bent in the middle, the distal part obliquely directed to the inner side (fig. 5). After having compared the parameres of *theryi* with those of my specimens agreeing with *fagniezi* in the two external characters just mentioned, I can only conclude that the difference is small (figs. 4, 8).

Material examined. — Morocco: labelled as follows: “Rabat”, “♂ Théryi Méq.”, red “type” (1 ♂ holotype); “fasciata F.”, “Marocco, Casablanca, Reitter”, “M. Théryi var. maroccana Méquign.”, red “type” (1 ♂); “1. Mimouna, Maroc. coll. Théry”, “M. Théryi var. nigrescens”, red “type” (1 ♂); “la Mimouna Maroc, coll. Théry”, “♂ Théryi n.sp.”, red “cotype” (1 ♂ with mounted parameres fig. 5); “Mamora”, “Maroc, coll. Théry”, “♀ Théryi Méquign.” and “♂ Théryi n.sp.”, red “cotype” (1 ♂ and 1 ♀) PT.

*Variimorda fagniezi* (Méquignon) figs. 1, 4, 8.

*fagniezi* Méquignon, 1946: 73, 74 (*Mordella* subgenus *Sulcatimorda*), Ermisch 1969: 166 (in *Variimorda*).

Material examined. — Many specimens from Spain, AC.

*Mordellistena micans* group

The species of the *micans* group belong to *Mordellistena* s.str., in fact the first section of Ermisch, having the first four segments of the antenna narrower and shorter than the fifth segment. The group is characterised by the following features: colour of pubescence light; at least three oblique ridges (apart from apical one) on the hind tibia parallel to the distal end; only the first and second tarsal segments with ridges; punctures on elytra finely asperate; pygidium long and slender (broader in *stenidea*); terminal segment of maxillary palp slender securiform, narrower and smaller than those of the *pumila* group.

Discussion. — It should be emphasized that smaller specimens frequently lack the third ridge on the hind tibia, and are then identified as belonging to the *gemellata* group. In that case only the genitalia of the males are decisive, whereas in the females there is a constant character in the eighth urosternite. The females of the *micans* and *gemellata* groups have in the middle of the posterior membrane of the eighth urosternite a chitinised axis being,

a. in the *micans* group:

slender with straight or concave sides (fig. 26) in *minima*, *stenidea*, *grisea*, *micantoides* and *wiebesi* n.sp.;

approximately circular with protruding apex (fig. 27) in *perroudi*, *hirtipes* and *pseudohirtipes*;

b. in the *gemellata* group:

long ovally with more or less convex sides (fig. 28) in *fuscogemellata* Ermisch, *carinthiaca* Ermisch and *dvoraki* Ermisch;


broadened to approximately circular at apex (fig. 29) in *gemellata* Schilsky, *pyrenea* Ermisch and *algeriensis* Ermisch.

In the reverse case this character can be used when there is a third ridge or the beginning thereof in the *gemellata* group.

Note. — *M. aegea* does not belong to the *micans* group, but to the second section of Ermisch, because the fourth segment of the antenna is as long as the fifth.

*Mordellistena wiebesi* n.sp. figs. 9—14.

Description. — Dimensions: ♂ length of body 3.2—5.0 mm without pygidium; length of elytra 2.3—3.5 mm, width 0.9—1.3 mm; pygidium 1.35—1.8 mm; fifth sternite 0.5—0.8 mm. ♀ length of body 3.4—4.2 mm; length of elytra 2.4—3.1 mm, width 0.95—1.25 mm; pygidium 1.1—1.5 mm; fifth sternite 0.5—0.6 mm.



Figs. 7—14. - 7, *Variimorda krikkeni* (holotype); 8, *V. fagniezi* (Spain, Barcelona, Avinyo); 9—14, *Mordellistena wiebesi* (holotype except 12). - 7, 8, 11 parameres inner side, right paramere on left hand, left paramere on right hand; 9, right maxillary palp; 10, eye dextrolateral; 12, right galea; 13, apex of penis; 14, phallobase.



General shape moderately elongate, sides anteriorly parallel, posteriorly slightly convex.

Integument black; pubescence sericeously yellow.

Head almost as long (from point of mandibles to hind margin) as broad, approximately circular, fine and densely punctate, with blue and yellow shagreening, hind margin approximately semicircular. Galea cf. fig. 12. Eyes finely granulated, hairy, temporal margin narrow (fig. 10). Terminal segment of maxillary palp slender, approximately securiform, in ♂ length/width ratio 2.5, inner side slightly rounded, largest at 0.6 of length from base (fig. 9), in ♀ smaller, length/width ratio 2.4, largest at 0.7 of length from base. Antenna in ♂ segment 11 1.3 times segment 10, segments 5—10 nearly parallel shaped, in ♂ 2 times as long as broad, in ♀ 1.7 times.

Pronotum with width/length ratio 1.1, broadest just in front of base; anterior lobe protruding, basal lobe prominent; lateral borders in dorsal view moderately convex, in lateral view concave; basal angles rectangular with a pointed edge; punctures shallowly impressed. Scutellum triangular, sides slightly convex.

Elytra with length/width ratio ♂ 2.5—2.7, ♀ 2.5—2.6; sides moderately elongate slightly convex; elytral apices separately rounded, punctures slightly asperate, pubescence densely covering integument.

Underside with bright pubescence except on last three segments, where it is infuscated.

Pygidium with length in male 2.2—2.9 times that of fifth sternite, in female 2.2—2.5 times; pygidium half as long as elytra; pygidium in dorsal view slender, evenly attenuated, in lateral view slightly convex.

Fore tibia in lateral view slightly convex, not dilated in male, nor with fringe of hairs along inner border; hind tibia with 3—5 ridges apart from apical one, first tarsal segment with 4—6 ridges, second one with 2—3 ridges; outer spur twice as long as inner one.

Parameres, apex of penis, phallobase, cf. figs. 11, 13, 14.

Material examined. — Greece (Mak.): 10 km S. of Kerkini, 100 m, 8.VII.1976, leg. R. Batten, 1 ♂ holotype, 1 ♀ allotype (ML), 5 ♂ and 4 ♀ paratypes (AC), on flowers of Umbelliferae, slopes to Lake Kerkinitis, deciduous trees, shrubs, *Euphorbia*, *Eringium*. Other paratypes as follows: Spain (Barcelona): Montseny, 522 m, 12.VII.1975, leg. R. Batten, (1 ♂ AC) on *Daucus* in deciduous forest; Spain: env. of Playa d'en Pere Fet Cadaques, 5.VIII.1950. leg. L.B. Holthuis (2 ♂ ML) on succulent Umbellifer; Yugoslavia: Dubrovnik 1974 leg. Hladil (2 ♂, one coll. Berger, one AC, kindly presented to me by Mr. C. J. M. Berger); Yugoslavia: Marja (nr. Split), 9.VI.1962, leg. M. H. de Boer (1 ♂ MA); Italy (Bresc.) Idro, 1.VII.1973, leg. P. Kanaar (1 ♂ AC, kindly presented to me by Mr. P. Kanaar).

Discussion. — The male of *M. wiebesi* n. sp. has no fringe of hairs along the inner border of the fore tibia, in which it agrees with *M. grisea*, *micantoides* and *stenidea*. The segments 5—10 of the antenna of the male are in *M. wiebesi* twice as long as broad, whereas they are almost square in *stenidea*, and 1.3 times as long as broad in *grisea* and *micantoides*. Other differences are in the terminal segment of the maxillary palp, the pygidium and the pronotum as described.

I dedicate this new species to Prof. Dr. J. T. Wiebes of the University of Leiden, who recommended to me the study of the Mordellidae.

#### Account of the *micans* group, species and material examined.

##### *Mordellistena hirtipes* Schilsky, fig. 15.

*hirtipes* Schilsky 1895: 46.

Material: Peloponnesis, Kyllini, VII, leg. Muche (1 ♂ and 1 ♀ SMD); Greece, (Mak.) Drama, 7.VII.1976, leg. R. Batten (1 ♂ AC); Greece (Mak.) Kerkini, 8.VII.1976, leg. R. Batten (11 ♂ and 11 ♀ AC).

##### *Mordellistena pseudohirtipes* Ermisch, fig. 16.

*pseudohirtipes* Ermisch 1965: 265, 268, 269.

Material: France (Var) Vidauban, 6.VII.1973, leg. R. Batten (25 ♂ and 25 ♀ AC).

##### *Mordellistena micans* (Germar) fig. 30.

*micans* Germar 1817: 212, *Mordella*; Ermisch 1954: 175—177, *Mordellistena*.




Material: Turkey, Kilios-Kôl., 11.VII.1954, leg. F. Schubert (1 ♂ and 1 ♀ det. Ermisch, parameres mounted SMD).

*Mordellistena minima* (Costa) figs. 20, 26.

*minima* Costa 1854: 18, *Mordella*; Mulsant 1856: 383, *Mordellistena*.

Material: Cyprus, Kyrenia, 9.VI.1939, leg. Håkon Lindberg (1 ♂ det. Ermisch SMD); France (Var) la Môle, 6.VII.1973, leg. R. Batten (3 ♂ AC), France (Aude) Quillan, 10.VII.1973, leg. R. Batten (6 ♂ and 10 ♀ AC).



Figs. 15—22. - Parameres of: 15, *Mordellistena hirtipes* (Greece, Mak., Drama); 16, *M. pseudo-hirtipes* (France, Var, Vidauban); 17, *M. purpurascens* (Italy, Sicily, Pachino); 18, *M. grisea* (Greece, Mak., Édhessa); 19, *M. micantoides* (France, Gironde, Soulac sur Mer); 20, *M. minima* (France, Var, La Môle); 21, *M. stenidea* (France, Aude, Quillan); 22, *M. perroudi* (Spain, Barcelona, Avinyo).

*Mordellistena grisea* Mulsant figs. 18, 24.

*grisea* Mulsant 1856: 376; Ermisch 1954: 175—177.

Material: France merid. leg. Puel (1 ♂ and 1 ♀ det. Ermisch 1952 SMD); Greece (Mak.) Édhezza, 6.VII.1976, leg. R. Batten (17 ♂ and 8 ♀ AC).

*Mordellistena perroudi* Mulsant figs. 22, 23.

*perroudi* Mulsant 1856: 382, 383.

Material: Turkey, Belgrader Wald, 1—7.VII.1954, leg. Schubert (1 ♂ and 1 ♀ SMD; Spain (Barcelona) Avinyo, 9.VII.1974, leg. R. Batten (1 ♂ and 1 ♀ AC).

*Mordellistena stenidea* Mulsant fig. 21.

*stenidea* Mulsant 1856: 381, 382.

Material: The type ♂ of Mulsant, labelled: „Museum Paris, 1843, coll. E. Mulsant”, red “Type”, handwritten “*Mordellistena stenidea*”. I extracted the parameres and hereby select this specimen as lectotype of *M. stenidea*; France (Aude) Quillan, 10.VII.1973, leg. R. Batten (3 ♂ AC).

*Mordellistena purpurascens* (Costa) fig. 17.

*purpurascens* Costa 1854: 17, *Mordella*; Ermisch 1954: 175—177, *Mordellistena*.

Material: Italy, Capo Circea, 27.VII.1939, leg. C.N.R. Com. Naz. Biol., parameres mounted (1 ♂ SMD); Italy (Sicily) Pachino, IV.1934, leg. Burlini (AC).

*Mordellistena micantoides* Ermisch, figs. 19, 25.

*micantoides* Ermisch 1954: 175—180.

Material: D.D.R., Kyffhäuser Gebirge Ochsenburg, 19.VIII.1959, leg. Mohr, genitalia were mounted (right paramere missing) (1 ♂ det. Ermisch 1959 SMD); Frankenhausen, 20.VIII.1963, leg. and det. Ermisch 1963 (1 ♀ SMD); France (Gironde) Soulac sur Mer, 5.VII.1974, leg. R. Batten (3 ♂ and 4 ♀ AC).

*Mordellistena balianii* Franciscolo.

*balianii* Franciscolo 1942: 77—79.

*Mordellistena ermischii* Compte

*ermischii* Compte 1966: 252, 254.

*Mordellistena wiebesi* n.sp. figs. 9—14.

Described in this paper.

Key to the West Palaearctic *micans* group species.

Note. — Ridges on hindlegs noted as follows: 3/4-3-2 means 3 or 4 ridges on tibia apart from apical one, 3 on first tarsal segment, 2 on second tarsal segment.


Length is measured without pygidium.

- 1y. Fore tibia in male more or less dilated and with fringe of hairs on inner border ..... 2
- z. Fore tibia in male not dilated nor with fringe of hairs. .... 7
- 2y. Second segment of maxillary palp in male distinctly dilated, with erect hairs on underside. General shape highly arched ..... 3
- z. This segment not dilated, lacking erect hairs on underside. General shape not highly arched ..... 4
- 3y. Segments 5—10 of antenna in male each with length/width ratio 1.5 (female slightly shorter). Terminal segment of maxillary palp elongately securiform. Head oval, much broader than long. Elytra with length/width ratio 2.7. Pygidium in male as long as elytra and two times as long as fifth sternite. Ridges 3/4-4/5-2/3. Parameres fig. 15. Length 3.3-5.5 mm. S. France, Greece ..... *hirtipes* Schilsky
- z. Segments 5—10 of antenna in male each with length/width ratio 1.25. In male terminal segment of maxillary palp broader triangular than in *hirtipes*. Head circular, almost as long as broad. Elytra with length/width ratio in male 2.5 in female 2.3. Pygidium in male 0.7 times as long as elytra and two times as long as fifth sternite. Ridges 3/4-3/5-2/3. Parameres fig. 16. Length 3.3-4.5 mm. S. Europe ..... *pseudohirtipes* Ermisch
- 4y. Maxillary palp in male reddish brown, in female brown. Antenna with basal segments reddish brown, other segments black, segments 5—10 each with length/width ratio 1.2. Fore and middle legs reddish brown, tarsus darkened distally. Pronotum broadest behind



middle, in lateral view slightly concave. Elytra with length/width ratio 2.1-2.5, finely punctate; pubescence yellowish with red-violet shine. Pygidium 2.5 times as long as fifth sternite. Ridges  $3/4-3/4-2/3$ . Parameres fig. 30. Length 3.65-4.25. Tirol, Hungary, Balkan ..... *micans* (Germar)  
 z. Maxillary palp black ..... 5

0.1mm  

Figs. 23—30. - 23, *Mordellistena perroudi* (♂ Spain, Barcelona, Avinyo); 24, *M. grisea* (♂ Greece, Mak., Édhessa); 25, *M. micantoides* (♂ France, Gironde, Soulac sur Mer); 26, *M. minima* (France, Aude, Quillan); 27, *M. perroudi* (Spain, Barcelona, Avinyo); 28, *M. fuscogemellata* (France, Pyr. Or., Canet Plage); 29, *M. gemellata* (Spain, Salamanca, Bejar); 30, *M. micans* (Turkey, Kilios-Köl.). - 23, right maxillary palp; 24, 25, eye dextrolateral; 26—29 posterior part of female eighth urosternite from dried specimen; 30, parameres.



- 5y. Pubescence of underside strikingly light and dense. Pronotum in lateral view concave, basal angles rectangular with a pointed edge. Segments 5—10 of antenna each with length/width ratio ca. 1.1—1.25. Elytra with length/width ratio in male 2.6, in female 2.4; pubescence shining red-violet. Length pygidium/fifth sternite ratio 1.6. Ridges 3-4-2. Parameres fig. 17. Length 4.5—5.0 mm. S. Italy, Sicily ..... *purpurascens* (Costa)
- z. Pubescence of underside not strikingly light and dense. .... 6
- 6x. Segments 5—10 of antenna almost square. Pronotum in lateral view almost straight, basal angles obtuse with a rounded edge. Galea long and slender, ending in a very fine point, and without appendages. Pubescence grey, sometimes slightly shining reddish to yellow-grey. Length pygidium/fifth sternite ratio 2; sometimes 2.5, then pygidium very slender at apex. Ridges 3-3/4-1/2. Phallobase long. Parameres fig. 20. Length 2.6—3.3 mm. S. France, Spain, Italy, Greece ..... *minima* (Costa)
- y. Segments 5—10 of antenna each with length/width ratio 1.25, segment 11 long oval. Galea short, inner side with appendages. Pronotum in lateral view slightly concave, basal angles rectangular with a pointed edge. Elytra almost parallel, length/width ratio ca. 2.5—2.6; pubescence grey, shining reddish golden. Length pygidium/fifth sternite ratio 2; sometimes 2.5, then pygidium very slender at apex. Ridges 4/5-4-2. Parameres fig. 22. Length 2.8—3.5 mm. S. France, Spain, Balkan ..... *perroudi* Mulsant
- z. Segments 5, 7—10 of antenna each with length/width ratio 1.35, but segment 6 shorter. Pronotum in lateral view concave, basal angles rectangular. Elytra with length/width ratio 2.2, sides in female parallel; pubescence sericeous greyish yellow. Terminal segment of maxillary palp in male triangular, in female spindle-like. Ridges 4-4-3. Length 3.5—4 mm. Spain (Ibiza) ..... *ermischi* Compte
- 7y. Elytra with length/width ratio more than 3. Head with width/length ratio ca. 2. Segments 5—10 of antenna each with length/width ratio a bit smaller than 2, these segments are triangular with a blunt inner angle. General shape elongate, posteriorly slightly dilated; pubescence sericeous, brown. Pronotum with basal angles rectangular. Length pygidium/fifth sternite ratio 2.5. Ridges 4-4-3. Length 5.0—5.6 mm. Sicily ..... *balianii* Franciscolo
- z. Elytra with length/width ratio shorter ..... 8
- 8y. Segments 5—10 of antenna each with length/width ratio in male 2, in female 1.7; nearly parallel shaped. Galea cf. fig. 12. Elytra with length/width ratio 2.5—2.7. Length pygidium/elytra ratio 0.5; length pygidium/fifth sternite ratio 2.2—2.9. Ridges 3/5-4/6-2/3. Parameres fig. 11. See figs. 9—14. Length 3.2—5.0 mm. Spain, Italy, Dalmatia, Greece *wiebesi* n.sp.
- z. Segments 5—10 of antenna shorter ..... 9
- 9y. Segments 5—10 of antenna almost square, segment 11 shortly oval. Pronotum in lateral view almost straight, basal angles obtuse with a rounded edge. Elytra with length/width ratio 2.5—2.7, sides almost parallel; pubescence light grey to yellow-grey. Pygidium conical, fifth sternite broad, length pygidium/fifth sternite ratio 2. Ridges 3/4-3/4-2. Phallobase short. Parameres fig. 21. Length 2.5—3.5 mm. S. Europe ..... *stenidea* Mulsant
- z. Segments 5—10 of antenna each with length/width ratio ca. 1.3 ..... 10
- 10y. Elytra with length/width ratio 2.5—3, from shoulders to apex evenly attenuated; pubescence grey with a light brown hue. Eyes elongately oval (fig. 24). Galea long and narrow, without appendages, apex more or less pointed. Pronotum with basal angles obtuse with a pointed edge. Ridges 3/5-3/4-2/3. Parameres fig. 18; distal part of penis slightly oval. Length 3.3—4.5 mm. S. Europe ..... *grisea* Mulsant
- z. Elytra with length/width ratio 2.3—2.5, sides almost parallel in anterior half, thence slightly attenuated; pubescence densely yellow-grey. Eyes broadly oval (fig. 25). Galea short, rounded apex with appendages. Pronotum in lateral view concave, basal angles obtuse with a rounded edge. Ridges 3/4-3/4-2. Parameres fig. 19; distal part of penis broadened, sometimes to elongately oval. Length 3.1—5.5 mm. Middle Europe, Hungary, Balkan, Russia, France ..... *micantoides* Ermisch



## ACKNOWLEDGEMENTS

For the loan of material I thank Mr. P. Teocchi, Sérignan near Orange, France, Mr. A. Descarpentries of the Muséum National d'Histoire Naturelle, Paris, Dr. R. Krause of the Staatliches Museum für Tierkunde, Dresden, Dr. J. P. Duffels of the Zoölogisch Museum van de Universiteit, Amsterdam, Mr. J. Krikken of the Rijksmuseum van Natuurlijke Historie, Leiden; Mr. C. J. M. Berger, Achel, Belgium and Mr. P. Kanaar, Leiderdorp for their gifts.

## REFERENCES

- Compte, A., 1966. Resultados de una expedición zoológica a las islas Pitiusas 2. Coleópteros. *Boln R. Soc. esp. Hist. nat. (Biol.)* 64: 239—275 (252—254).
- Costa, A., 1854. Coleotteri. - *Fauna Regno Napoli, Mordellidea*: 1—32, Pl. 19—24.
- Ermisch, K., 1954. Mordellistena Studien I. - *Beitr. Ent.* 4: 173—180.
- , 1965 - Neue Mordelliden von der Balkanhalbinsel. - *Reichenbachia* 5: 252—272.
- , 1969 Mordellidae. - *Käfer Mitteleuropas* 8: 160—196.
- Franciscolo, M., 1942. Diagnosi di nuovi Mordellidi Italiani. - *Boll. Soc. ent. ital.* 74: 76—80.
- , 1949. 13° Contributo alla conoscenza di Mordellidi. - *Memorie Soc. ent. ital.* 28: 81—95.
- Germar, E. F., 1817. *Reise nach Dalmatien und in das Gebiet von Ragusa*, 2: xii, 323 (212). Brockhaus, Leipzig (not seen).
- Méquignon, A., 1946. Contribution à l'étude des Mordellides Palaearctiques. - *Revue fr. Ent.* 13: 52—76.
- Mulsant, E., 1856. Longipèdes. - *Ann. Soc. linn. Lyon* 3: 376—385.
- Schilsky, J., 1895. *Käfer Europas* 31: 1—100.

Sprencklaan 3, 4333 HC Middelburg, the Netherlands.

---

## Ist EUROPEAN CONGRESS OF ENTOMOLOGY

Following the initiative of representatives of a number of European Entomological Societies that met in Giessen, Germany in March 1976, the Royal Entomological Society will sponsor the first European Congress of Entomology, which will be held at Reading University in Britain from 19th-22nd September 1978.

Entomologists, whether amateur or professional, are cordially invited to the Congress. Contributions concerning recent research into problems related to European entomology will be welcome, however, papers on entomological research in other areas will also be considered. It is hoped that young research workers will contribute a large part of the programme. All interested entomologists are asked to send their name and address to the above address before 31st December 1977, and they will then be sent further details of the Congress in due course.

Entomologists wishing to offer a contribution should send the title and 150 word abstract to the above address by 1st December 1977 at the latest. Contributions in English, French and German will be accepted.

It is thought unlikely that the programme will be able to accommodate all of the contributions offered, but the Planning Committee will select papers from those received by the date stated and will produce a stimulating and structured programme. Contributions will not be published, other than as preprints of extended abstracts issued to participants at registration.

It is anticipated that attendance at the Congress will cost about £ 50 for those booking full accommodation, and participating in all events and visits. There will be a reduction for family members and bona fide students. A sightseeing programme will be arranged for accompanying family members if there is sufficient interest.

Department of Zoology, The University, Early Gate, Reading RG 6 2 AT, Britain.