long as broad and as long as 3rd segment (fig. 10). Face : dark yellow. Thorax ochreous. Wings hyaline, third vein reaching costa just beyond the tip of the wing. Legs ochreous, tibiae and tarsi darker, claws all simple, slightly longer than empodium (fig. 11). Abdomen bright orange-red with a dark spot and tranverse bands. Genitalia : basal clasp segment long with a distinct basal triangular sharp process on inner surface ; distal clasp segment swollen at base, glabrous, slightly curved ; dorsal plate with deep emargination, lobes broadly rounded ; ventral plate complete, broadly rounded ; style long (fig. 12).

Holotype : Cecid. 374 ; paratypes : Cecid. 375—385 (in my collection).

F e m a l e. Length about 2 mm. Antennae : 2 + 12, 1st and 2nd flagellar segments fused, each flagellar segment elongated cylindrical, slightly constricted at centre, each bearing a distinct neck, except the terminal segment ; the neck of the 3rd flagellar segment 5 times as long as broad, neck of the 10th flagellar segment 7 times as long as broad and the 12th flagellar segment 3½ times as long as broad at its base (fig. 13). Palpi : 4 segments, 1st segment about 1½ times as long as broad, 2nd segment 2 times as long as broad, 3rd segment 2 or 2½ times as long as broad, 4th segment 3½ or 4 times as long as broad and longer than the 3rd (fig. 10). Wings spotted. Ovipositor lamelliform, lobes elliptical (fig. 14). Otherwise about as in male.

Allotype : Cecid. 386 ; paratypes : Cecid. 387—395 (in my collection).

Amsterdam, Entomologisch Laboratorium van het I.P.O., September 1952.

Araschnia levana L. Naar aanleiding van Uw artikel over de verbreiding van *Araschnia levana* L. in Ent. Ber. **14** : 216 (1953), kan ik U nog enige mededelingen doen over deze vlinder.

Op 4.V.1952 zag ik een vers ex. te Bolnes in het park van „Het Huis ten Donck".

In de laatste week van Augustus 1952 vond ik in de Roozendaalse- en Beekhuizerbossen bij Velp een groot aantal „rupsennesten". Bijna bij iedere groep brandnetels, die een beetje open stond, vond ik rupsen. Als ik zeg, dat we zonder goed zoeken een dertig nesten vonden, overdrijf ik zeker niet. Daarbij komt dan nog het feit, dat de „nesten" van kleine rupsjes zeer makkelijk over het hoofd gezien worden omdat ze aan de onderzijde der bladeren hangen, en de diertjes lang zoveel niet spinnen als bijv. *Aglais urticae*. Toch nam ik slechts één imago waar n.l. op 30.VIII, een vers mannetje. Ik nam een flink aantal rupsen mee en kweekte ze buiten, in zo natuurlijk mogelijke omgeving. Geen enkele pop leverde een ex. van de derde generatie.

In 1951 vond ik op dezelfde plaats na lang zoeken slechts één nest rupsen (24.VIII). Vóór half September was de familie (25 stuks) verpopt en op 25 September kwamen er twee vlinders uit. In September was het toen wel veel warmer weer geweest.

N. W. Elfferich, Mathenesserdijk 101a, Rotterdam (West).