

	L	K.N.		B		N.F.		Div.	Opm.
		z	n	x	xx	x	xx		
<i>Dicranum scoparium</i>								Nieuwkoop, "de Haeck"	rietland
<i>Drepanocladus aduncus</i>				x					
<i>Drepanocladus fluitans</i>					x	x			
<i>Drepanocladus intermedius</i>		x							
<i>Drepanocladus lycopodioides</i>			x			x			
<i>Eurhynchium praelongum</i>	x								
<i>Eurhynchium speciosum</i>		x							
<i>Eurhynchium stokesii</i>	x					x			
<i>Eurhynchium striatum</i>	x							Zegveldse	in wilgenbosje
<i>Fissidens adianthoides</i> c.sp.			x						
<i>Fissidens bryoides</i> c.sp.	x								tussen
<i>Fissidens exilis</i> c.sp.	x								<i>F. taxifolius</i>
<i>Fissidens taxifolius</i> c.sp.	x								
<i>Pontinalis antipyretica</i>		x				x			in sloten
<i>Punaria hygrometrica</i>	x			x					
<i>Homalia trichomanoides</i>	x								knotwilg en
<i>Homalothecium sericeum</i>	x								<i>Corylus</i> -stronk
<i>Hypnum cupressiforme</i> c.sp.		x	x			x			
<i>Isoetium myurum</i>	x								Ook op <i>Quercus</i>
<i>Leptodictyum riparium</i>	x	x					x		
<i>Leucobryum glaucum</i>					x				
<i>Anium affine</i> s.l.	x	x	x						
<i>Anium cuspidatum</i> c.sp.	x								
<i>Anium hornum</i>	x			x		x		Nieuwkoop, "de Haeck"	rietland
<i>Anium seligeri</i> (= <i>affine</i> var. <i>elatum</i>)		x							
<i>Anium undulatum</i>	x								
<i>Heckera complanata</i>	x					x			op <i>Salix</i>
<i>Orthotrichum diaphanum</i>	x								op steenhoop
<i>Plagiothecium denticulatum</i> s.l.						x		Nieuwkoop, "de Haeck"	rietland
<i>Plagiothecium roeseanum</i>	x								

	L	K.N.			B		N.P.		Div.	Opm.
		z	n	x	xx	x	xx	xx		
Plagiothecium ruthei c.sp.		x	x	x						
Plagiothecium silvaticum	x					x	x			
Plagiorth. silv. var. neglectum fo. orthocladum	x									
Pohlia nutans c.sp.						x				
Polytrichum alpestre c.sp.					x					
Polytrichum commune						x				
Polytrichum marginatum (= P. gracilis) c.sp.				x	x	x				
Pseudoscleropodium purum	x								Nieuwkoop, "de Haeck"	rietland
Rhynchostegium confertum c.sp.	x									
Rhynchostegium murale	x									
Rhytidiadelphus squarrosus	x	x	x	x						
Rhytidiadelphus triquetrus	x									
Scorpidium scorpioides		x		x						
Sphagnum acutifolium				x						
Sphagnum auriculatum				x						
Sphagnum cuspidatum				x	x					
Sphagnum fimbriatum				x	x		x		Nieuwkoop, "de Haeck"	rietland
Sphagnum palustre				x	x	x	x			
Sphagnum plumulosum				x						
Sphagnum recurvum var. amblyphyllum							x			
Sphagnum recurvum var. mucronatum							x			
Thamnum alopecurum	x									
Tortula aestiva	x									
Tortula muralis c.sp.	x									
HEPATICAE										
Calypogeia fissa				x		x				
Calypogeia trichomanis						x	x			

	L	K.N.		B		N.P.		Div.	Opm.
		z	n	x	xx	x	xx		
<i>Cephaloziella starkei</i>				x					
<i>Chiloscyphus pallescens</i>		x				x			
<i>Cladopodiella fluitans</i>		x							
<i>Prullania dilatata</i>	x								
<i>Lophocolea bidentata</i>	x					x			
<i>Lophocolea heterophylla</i>	x	x		x			x		
<i>Marchantia polymorpha</i> fo. <i>aquatica</i>		x							tussen Drep. interm.
<i>Metzgeria furcata</i>	x								op <i>Acer negunda</i> en <i>Fraxinus ornus</i>
<i>Pallavicinia lyellii</i>						x			
<i>Pellia epiphylla</i>		x	x					Nieuwkoop, "de Haeck"	rietland
<i>Pellia neesiana</i>						x			
<i>Riccardia pinguis</i> c.sp.		x							
<i>Riccia fluitans</i>						x			

Summary:

The spring field meeting was held this year in the vicinity of Woerden, province of Zuid-Holland.

An old manor was visited the first day. The surrounding park, mainly composed of deciduous trees, harboured some interesting species. *Barbula gracilis* was recorded, the second find in the Netherlands! The loamy soil was favourable for the growth of *Fissidens* and three species were collected, among which the minuscule *F. exilis*. The ash trees bore a luxuriant vegetation; its composition indicated that this was a special variant of the *Homodonteto-Isothecietum*. The next day some very old fens were examined. Their origin dates from 1494. The vegetation is peculiar, for, although a thick carpet of mosses is present, no *Sphagnum* was encountered, the water being more or less oligo-balnic. Notable finds were *Campylium elodes*, *Riccardia pinguis*, *Eurhynchium speciosum*, *Cladopodiella fluitans*, *Drepanocladus intermedius*, *Scorpidium scorpioides*. The last day was spent in the extensive lakes and very wet carr of Nieuwkoop. In an *Alnetum*, *Calypogeia fissa*, *Pellia neesiana* and the rare liverwort *Pallavicinia lyellii* were gathered.