

IV. Naamlijst

Mollusca

Abra, I - 8,9; II - 62; A.alba, I - 42; II - 31; A. antwerpiensis, I - 52; II - 49/50; A. cf fragilis, I - 48/49; A. prismatica, I - 39; Acamptogenotia, II - 10; Acirsa lanceolata, I - 52; II - 54, 57; Acmaea compressiuscula, I - 52/53; II - 41; Acrilla subreticulata, I - 52; Actaeon, I - 10; A.levi-densis, II - 10; A. semistriatus, I - 53; II - 54; A. subulatus, I - 40; A. tornatilis, I - 40; Admete fusiformis, I - 9,53; A. gracilentata, I - 40, A. wouweri, I - 40; Aligena orbicularis, I - 39; Alvania, I-40, II-53,56,60; A. antwerpiensis, II-53,85; A. beani, I-40; A. belgica, I-52, II-53; Amaea, II-49; A. woodi, II-10; Amussium woodi, I-52, II-52; Amyclina facki, II-54; A. labiosa, I-40; Anachis cf gumbeli, I-53; Anadara, II-55; A. diluvii, I-7/8, 52, II 52,55; A. suessi, I-52, II-52,55; Ancilla, I-7; A. obsoleta, I-9,53; Ancistrocyrinx corneti, I-53, II-54; Angulus benedeni, II-63; A. distortus, I-13; A. donacinus, I-40; A. fallax, II-49; A. nysti, I-32; A. posterus, II-53; Anisus, II-69; A. leucostomus, II-27, 31,33; A. vortex, II-27, 33; Anomia, I-45; II-63; A. ehippium, I-13, A. patelliformis, I-13; Aplexa hypnorum, II-26,33; Aporrhais, II-49,59; A. alata, I-9; A. pespelicani quadrifidus, I-13; A. scaldensis, I-40; Aquilofusus festivus, I-9; Arca diluvii, I-36; A. tetragona, I-39; Architectonica, II-53,56; A. berthae, II-56; Arco-pagia, II-10; A. balaustina, II-53; A. crassa, I-40, II-10; A. heberti, I-48/49; Arcoperna sericea, I - 52, II-10,52; Armiger crista, II-27; A. crista cristata, II-33; A. c. nautilus, II-33; Astarte, II-10,62; A. angulata, I-9, 39, II 10,52,55; A. basteroti, I-37,39, II - 52; A. beyschlagi, I-52, II-52; A. concentrica, I-52; A. corbuloides, II-10; A. galeotti, II-10; A. goldfussi, I-8,9, II-40,52; A. gracilis, II-52; A. henckeliusiana, I-20/21, A. incerta, I-39; A. mutabilis, II-10,52,55; A. obliquata, I-39, A. omalii, I-37,39; A. parva, I-39; A. parvula, II-10; A. pseudopygmaea, I-39, A. radiata, I-9,52, II-40,52; A. triangularis, I-39, II-55; A. trigonella, I-20; Asthenotoma ornata, I-10,40; Astraea belgica, I-51/52, II 39/40, 53, 60; Athleta ficulina, I-9, A. rathieri, I-32; Aturia, II-11.

Babylonella fusiformis, II-54; Balcis eichwaldi, I-52; Barnea candida, I-13; B. cylindrica, I-42; Bathyarca pectunculoides, I-9,39; Bathytoma, I-8, B. cataphracta, I-10, B. crenata, I-20; Bela neerlandica, I-40; Belosepia sepicidea, II-67; Bittium reticulatum, I-13, II-26,32; B. robustum, I-55; B. rubanocinctum, I-40, B. spina, I-9; Bithynia tentaculata, II-26,33; Boreodrillia toftlundensis, I-9,53; Bornia deltoideum, I-39; Borsonia, II-49; B. deluci, I-20; B. unipliata, I-53; Brachitoma grateloupi, I-53, B. obtusangula, I-53; Brachyodontes lineatus, I-13; Brocchinia mitraeformis I-40,53; Buccinum undatum, I-13.

Cadulus, II-71; C. gadus, II-81; C. politus, I-40, 52/53, II-50, 53, 81; Caecum cf glabrum, II-10; C. mamillatum, I-40; Callista chione, I-39; Calliostoma, II-40,50, 53, 60, 62; C. simile, I-40; C. zizyphinum, I-40; C. z. simile, I-42; Calopodium inaequalis pinna, II-10; Calyptraea, I-45; II-49; C. chinensis, I-9,13,40,52; C. striatella, I-20,21,32; Cancellaria, II-40;

C. cf aperta, I-53; C. beyrichi, I-53; II-54; C. varicosa, I-9; Capulaomea radiata, I-40; Capulus ungaricus, I-40; II-54; Cardita, I-39; C. chamaeformis, I-9,52, II-52; C. corbis, I-39; C. omaliana, I-20/21, C. orbicularis, I-7/8, C. planicosta, II-69, C. squamulosa, II-52; Carditopsis chavani, II-52; Cardium, II-53, 56,59,68,70; C. edule, I-13, II-21,30,67; C.e.belgicum, I-42, C. exiguum, I-13; C. nodosum, I-39; C. parkinsoni, I-43; C. paucicostatum, I-13; C. straeleni, I-9,52, II-53,56; C. tuberculatum, I-13, Cardiomya costellata, II-41; Cavilucina, II-49; C. droueti, I-52, II-40; Cerithiella, II-54,61; C. asperulata, II-54; C. degrangei, II-54; Cerithiopsis, II-59; C. tubercularis, I-13,40; II 54,56; C. subulata, II-56; Cerithium kunthi, II-6; Chama gryphoides, II-53; Chlamys, II-52,55; C. angeloni, I-52,II-52,58,60, C. clavata, II-62; C. edegemensis, II-52,55; C. ercolaniana, I-52,II-52,59; C. gerardi, I-39, II-14,63; C. harmeri, I-40, 42; C. hoeninghausi, I-19,21,32; C. lilli, I-9,50,52; II-55; C. multistriata harmeri, II-10; C. opercularis, I-39,42, II-63; C. radians, I-39,52, II-52,55; C. sarmenticius, II-55; C. similis, I-39, II-52,59,60; C. tigerina, I-39, II-52,55,62; C. woodi, I-8/9, Chrysallida, I-40; C. pygmaea belgica, II-54, C. indistincta, I-40, C. jeffreysi, I-40; Chiton, I-52,53; Chitonidae, II-53,56; Cingula, I-52,53, II-53,56; C. inusitata, I-40; C. cf proxima, II-10; C. p. laevigata, I-52, II-53, Circulus hennei, I-52, II-10,53; C. praecedens, I-40,52, II-53; C. striatus, I-40; C. supranitidus, II-10; Cirsotrema crassicosatum, II-54; C. fimbriosum, I-40; Clavus diensti, I-53; Cochlicopa, II-70; Cocculina miocaenica, II-53; Columella, II-70; C. columella, I-46, II-27,32,33; Congeria nysti, I-20,48; Comus antediluvianus, I-10; C. dujardini, I-10,53,II-54; Cor alliophaga, II-40,51; C. lithophagella, I-51,52,54, II-40,52,55,58/9; Corbicula fluminalis, II-14,20,28,31; C. semistriata, I-18,20,22,48; Corbula, II-50; C. gibba, I-20,40,42,48,52, II-31,32; Crenella rhombea I-39, II-40,52; Cultellus cf pellucidus, I-52/3; Cuspidaria costellata, I-9,52, II-53; C. jugosa, I-40; C. rostrata, I-40; Cyclocardia orbicularis, II-10; C. scalaris, II-10, Cyclostrema, I-40; Cylichna cylindracea, I-10,40,53, II-54; Cypraea, I-52, Cyprina, I-9,39, II-62; C. islandica, I-36,52, II-62/3; C. i. rotundata, I-32; C. rotundata, I-20-21; C. rustica defrancei, I-43; C. rustica, I-37, II-62,63; Cyrtodaria angusta, I-42.

Dentalium, I-7/9,32,40,45,52, II-71, 78,80,81; D. (Fustiaria), II-81, D. (Laovidentalium), II-80; D. cf anomalocostatum, II-77; D. badense, II-72/4; D. bouei, II-75; D. costatum, II-53,76; D. dollfusi, II-72/4. D. dumasi, II-75; D. entale, II-50;53,56,79; D. fissura, II-81; D. cf kickxi, I-20,21; D. michelotti, II-75; D. novemcostatum f. mutabile II-74,76; D. sandbergeri, I-20; D. sexangulum, II-75,77; D. vitreum, I-52, II-79,80; D. vulgare, II-79/80; D. vulgaris, I-13; Delphinula speyeri, II-6,7; Deroceras, II-28,33; Digitaria digitaria, I-39; D. excurrens, I-39; Diodora apertura, I-40; Diplozonta, II-10,50,52; D. rotundata, I-39; D. trigonula astartea, I-39; Divaricella, I-20; D. divaricata, I-13; Dolium, I-8; Donax variegata, I-39, II-10; Dorsanum subpolitum, I-9; Dosinia exoleta, I-39,42; D. lupinus lineta, I-13.

Emarginula, II-39,53,56,59,60; E. cancellata, II-53,56,60; E. crassa, I-40, E. nysti, I-32; E. reticulata, I-40, Ensis, I-52,53,56; E. ensis, I-13; E. siliqua, II-10; Eocypraca, II-54; Eomiltha, II-49; E. transversa, I-52, Epitonium, II-50,54,57; E. clathrus, I-13; E. frondiculum, II-54,57; E.f.antwerpiense, I-52; E. weyersi, I-52; Erato, I-40, E. germanica, II-41,54; Erycina, II-53,55, E. cimbrica, II-52; E. degrangei, II-52; E. depressa, I-39; E. striatissima, II-52; Eulima, I-52; Eulimella, I-53, II-50,54,57; E. acicula, I-10,40, II-57; E. neumayeri, I-10, II-57; Exilia ordita, I-9.

Fusiturris duchasteli, I-53; Fusus sexcostatus, I-9.

Galba palustris, II-26,33; G. truncatula, II-26,33; Gari depressa, I-13, Gastrana fragilis, I-13; G. laminosa, I-40; Gastrochoena dubia, II-40,58; Gemmula, I-53, II,49; G. stoffelsi, I-53; G. zimmermanni, I-53; Gibbula, I-40, II-62,63; G. cineraria, I-13; G. gelriana I-55; G. magus, I-13; G. octosulcata, I-42; G. solarium, I-55; Glibertia prosperi, I-39; Glycymeris, I-36; G. glycymeris variabilis, I-39; G. lunulata, I-19,21,32; G. obovata, I-19,21,32; G. pilosa deshayesi, II-52; G. variabilis, I-42, II-62; G. v. deshayesi, I-52; Gouldia minima, I-9,39, II-53; Gregariella barbatella, II-52,55; Gyraulus acronicus, II-27,33.

Haedropleura septangularis, I-13; Hemilepton kautskyi, I-39, Heteranomia squamula, I-13,39,42,52; Hiatella, II-40,51; H. arctica, I-13,48,49,51,52,54, II-40,50,58; Hinia bocholtensis, II-54; H. dollfusi, I-40; H. cf. ligustica, I-40; H. tenuistriata, II-54, Hydrobia antwerpiensis, II-53,56; H. duboissoni, I-48; H. elongata, I-22; H. stagnorum, I-13; H. ulvae I-13.

Inquisitor borealis, I-53; I. obeliscus, I-10; Isocardia, II-49; I. humana, I-39; I. lunulata, I-7/9,50,52, II-41; I. subtransversa, I-20,32.

Jagonia decorata, II-10; Jupiteria pygmaea, I-8,39,52.

Kellia, II-40,53,55,58,60; K. suborbicularis, II-55; Kleinella nordmanni, II-54,57.

Lacuna, I-56, II-53,56; Laevicardium, I-39; L. cingulatum I-21; L. cyprium, I-8/9, II-49; L. decorticatum, I-37,39,42; L. dingdense, I-9, II-53; L. parkinsoni, I-42,55; L. subturgidum, I-52; Laiocochlis sinistratum, II-10; L. woodi, I-40; Lajonkairia rupestris lupinoides, II-10; Lathyrus rothi, I-9; Leda deshayesiana, II-11; Leiostraca glabra, I-40; Lentidium complanatum, I-41,42; L. donaciformis, I-20,48,49; L. nysti, I-18,22; L. triangula, I-20,48; Lepeta scaldensis II-10; Lepton transversarium, I-52, II-41,53; Lima, II-10,55; L. loscombi, I-39, II-52; L. strigillata, II-52; L. subauriculata, II-52,55; Limatula ovata, I-39; L. subauriculata, I-39; Limopsis, I-7; L. aromala, I-8/9,39,52; II-48,50,52,59,60; L. aurita, I-8/9,52, II-52,48,60; L. goldfussi, I-19,21; L. lamellata, I-9; Liomesus dalei, I-40; Littorina littorea, I-13; L. saxatilis, I-13, II-25; Lucina orenulata, I-39; L. thierensi, I-48; Lucinoma, II-49; L. borealis, I-39,52, II-40,59; Lutetia nitida, II-53/6; Lutraria lutraria, I-13; Lymnaea, I-18,22,49; II-69,70,85; L. stagnalis, II-26,33; Lymnaeidae, II-26,33; Lyrotyphis priscus, II-54.

Macoma, II-49; M. balthica, I-13,II-31; M. elliptica, I-52; M. obliqua, I-40, M. praetenuis, I-41,42,55; Mactra corallina plistoncerlandica, I-13; Mangelia, I-40, II-50; M. costellaria, I-49; M. elegans, I-49; M. smithi, I-40; Margarites belli, II-10; M. trochoideus, II-10; Martesia, II-40,51; M. rugosa, I-51,52,54, II-40,53,58,60; Melampus pyramidalis, I-43,55,56; Melanella, I-9; M. alba, I-40; M. eichwaldi, II-54; Melania acuta, I-22; Melanoides fasciatus I-22,48; Microdrillia crispata, I-9; Mitra, II-54; Mitrella scaldensis, I-40; Modiolus cf. faujasi, I-20,48,49; M. modiolus, I-38,II-10; M. phaseolinus, I-39,38; Montacuta, II-53,56; M. antwerpiensis, II-53,55; M. coarctata, I-39; M. ferruginosa, I-39; M. substriata, II-10; Murex inornata, I-9; M. nysti, I-52, II-41; M. scalariformis, I-52; Musculus, I-39, II-52,55; M. barbatella, II-40; M. nysti, I-48; Mya arenaria, I-13,43; M. truncata, I-13,40; Mysella, II-40,53,55,56; M. bidentata, I-39,II-56; Mytilus edulis, I-13,39,42, II-28; M. subfragilis, I-20,48,49.

Narona mitraeformis parvula, II-54; N. varicosa, II-54; Nassarius, I-9; N. bocholtensis, I-9; N. cf. facki, I-53; N. holsaticus, I-9; N. pygmaeus, I-13; N. reticulatus, I-13, II-26; N. cf. tenuistriatus, I-53; Natica, I-52, II-50; N. achatensis, I-48,49,50; N. catena achatensis, II-6; N. fissurata, II-6; N. multipunctata, I-40; Nautilus lamarcki, II-67; Neptunea contraria, I-42; Neurina duchastoli, I-22,48, II-8,85; Newtoniella genei, I-9,52,53; Niso acarinatoconica, I-52; Northia gossardi, I-21,48; Nucella lapillis incrassatus,

Nucella lapillus incrassatus, I-43; Nucula, I-52, II-28; N. compta, I-32; N. nucleus, I-13, 32, 39, II-50, 52; Nuculana pygmaea, II-40, 49; N. westendorpi, I-8, 32, 52; II-48, 50, 52, 59, 60; Nuculoma, I-8; N. haesendoncki, I-52; N. laevigata, I-39; Nystia, I-48; N. duchasteli, I-48, N. plicata, I-18, 22; N. pseudoplicata, I-22.

Ocenebra erinaceus, I-13; Odostomia, I-49, II-54, 57; O. conoidea, I-40, 53, II-50, 54; O. semperi, I-49; Oliva dufresnii, I-53; Opalia, II-57; O. pertusa, I-52; II-54; O. vilandti, I-9; Orbitolites, II-67, 70; Ostrea, I-37, II-14, 52, 67; O. callifera, I-45; O. cyathula, I-22; O. cymbula, II-68; O. edulis, I-13, 39, 42; O. gryphina, II-67.

Pandora, II-63; Panopea, I-36, II-50, 51; P. faujasi, I-42; P. menardi, I-50, 52, II-41; Paphia rhomboides, I-42; Parvicardium scobinula, I-20, 22; P. s. tongrica, I-48; Pecten, I-52; P. pictus, I-32; P. plebeius, II-67; P. subgrandis, I-39; P. westendorpianus, II-62; Phacoides, I-20; P. borealis, I-9, 50; Phalium bicoronatum, I-52, II-41; P. miolaevigatum, I-52; Phasianema, II-10; P. sulcatum, I-40, II-10; Phaxas pellucidus, I-40; II-53; Philbertia praehispida, II-54; Philine quadrata, I-40; P. scabra, I-40; Pinna, I-37; Pirenella monilifera, I-22, 48; Pirula concinna, I-21; P. condita, I-52; Pisidium, I-13; P. amnicum, I-13; II-29, 31; P. casertanum, II-30; P. henslowanum, II-29; P. hibernicum, II-30, 32; P. lilljeborgi, II-30; 32; P. milium, II-29; P. nitidum, II-30; P. obtusale, II-30; P. o. lapponicum, I-46, II-32; P. subtruncatum, II-29; P. supinum, II-29, 32; P. vincentianum, II-30, 32; Pitar chione, II-53; P. polytropa, I-50; P. p. nysti, II-40, 60; P. rudis, I-39, 52, II-53; Pitaria incrassata, I-20, 21, 48, 50; P. kickxi, I-20; P. polytropa, I-36; P. p. subrotundata, I-32; P. splendida, I-20; Planorbis, I-9; P. acuticarinatus, I-18, 22; P. bosqueti, I-18, 49; P. planorbis, II-27, 31, 33; Plejona rathieri, I-21; Pleurodon cinctus, I-32; P. ovalis, I-39; Polinices, I-9, 40, II-50, 54; P. catena achatensis, I-50; P. c. helicina, I-52; P. cyclostomoides transferta, I-52; P. olla, II-54; P. polianus, I-13; P. submamillaris dertomamilla, I-52; P. varians protracta, I-52; Pomatias fragilis, I-48; Potamides lamarecki, I-22, 48; Psammobia, I-39, II-10, 53, 56; P. affinis, II-56; P. costulata, II-10; P. fervensis, I-39; P. tellinella, I-39; Pseudamnicola helicella, I-18, 22; Ptychopotamides thenensis, I-22; Pterynotus, II-49; Punctum, II-70; Pupilla, II-69/70; P. muscorum, II-27, 33; Pycnodonta callifera, I-20, 32; P. cochlear, I-8; Pyramidella laeviuscula, I-40; P. plicosa, I-53, II-54; Pyrrula condita, I-9.

Raphitoma, I-40; Retusa elongata, II-50, 54, 60; R. e. conuloidea, I-40; R. subcylindrica, I-53; Rhizorus acuminatus, I-10, 53, II-54; Ringicula, II-40; R. buccinea, I-8, 10, 40, 53, II-50, 54; R. ventricosa, I-40, II-54; Rissoa membranacea, I-13; R. obsoleta, I-40; R. parva, I-40; R. turbinata, I-48.

Sabatia utricula, I-40, 53; Sandbergeria cancellata, I-48; Sanguinolaria, I-48, 49; S. brabantina, I-22; Saxicava arctica, I-40; Saxicavella carinata, I-40; S. jeffreysi, I-40, 52, II-50, 53; S. pusilla, II-53; Saxolucina omaliusi, I-22, 48; S. tenuistriata, I-20; Scacchia, II-55; S. elliptica, I-39; Scala clathratula minuta, I-40; S. frondicula, I-40; S. subulata, I-40; Scaphander, II-62; S. lignarius, I-40, 53; Scaphella bolli, I-8, 36; S. lamberti, I-40; Scrobicularia plana, I-13, 41, II-31, 67; Scurria compressiuscula, II-53; Seila angusta, II-6; S. spicula, II-6; S. trilincata, I-40; II-6; Semicassis, I-36; S. bicoronata, I-9; Sipho gregarius, I-40; Sigatica hantoniensis, I-32; Solaricella, I-52; S. maculata, I-40; Solecurtis basteroti, II-53; Solen marginatus, I-13; Solenocurtis basteroti, I-52; Spaniorinus ambiguus, I-39; Sphenia, I-50; S. angustata, I-48; Sphaerium corneum, II-29; S. solidum, I-13, II-29, 31; Spiratella atlanta, II-85; S. microstralis, I-53, II-54, 57; S. umbilicata, II-7; Spisula arcuata, I-42; S. elliptica, I-9; S. ovalis, I-39; S. triangula, I-52; S. subtruncata, I-13; S. s. triangula, II-53; Splendrillia selenkae, I-9;

Splendrillia selenkae, I-9; Stenothyra bidens, I-22; S. pupa, I-48; Streptochetus, II-49, 51, 59; S. sexcostatus, I-53; Striarca lactea, II-39, 40, 52, 58; Strombiformis tauriniensis, II-54; Succinea, II-28, 69, 70; S. elegans, II-28; S. e. schumacheri, II-32; S. oblonga, I-13, II-28, 33; Sulcotrivia testudinella, I-55; Syrnola, I-22; S. laevis, I-49.

Taras trigonulus astartea, I-42; Teinostoma decussatum, I-48; Tellimya suborbicularis, II-10; Tellina benedeni, I-43; Terebra hoernesii, I-10, 53; T. inversa, I-40; Teretia anceps, II-54; Theodoxus fluviatilis, II-25, 31, 33; Theridium intradentatum, I-22; Timoclea ovata, I-39; Thracia, I-40; II-50; T. ventricosa, I-9, 52; Thyasira, II-10; T. flexuosa, I-39, 52; II-52; Tornus, II-53, 56, 59; T. belgicus, I-40; Trichia, II-70; T. hispida, II-28, 33; Trigonostoma protrigonostoma, II-54; T. umb. pluricostatum, II-54; Triphora adversa, I-13; T. fritschi, II-54; T. perversa, II-54; T. p. adversa, I-40; Trivia, I-13, 40; Trophonopsis muricata, I-40; Turbonilla, I-10, 53, II-50, 54, 57; T. amoena, II-57; T. densecostata borealis, II-57; T. filosa, I-40; T. cf. gastaldi, I-53, II-57; T. internodula, I-40; T. kendalli, I-40; T. cf. miomutinensis, II-57; T. miostriatula, II-57; T. m. edegemensis, I-53; T. pseudoterebralis, I-53, II-57; T. senistriata, I-40; T. undulata, II-57; Turricula, I-32; T. steinvorthei, I-9, 53; Turris, I-9, II-49, T. antwerpiensis, I-40; T. duchasteli, I-9; T. zimmermanni, I-9; Turritella, II-14, 67, 68; T. incrassata, I-40; T. subangulata, I-9, 52; Tympanotonus labyrinthus, I-22, 48; Typhis fistulosus, I-9, 52; T. pungens, II-54.

Vallonia, II-70; Valvata piscinalis, I-13, II-25, 33; V. p. alpestris, II-33; V. pulchella, II-25, 27, 32; Valvatina umbilicata, II-7; Varicorbula gibba, I-8, 9, 13; Venerupis aurea senescens, I-13, II-30, 32; V. decussata, I-13; V. rhomboides, I-39; Venus, II-49, 59; V. casina, I-39, II-63; V. imbricata, I-39; V. multilamella, I-7, 8, 9, 52, II-40; V. ovata, I-13; Vermetus ingens, II-53; V. intortus, I-40; Vertigo, II-69, 70; V. parcedentata, II-27, 32, 33; Voxillum aciculum, I-53, II-54; V. a. proccra, I-53; Villorita neglecta, I-20, 48; Volvula acuminata, I-40.

Xenophora II-49, 63; X. deshayesi, I-9, 52.

Yoldia glaberrima, I-8, 52, II-49; Y. semistriata, I-39.

Zirfaea crispata, I-13.

BRACHIOPODA

Lingula, II-63; L. dumortieri, II-63, Terebratulina, I-45, II-62, Terebratulina, II-10.

ECHINODERMATA

Echinocardium, II-8.

VERMES

Ditrupa, II-62, 63; D. strangulata, II-67.

FORAMINIFERA

Nummulites, II-67; N. planulatus, II-68

FACIES

Alnus, II-24; Betula, II-24; Corylus avellana, II-24; Cytisus, II-24; Pinus, II-24; Quercus robur, II-24; Sphagnum, II-24.

PISCES

Galeocerda aduncus, I-8; Notidanus cf. primigenius, I-8; Odontaspis acutissima, I-7; O. vorax, I-7; Oxyrhina hastalis, I-8.