

# Sharks, bony fishes and endodontal borings from the Miocene Montpelier Formation (White Limestone Group) of Jamaica

Charlie J. Underwood<sup>1</sup> & Simon F. Mitchell<sup>2</sup>

<sup>1</sup>*School of Earth Sciences, Birkbeck College, Malet Street, London WC1E 7HX, England;*  
*e-mail: c.underwood@geology.bbk.ac.uk*

<sup>2</sup>*Department of Geography & Geology, University of the West Indies, Mona, Kingston 7, Jamaica;*  
*e-mail: smitchell@cwjamaica.com*

Received 12 March 2001; revised version accepted 7 April 2003

Bulk samples of Miocene carbonate sedimentary rocks (deep-water chalks and shallow-water-derived calcarenites) from the Montpelier Formation (White Limestone Group) at Duncans Quarry (Jamaica), have yielded a small, but diverse, fauna of dissociated fish remains. Shark remains include the teeth of five species, four of which are squalids. A diverse, but taxonomically indeterminate, osteichthyan tooth assemblage is also present. A number of the teeth contain microborings of two ichnotaxa. The assemblage is considered to be typical of a deep-water continental slope fauna and indicates an abrupt northern margin of the Miocene shallow-water carbonate platforms.

KEY WORDS: Montpelier Formation, White Limestone Group, fish faunas, Miocene, Jamaica.

---

## Introduction

Although fish comprise very important elements of the vast majority of marine ecosystems, their study has typically been extremely piecemeal. The bulk of well-described fossil fish assemblages comprise either unusually rich concentrations of chondrichthyan teeth (often at hiatal levels) or exceptionally well-preserved assemblages of bony fish preserved as largely complete skeletons (often within atypical depositional settings). Despite these biases within the described fossil fish record, isolated fish remains, especially teeth, are present within many fossiliferous marine sedimentary rocks and represent a significant component of fossil biotas.

Deep-water fish assemblages are poorly known and have rarely been described in detail. This is especially true of bony fish faunas, where little work has been carried out on the tooth morphologies present within the different taxonomic groups. Deep-water shark assemblages are known from several levels within the Neogene, such as the Miocene of southern France (Ledoux, 1972) and the Pliocene of northern Italy (Cigala-Fulgosi, 1986). These assemblages are typically dominated by diverse Squaliformes and deep-water Lamniformes, along with some pelagic taxa.

In this paper, a deep-water fish assemblage from the Miocene limestones of the Montpelier Formation (White

Limestone Group) at Duncans Quarry is described. All specimens carry requisition numbers of the Florida Museum of Natural History, University of Florida, Gainesville (UF).


## Previous work and geological setting

There has been relatively little work on the Neogene fish assemblages from Jamaica. Eyles (1958) recorded abundant small fish teeth from a phosphate band at the base of the bauxite deposits that rest on the shallow-water limestones of the White Limestone Group. Purdy *et al.* (1996) recorded a shark tooth and other fragments from the Pleistocene of Jamaica. Stringer (1998) described teleostean otoliths from the Upper Pliocene Bowden shell bed. Donovan & Gunter (2002) described several large teeth found in the Geology Museum at the University of the West Indies, but without locality or stratigraphic data. Therefore, the presence of a small, but diverse, assemblage of fish teeth from the Montpelier Formation adds considerably to this dataset.

The Montpelier Formation (White Limestone Group) of Jamaica consists of deep-water chalks and calcarenites that contain bands of nodular chert and bentonite clay bands (Mitchell, 2004). The Montpelier Formation is well exposed at Duncans Quarry, which is

situated on the north coast of Jamaica some 5 km west of Duncans (Figure 1). About 50 m of the Montpelier Formation is exposed in the quarry. The lower part consists of deep-water chalks with chert bands and thin bentonites, and the upper part, deep-water chalks with abundant graded calcarenite beds. The graded calcarenites contain sand-grade carbonate that includes faunal elements, such as larger benthic foraminifera,

derived from the nearby platform carbonate successions of the Clarendon Block (Figure 1). The deep-water sedimentary rocks at Duncans Quarry are Miocene (Steineck, 1974; Donovan *et al.*, 1995). The fish assemblages described in this paper were picked from bulk samples collected by Roger Portell (Florida Museum of Natural History, Gainesville) from the upper part of the succession at Duncans.



**Figure 1.** Location of Duncans Quarry and its relationship to the shallow-water platforms in the Miocene.

### Systematic palaeontology

Class Chondrichthyes Huxley, 1880  
Order Squaliformes Goodrich, 1909

*Remarks* — It is possible that the order Squaliformes as defined by Compagno (1973) is paraphyletic (Shirai, 1996).

Genus *Deania* Jordan & Snyder, 1902

*Type species* — *Acanthidium calceum* Lowe, 1839; Recent.

*Deania* sp.  
Figure 2/1, 2

1972 *Deania calceus* (Lowe) — Ledoux, pp. 149-153, figs 7, 8.


*Material* — One lower tooth, UF 206535.

*Description* — The tooth is somewhat higher than wide and strongly compressed. A large principal cusp is directed towards the posterior, the apical part overhanging a small distal heel. An elongate, but narrow, labial apron reaches almost to the base of the root, and has large foramina on its anterior and posterior side. The lingual face of the root has a pair of well-developed foramina above a narrow antero-basally directed groove. A large indentation is present on the posterior part of the root lingual face. A strongly compressed flange projects from the anterior side of the root.

*Comparisons* — This single lower tooth appears almost identical to teeth described as *Deania calceus* [*sic*] by Ledoux (1972). These teeth differ from those of extant *D. calcea* figured by Herman *et al.* (1989) in having a deeper root, less erect cusp with a more convex cutting edge and narrower labial apron. These teeth can be distinguished from those of *Centrophorus* in having clearly separated lingual foramina and no lingual protuberance.

### Figure 2.

- 1, 2 - *Deania* sp., UF 206535, in labial and lingual view, respectively.  
3-9 - *Squaliolus schaubi* (Casier, 1958), UF 206537 in labial (3) and lingual (4) view; UF 206538 in labial (5) and lingual (6) view; UF 206539 in labial (7) and lingual (8) view; UF 206464, in labial (9) view.  
10, 11 - *Squaliolus* sp., UF 206536, in labial and lingual view, respectively.  
12-17 - *Scymnodon* aff. *obscurus* (Vaillant, 1888), UF 206534 in labial (12), lingual (13) and lateral (14) view; UF 206553, in labial (15), lateral (16) and lingual (17) view.



Genus *Squaliolus* Smith & Radcliffe, 1912

*Type species* — *Squaliolus laticaudus* Smith & Radcliffe, 1912; Recent.

*Squaliolus schaubi* (Casier, 1958)

Figure 2/3-9

1958 *Centroscymnus schaubi* Casier, pl. 1, fig. 11.

1972 *Squaliolus schaubi* (Casier) — Ledoux, fig. 11.

*Material* — One partial and three near-complete lower teeth, UF 206537, 206538, 206539 and 206464.

*Description* — Although all of the teeth recorded of this taxon are from the lower dentition, there is some degree of variation due to differing positions on the jaw. The teeth are considerably higher than wide and very strongly compressed. The principal cusp is very low and short, with a strongly convex anterior edge. This cusp completely overhangs the small distal heel. The lingual face of the tooth is flat, but with a recessed posterior section below the distal heel. A very well-developed foramen is roughly in the centre of the lingual face, immediately below the lower edge of the enameloid crown. The labial face of the tooth has an extensive enameloid covering, other than within a recessed region comprising approximately the anterior third of the tooth, the upper edge of which forms a strongly arcuate anterobasal edge of the crown. A very well-developed foramen in the lower part of the labial face forms the upper limit of a distinct groove, which in turn forms a notch on the root basal edge. This foramen and groove are separated from the anterior recess by a narrow ridge of enameloid. Several small foramina are present posterior to the labial groove.

*Comparisons* — These teeth appear almost indistinguishable from lower teeth of *S. schaubi* figured by Ledoux (1972), although no associated upper teeth or more erect (?male) lower teeth, as noted by Ledoux (1972), were recorded in this study. These teeth may be separated from the extant Atlantic species of *Squaliolus*, *S. laticaudus* (Smith & Radcliffe) as figured by Herman *et al.* (1989), by the lower and shorter principal cusp and more rectangular overall shape. Teeth of this taxon may be separated from those of the closely related genus *Euprotomicrus* Gill by the presence of a better-developed labial anterior recess and a narrower labial groove.

*Squaliolus* sp.

Figure 2/10, 11

*Material* — One lower tooth, UF 206536.

*Description* — This single lower tooth is very strongly

compressed and has a basal edge at an angle to the anterior and posterior edges of the tooth, giving an overall rhombic shape with a distally directed root. There is a large and erect principal cusp with a straight anterior edge, and a very small distal heel. The flat lingual face of the root has a very large central foramen and a posterior recessed region, which is somewhat narrower than the distal heel above it. The labial face of the tooth has narrow anterior recessed region, the upper edge of which forms a strongly notched anterobasal edge of the crown. A very well-developed foramen in the lower part of the labial face opens into a strong groove. This foramen and groove are separated from the anterior recess by a narrow ridge of enameloid. Two small, oval foramina are present posterior to the labial groove.

*Comparisons* — This tooth is larger than any known lower tooth of *S. schaubi*, and differs in having a far larger and more erect cusp. The narrow labial groove suggests an affinity with *Squaliolus* rather than *Euprotomicrus* or *Heteroscymnoides* Fowler.

Genus *Scymnodon* du Bocage & de Capello, 1864


*Type species* — *Scymnodon ringens* du Bocage & de Capello, 1864; Recent.

*Scymnodon* aff. *obscurus* (Vaillant, 1888)

Figure 2/12-17

*Material* — Two upper teeth, one partial and one complete, UF 206534 and 206553.

*Description* — These teeth comprise a larger, if somewhat damaged, anterior tooth and a smaller lateral tooth. In the lateral tooth (where the crown is well-preserved), the single cusp is elongate, about four times as long as wide, and slightly posteriorly inclined. Cutting edges are present along both sides of the cusp, except for within a slightly constricted 'collar' at the base of the cusp. This cutting edge separates a fairly flat labial face from a strongly convex lingual face. The preserved part of the cusp of the anterior tooth is similar, differing only in the possession of two faint ridges on the basal part of the labial face and lacking posterior inclination. The root of both teeth is strongly bilobate, with a distinct notch on the basal edge. The lingual face is flat and has a pair of well-developed foramina, one at the base of the cusp and one within the basal notch. The labial edge is more convex with several small marginal foramina. The root lobes of the anterior tooth are similar in size and oval in lingual view. There is strong asymmetry of the root lobes of the lateral tooth, with an anterior pointed lobe and a posterior lobe of almost rectangular lingual profile.



**Figure 3.**

- 1, 2 - Squaliform? indet., UF 206474, in oblique and surface view, respectively.  
 3 - *Carcharhinus* sp., UF 206554, labial? view.  
 4 - Tooth type 1, UF 206540.  
 5 - Tooth type 2, UF 206541.  
 6 - Tooth type 3, UF 206478.  
 7 - Tooth type 4, UF 206477.  
 8 - Tooth type 5, UF 206542.  
 9 - Tooth type 6, UF 206480.  
 10 - Tooth type 7, UF 206476.  
 11, 12 - Tooth type 8; UF 206475, lateral view of oval tooth (11); UF 206479, occlusal view of rounded tooth (12).  
 13 - Circular boring, UF 206534.  
 14, 15 - *Mycelites ossifragus* Roux, 1887; UF 206535, large and small ramifying mass, respectively.

*Comparisons* — The upper teeth of heterodont squaliforms are very variable in form between species and often bear little resemblance to the lower teeth of the same taxa. The teeth recorded here appear extremely

similar to upper teeth of *Scymnodon obscurus* (Vaillant, 1888) figured by Herman *et al.* (1993). The strongly bilobed root and rather oval root lobes separate upper teeth of this taxon from those of other species of *Scymnodon* (Herman *et al.*, 1989, 1993). These teeth can be separated from the rather similar upper teeth of *Squaliolus*, *Euprotomicrus* and *Heteroscymnoides* by the presence of a constriction at the base of the cusp, a pronounced notch between the root lobes and lack of a deep excavation at the base of the cusp on the labial face. Therefore, these teeth do not appear to be conspecific with any of the lower teeth recorded in this study, this not being unsurprising considering the low total number of specimens.

Squaliform? indet.  
Figure 3/1, 2

*Material* — One scale, UF 206474.

*Description* — The roughly diamond-shaped face of this scale is ornamented with several longitudinal ridges. The scale is low with no well-defined neck, but the somewhat concave lateral faces are ornamented with a very weak reticulate pattern.

*Comparisons* — Selachian scales are generally regarded as poorly diagnostic, but the scale recorded here is of typical squalid form, being similar to scales of many Recent and fossil heterodont squalids (*e.g.*, Cappetta, 1980; Ledoux, 1972).

Order Carcharhiniiformes Compagno, 1973  
Genus *Carcharhinus* de Blainville, 1816

*Type species* — *Carcharias melanopterus* Quoy & Gaimard, 1825; Recent.

*Carcharhinus* sp.  
Figure 3/3

*Material* — One partial upper tooth, UF 206554.

*Description* — This single isolated cusp is triangular, being rather higher than wide, and robust. Both lingual and labial faces are convex and unornamented, being separated by a well-developed cutting edge. The cutting edge is lightly and irregularly serrated other than at the extreme apex.

*Comparisons* — The robust and serrated shape of the cusp suggests that this specimen represents an upper tooth. Although there is great variation in the dentition of various species of *Carcharhinus*, identification of different species is extremely difficult even when deal-

ing with intact teeth (*e.g.*, Purdy *et al.*, 2001). This specimen must therefore be regarded as indeterminate.

Class Osteichthyes Huxley, 1880

*Remarks* — Although some teeth of osteichthyans are highly characteristic, in general isolated dental remains of osteichthyans are probably best regarded as being of relatively little taxonomic use. This is especially true of deeper-water fishes, where strong convergence is seen within relatively simple teeth morphologies. The difficulty in applying standard taxonomic methodology to isolated dental material of deep-water osteichthyans (ichthyoliths) has led to the development of generic descriptive methods for this material (see Doyle & Riedel, 1985). This generic descriptive terminology is here regarded as cumbersome and is not followed within this study. Despite this general difficulty in assigning taxonomic position to the isolated osteichthyan teeth described here, it may be possible to tentatively assign the molariform teeth of Tooth type 8 of this study to the Sciaenidae.

Tooth type 1  
Figure 3/4

*Material* — One partial tooth, UF 206540.

*Description* — Although missing the apex, this tooth is extremely elongate. It is gently curved and oval in cross section. A narrow and delicate cutting edge is present on the upper part of the posterior edge of the tooth.

Tooth type 2  
Figure 3/5

*Material* — Several teeth, including UF 206541.

*Description* — These teeth are high and somewhat curved, sometimes faintly sigmoidally. The tooth is oval in cross section and hollow for a large proportion of its length. A small, clear, enamel tip is present, which is expanded posteriorly into a short cutting edge. This cutting edge takes up less than a third of the posterior edge, terminating sharply to form a distinct barb.

Tooth type 3  
Figure 3/6

*Material* — One tooth, UF 206478.

*Description* — The tooth is elongate and somewhat laterally compressed. The anterior edge of the tooth is rounded for the lower two thirds, with a well-developed cutting edge in the upper part. The base of the anterior cutting edge is coincident with a slight posterior curva-



ture. A well-developed cutting edge is present along the entire posterior edge.

Tooth type 4  
Figure 3/7

*Material* — Common teeth, including UF 206477.

*Description* — These high and triangular teeth are very strongly compressed. Well-developed cutting edges are present on both sides of these symmetrical teeth. The preserved part of these teeth is composed entirely of clear enamel, with a conical basal cavity being clearly visible.

Tooth type 5  
Figure 3/8

*Material* — One tooth, UF 206542.

*Description* — The single tooth of this morphology is triangular and very strongly compressed. A weakly-developed cutting edge covers both the convex anterior edge and the concave posterior edge. The central part of the tooth is taken up with a large basal cavity. There is no obvious clear enamel cap.

Tooth type 6  
Figure 3/9

*Material* — Several teeth, including UF 206480.

*Description* — These somewhat elongate teeth are round in cross section and have slight curvature. A small conical clear enamel cap is present.

Tooth type 7  
Figure 3/10

*Material* — One tooth, UF 206476.

*Description* — This conical tooth is elongate with no curvature. The apex has a small and blunt clear enamel cap. The basal region is faintly flared.

Tooth type 8  
Figure 3/11, 12

*Material* — Several teeth, including UF 206475, 206479.

*Description* — Although very variable in size, molariform teeth comprise a relatively small number of different morphologies. The bulk are circular to faintly oval in occlusal view and approximately hemispherical in overall shape. A weakly developed apical boss is sometimes present. Rare examples of oval teeth have the api-

cal boss stretched out to form a poorly-developed occlusal crest. No clear enamel tip was observed on any teeth.

### Endodontal borings

Borings are commonly present within fossil bone material, especially that deposited within a marine setting. In many cases this boring can occur with sufficient intensity to severely damage or destroy parts of the fossil. Despite the abundance of these endodontal borings, they have rarely been described.

Circular borings (possibly attributable to *Oichnus* Bromley, 1981)  
Figure 3/13

*Material* — Borings in three shark teeth, including UF 206534.

*Description* — Widely dispersed circular holes of between 30 and 50  $\mu\text{m}$  are seen on the surface of several shark teeth. They usually penetrate the root, although at least two were seen cutting the lower part of the enameloid crown. It was not possible to see the morphology of the borings within the tooth, although it is evident that they are deep and are not straight.

*Comparisons* — It is unclear whether these borings are the same as the flask-shaped borings in osteichthyan teeth described by Underwood *et al.* (1999). Despite the similarities in the shape and general size of the entrance holes, the deep and curved shape of the borings recorded here do not seem to be very similar to the Cretaceous borings of Underwood *et al.* (1999).

Ichnogenus *Mycelites* Roux, 1887

*Type species* - *Mycelites ossifragus* Roux, 1887; Jurassic to Miocene.

*Mycelites ossifragus* Roux, 1887  
Figure 3/14, 15

- 1887 *Mycelites ossifragus* Roux, pls 14, 15; figs 1-7.
- 1989 *Mycelites enameloides* Martill, pl. 1.
- 1999 *Mycelites ossifragus* Roux — Underwood *et al.*, fig.1b-h.

*Material* — Borings in many teeth, including UF 206535.

*Description* — These borings are relatively tubular and are generally about 5  $\mu\text{m}$  in diameter. Bifurcate branching present, but relatively uncommon. They are

present both within the tooth and embedded within the tooth surface. These borings are seen to form irregular radiating masses up to 400 µm across.

*Comparisons* — Although many of the examples of *Mycelites* show a clear radial form, this is far more irregular than that of *Abeliella riccioides* Mägdefrau, 1937.

### Palaeoecology

Despite the relatively small sample size, the material described here has important implications as to the water depth at the time of deposition of the Montpelier Formation at Duncans Quarry. Tropical shelf seas are typically inhabited by a diverse suite of selachians, including orectolobids, triakids, carcharhinids, rhinobatids and dasyatids. The absence of teeth of these forms, and the presence of a squalid-dominated assemblage, is highly indicative of a deeper-water environment.

All three of the genera of Squaliformes recorded in the present study are basically deep-water forms (*e.g.*, Compagno, 1984; Froese & Pauly, 2001). Extant species of both *Deania* and *Scymnodon* are essentially benthopelagic inhabitants of continental slopes, being rarely found in less than 200 m of water. Although the maximum depths of many species are poorly known, most living species have been recorded at depths of between 1,000 and 2,000 m. Extant species of *Squaliolus* are bathypelagic with well-developed luminescent organs. They undergo vertical migration, rising to about 200 m during the night and returning to between 1,000 and 2,000 m depth during the day. They are typically commonest near oceanic islands.

The numerous living species of *Carcharhinus* are very variable in their habitat. Although many species are present in shallow water, others are truly oceanic, living in surface waters away from land. The presence of indeterminate *Carcharhinus* is, therefore, palaeoenvironmentally undiagnostic, although teeth of oceanic taxa are to be expected within offshore deposits.

The osteichthyan assemblage is not readily interpreted due to the inherent difficulties of working with isolated teeth. Despite this, the dominance of elongate 'grasping' teeth is highly suggestive of the presence of mesopelagic and bathypelagic predators as opposed to more tropically diverse shallow-water forms.

Consequently, the fish assemblage indicates that the chalks at Duncans were deposited in water depths of more than 200 m and possibly significantly more. This is particularly significant, since Duncans Quarry is situated only a few km north of the platform margin (Figure 1). It indicates that there was a well-defined, fault-bounded northern margin to the Miocene carbonate platforms, similar to that of the modern north coast

of Jamaica.

The presence of borings within many of the teeth is highly suggestive of significant reworking of biogenic phosphate (Martill, 1989; Underwood *et al.*, 1999). However, it should be noted that the intensity of boring is far lower than that commonly recorded within shelfal assemblages. The implications of this are unclear, as the nature of the organisms involved in endodontal boring and the controls on the growth of these organisms are still very poorly known (but see Gouget & Locquin, 1979).

### Acknowledgements

We thank Roger Portell (UF) for allowing us to study the fish material from Duncans Quarry. We would also like to thank David Ward for his advice and help with the literature, and Andy Beard for his assistance with the SEM. We gratefully acknowledge Barbara and Reed Toomey, who not only assisted in the collection of this material in the field, but also provided the funding for several field trips to Jamaica that facilitated the collection of this material. Bob Purdy (National Museum of Natural History, Smithsonian Institution) kindly made comments on an earlier version of this paper.

### References

- Blainville, H.M.D. de. 1816. Prodrome d'une distribution systématique du règne animal. *Bulletin de la Société Philomatique de Paris* 8, 105-124.
- Bocage, J.V.B. du & Capello, F. de B. 1864. Sur quelques espèces inédites de Squalidae de la tribu Acanthiana, Gray, qui fréquentent les côtes du Portugal. *Proceedings of the Zoological Society of London* 1864, 260-263.
- Bromley, R.G. 1981. Concepts in ichnotaxonomy illustrated by small round holes in shells. *Acta Geológica Hispánica* 16, 55-64.
- Cappetta, H. 1980. Les séliaciens du Crétacé supérieur du Liban. *Palaeontographica* A168, 69-229.
- Casier, E. 1958. Contribution à l'étude des poissons fossiles des Antilles. *Mémoires Suisses de Paléontologie* 74, 1-95.
- Cigala-Fulgosi, F. 1986. A deep water elasmobranch fauna from a Lower Pliocene outcropping (northern Italy). In: Uyeno, T., Arai, R., Taniuchi, T. & Matsuura, K. (eds). *Indo-Pacific fish biology. Proceedings of the Second International Conference on Indo-Pacific fishes*, 133-139. Tokyo (Ichthyological Society of Japan).
- Compagno, L.J.V. 1973. Interrelationships of living elasmobranchs. In: Greenwood, P.H., Miles, R.S. & Patterson, C. (eds). *Interrelationships of fishes. Zoological Journal of the Linnean Society* 53 (Supplement), 15-61.
- Compagno, L.J.V., 1984. FAO species catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 1. Hexanchiformes to Lamniformes. *FAO Fisheries Symposium* 125 (4, 1), 1-249.
- Donovan, S.K. & Gunter, G. 2002. Fossil sharks from Ja-



- maica. *Bulletin of the Mizunami Fossil Museum* 28 (2001), 211-215.
- Donovan, S.K., Jackson, T.A., Dixon, H.L. & Doyle, E.N. 1995. Eastern and central Jamaica. *Geologists' Association Guides* 53, i + 1-62.
- Doyle, P.S. & Riedel, W.R. 1985. Cenozoic and late Cretaceous Ichthyoliths. In: Bolli, H.M., Saunders, J.B. & Perch-Nielsen, K. (eds). *Plankton stratigraphy*, 865-996. Cambridge (Cambridge University Press).
- Eyles, V.A. 1958. A phosphatic band underlying bauxite deposits in Jamaica. *Nature* 182, 1367, 1368.
- Froese, R. & Pauly, D. (eds). 2001. *FishBase*. [Live 5 November 2001.] <www.fishbase.org>
- Goodrich, E.S. 1909. Vertebrate Craniata (first fascicle: cyclostomes and fishes). In: Lankester, E.R. (ed.). *A treatise on zoology, part 10*, 518 pp. London (Adam and Charles Black).
- Gouget, D. & Locquin, M.V. 1979. Découverte de spores fongiques dans les écailles de poissons et d'agnathes paléozoïques: *Mycobystrovia lepidophaga* gen. et. sp. nov. *104e Congrès national des Sociétés savantes, 1979, Sciences* 1, 87-99.
- Herman, J., Hovestadt-Euler, M. & Hovestadt, D.C. 1989. Contributions to the study of the comparative morphology of teeth and other relevant ichthyodorulites in living supraspecific taxa of chondrichthyan fishes. Part A: Selachii. No. 3: Order: Squaliformes - Families: Echinorhinidae, Oxynotidae and Squalidae. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Biologie* 59, 101-157.
- Herman, J., Hovestadt-Euler, M. & Hovestadt, D.C. 1993. Contributions to the study of the comparative morphology of teeth and other relevant ichthyodorulites in living supraspecific taxa of chondrichthyan fishes. Part A: Selachii. No. 1b: Order: Hexanchiformes - Family: Chlamydoselachidae; No. 5: Order: Heterodontiformes - Family: Heterodontidae No. 6: Order: Lamniformes - Families: Cetorhinidae, Megachasmidae; Addendum 1 to No. 3: Order: Squaliformes; Addendum 1 to No. 4: Order: Orectolobiformes; General Glossary; Summary Part A. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Biologie* 63, 185-256.
- Huxley, T.H. 1880. On the application of the laws of evolution to the arrangement of the Vertebrata and more particularly of the Mammalia. *Proceedings of the Zoological Society of London* 1880, 649-662.
- Jordan, D.S. & Snyder, J.O. 1902. Descriptions of two new species of squaloid sharks from Japan. *Proceedings of the United States National Museum* 25, 79-81.
- Ledoux, J.-C. 1972. Les Squalidae (Euselachii) miocènes des environs d'Avignon (Vaucluse). *Documents des Laboratoires de Géologie de la Faculté des Sciences de Lyon* 52, 133-175.
- Lowe, R.T. 1839. A supplement to a synopsis of the fishes of Madeira. *Proceedings of the Zoological Society of London* 1839, 76-92.
- Mägdefrau, K. 1937. Lebensspuren fossiler "Bohr"- Organismen. *Beiträge zur naturkundlichen Forschung in Südwestdeutschland* 2, 54-67.
- Martill, D.M. 1989. Fungal borings in neoselachian teeth from the lower Oxford Clay of Peterborough. *Mercian Geologist* 12, 1-5.
- Mitchell, S.F. 2004. Lithostratigraphy and palaeogeography of the White Limestone Group. In: Donovan, S.K. (ed.). The Mid-Cainozoic White Limestone Group of Jamaica. *Cainozoic Research* 3, 5-29.
- Purdy, R.W., Donovan, S.K., Pickerill, R.K. & Dixon, H.L. 1996. Fish teeth from the Pleistocene of Jamaica. *Journal of Vertebrate Paleontology* 16, 165-167.
- Purdy, R.W., Schneider, V., Applegate, S., McLellan, J., Meyer, R. & Slaughter, R. 2001. The Neogene sharks, rays, and bony fishes from Lee Creek Mine, Aurora, North Carolina. In: Ray, C.E. & Bohaska, D.J. (eds). *Geology and paleontology of the Lee Creek Mine, North Carolina, III. Smithsonian Contributions to Paleobiology* 90, 71-202.
- Quoy, J.R.C. & Gaimard, J.P. 1824-1825. Description des Poissons. In: Freycinet, L. de, *Voyage autour du Monde exécuté sur les corvettes de L.M. "L'Uranie" et "La Physicienne," pendant les années 1817, 1818, 1819 et 1820*, 192-401. Paris (Voyage Uranie, Zoologie).
- Roux, W. 1887. Über eine im Knochen lebende Gruppe von Fadenpilzen (*Mycelites ossifragus*). *Zeitschrift der wissenschaftlichen Zoologie* 16, 227.
- Shirai, S. 1996. Phylogenetic interrelationships of neoselachians (Chondrichthyes: Euselachii). In: Stiassey, M.L.J., Parenti, L.R. & Johnson, G.D. (eds). *Interrelationships of fishes*, 9-34. San Diego (Academic Press).
- Smith, H.M. & Radcliffe, L. 1912. Description of a new family of pediculate fishes from Celebes [Scientific results of the Philippine cruise of the Fisheries steamer "Albatross," 1907-1910, No. 20]. *Proceedings of the United States National Museum* 42, 579-581.
- Steineck, P.L. 1974. Foraminiferal paleoecology of the Montpelier and lower Coastal Groups (Eocene-Miocene), Jamaica, West Indies. *Palaeogeography, Palaeoecology, Palaeoceanography* 16, 217-242.
- Stringer, G.L. 1998. Otolith-based fishes from the Bowden shell bed (Pliocene) of Jamaica: systematics and palaeoecology. *Contributions to Tertiary and Quaternary Geology* 35, 147-160.
- Underwood, C.J., Mitchell, S.F. & Veltkamp, C.J. 1999. Microborings in mid Cretaceous fish teeth. *Proceedings of the Yorkshire Geological Society* 52, 269-274.
- Vaillant, L., 1888. *Poissons. Expéditions scientifiques du 'Travailleur' et du 'Talisman' pendant les années 1880-1883*, 406 pp. Paris.