

THE INDIRECT ACTION OF EXTERNAL FACTORS ON PHOTOSYNTHESIS

by

F. VAN DER PAAUW.

Contents.

Chapter	Page
I. INTRODUCTION.....	500
A. Blackman's Formula of Limiting Factors	500
B. Do the External Factors Act Directly on the Photosynthetic Process, or Do Internal Protoplasmic Processes Interact?	502
II. EXPERIMENTAL METHOD	509
A. Material	509
B. The Apparatus of Van den Honert Used in my First Experiments	511
C. The Apparatus Used in my Further Expe- riments.....	512
1. Description of the apparatus	512
2. Fixation of the algal film on the glass cover	514
3. The regulation of temperature.....	515
4. The regulation of CO ₂ concentration	516
5. Illumination	516
D. Course of an Experiment and Corrections	518
III. THE RESPIRATION DURING THE ASSIMILATION	522
A. Introduction	522
B. Experiments	524
IV. THE ASSIMILATORY QUOTIENT	532
V. THE INFLUENCE OF CARBON DIOXIDE CONCEN- TRATION ON ASSIMILATION	534

Chapter	Page
A. The CO_2 Concentration-Assimilation Curve	534
B. The Influence of Carbon Dioxide Concentration when the Assimilation has been Retarded by Narcotics.....	536
C. The Influence of Temperature on the CO_2 Concentration-Assimilation Curve when Phenylurethane has been Added	540
VI. THE INFLUENCE OF LIGHT INTENSITY ON ASSIMILATION.....	541
A. Introduction	541
B. Experiments on the Influence of Light Intensity on the Assimilation of Hormidium Cultivated in Daylight.....	543
C. Experiments on the Influence of Light Intensity on the Assimilation of Hormidium Cultivated in Weak Artificial Light	544
D. The Influence of Light Intensity on Assimilation at Different Temperatures....	546
VII. THE INFLUENCE OF TEMPERATURE ON ASSIMILATION AND RESPIRATION	550
A. Introduction	550
B. The Influence of Temperature on Assimilation.....	555
C. The Influence of Temperature on Respiration	560
VIII. THE INFLUENCE OF POTASSIUM CYANIDE ON ASSIMILATION AND RESPIRATION	564
A. Introduction	564
B. Experiments with Low Concentrations of KCN, Showing a Stimulating Influence	565
C. Experiments with Higher Concentrations of KCN, Showing a Retarding Influence	570
D. Experiments, Showing a Complete Inhi-	

Chapter	Page
bition of Assimilation after the Addition of KCN.....	574
E. Discussion of the Experimental Results	576
IX. THE INFLUENCE OF PHENYLURETHANE ON ASSI- MILATION AND RESPIRATION	578
X. THE INFLUENCE OF GLUCOSE AND FRUCTOSE ON ASSIMILATION AND RESPIRATION	585
XI. THE INFLUENCE OF BaCl_2 ON ASSIMILATION AND RESPIRATION	590
XII. THE TIME COURSE OF PHOTOSYNTHESIS.....	593
A. The Induction-Time of Photosynthesis	593
1. Introduction	593
2. Experiments on the Induction of Pho- tosynthesis at Different Temperatures	595
B. The Course of Photosynthesis after a Long Period of Darkness	599
XIII. DISCUSSION	601
A. The Assimilation Reacts to the Addition of Uncommon Substances in the Same Way under Different Conditions.....	601
B. The Dependence of Photosynthesis of Internal Factors	602
C. The Explanation of the Parallelism of Respiration and Photosynthesis	607
D. The Reaction of Photosynthesis to Inju- rious Influences	610
E. An Attempt to Explain the too Large Deviations of Some Determinations on the Influence of Temperature on Photo- synthesis from the Average, and a Few Remarks on Optimum Curves.....	612
F. Blackman's Rule of Limiting Factors..	615
Summary of the Principal Results	616
Literature	618

CHAPTER I.

Introduction.

In this chapter a full discussion of the literature will not be aimed at. Very extensive surveys are a.o. to be found in the monographs of Stiles and Spoehr; I also refer to the short review of literature of van den Honert. In this place only that which pertains to the problem of this paper will be discussed.

A. Blackman's Formula of Limiting Factors.

In the year 1905 Blackman stated in his well-known paper: „When a process is conditioned as to its rapidity by a number of separate factors, the rate of the process is limited by the pace of the „slowest” factor.” In Blackman's opinion the relation between an external factor and the rate of photosynthesis can be represented by a straight curve with a sharp break.

This statement is founded on experiments of Miss Matthaei (1904), and is further supported by experiments of Blackman and Matthaei (1905), and of Blackman and Smith (1911).

The accuracy of these experiments was of such a nature that a possibility of other interpretation remained (criticism of Brown and Heise (1917, 18). Experimental attacks on the theory of limiting factors came from the part of many investigators. Especially the paper of Harder (1921), which was altogether devoted to this question, is of importance. He stated that in case of low „concentration” of one factor, an increase in the concentration of the other (non-limiting) factor, will cause an acceleration of the process. According to Harder the relationship between photosynthesis and the external factors can be expressed by a smooth curve.

A theoretical paper against the theory was published by Romell (1926).

So, the problem seemed to be settled against Blackman's formulation, were it not that van den Honert (1928, 30) seemed to have confirmed this theory experimentally, in very accurate experiments on the influence of the CO_2 factor, made with films of the filamentous alga *Hormidium flaccidum*. The relation between CO_2 concentration and assimilation, represented in a graph, shows a straight curve with a sharp break, resembling the scheme proposed by Blackman.

It must also be mentioned that Boysen Jensen and Müller (1929) discovered a Blackman curve for the action of light on shade plants of *Marchantia polymorpha*. The number of points, determined on this curve is, however, rather small.

The light intensity-assimilation curve of van den Honert, on the other hand, shows a logarithmic type (as Harder's curve), but van den Honert is of opinion that a better agreement with Blackman's scheme should be possible, if the light intensity on the light- and shade-side of the assimilating cells could be equalized.

A reliable curve of the Blackman-type obtained experimentally is an argument of more importance than ten curves showing a gradual course, and constitutes therefore a great support for the adherents of Blackman's formula. A smooth curve may also occur (even if Blackman's postulate were true), if the experimental technique were inexact and not all other factors than the one studied were kept constant (cf. also Boysen Jensen and Müller).

The fact that van den Honert did not find a Blackman-curve when the influence of light was studied, therefore, does not preclude the possibility that under improved experimental conditions a Blackman-curve would indeed appear. It is true that van den Honert worked with a film one-cell in thickness, but objections arise against the manner of illumination used. A solution of the

question how *Homidium* behaves under improved experimental conditions has been sought in Chapter VI.

Van den Honert explains his CO_2 concentration-assimilation curve by assuming that in case of low CO_2 concentrations the diffusion of CO_2 will limit the assimilation, and that the direct proportionality arises from the quick absorption of CO_2 in the organism.

Van den Honert succeeded in making it quite plausible that the diffusion process limits indeed, by showing that the process is not at all sensitive to temperature, which is in conformity with the properties of the diffusion of CO_2 in water (cf. also Bohr 1897 and 99).

Though van den Honert has shown that a curve with a proper resemblance to the Blackman-scheme is obtainable, this may be a coincidence and an exception. The same opinion is held by van den Honert in the theoretical discussion of his results (v. d. H. 1930, p. 245). The fact itself does not speak for the *general* validity of Blackman's theory.

In the experiments of Warburg (1919, 20), in which the diffusion plays no part, as he could prove by change in temperature, a logarithmic curve was obtained.

In other experiments Warburg ascertained that narcotics retard the assimilation. I, therefore, have asked myself how the CO_2 concentration-assimilation curve of narcotized *Homidium* would proceed, and whether it would be possible in this way to cause the Blackman-type to disappear. It may be possible that the limiting action of diffusion of CO_2 will be excluded by this interference. This question will be treated in Chapter V.

B. Do the External Factors Act Directly on the Photosynthetic Process, or Do Internal Protoplasmic Processes Interact?

Willstätter and Stoll (1918) and O. Warburg

(1919, 20, 22, 24) have very thoroughly studied the assimilation of carbon dioxide, and they have arrived experimentally at a theory of photosynthesis. I am not going to consider the contents of these theories here, in which case the opinion of other investigators should have to be mentioned too, but only a tendency which these theories have in common.

Photosynthesis is considered here as a reaction in an „unalterable” system (test-tube), while the possibility that this system itself (the organism) changes under the influence of the varying factors, is ignored and not even suggested ¹⁾.

With high light intensity and high concentration of CO₂, temperature is the limiting factor of photosynthesis. It is supposed that under these conditions the rate of the whole process is controlled by the speed of a pure chemical part of it (Blackman reaction). In the theories of Willstätter and Stoll, Warburg, a.o., this chemical reaction or reaction complex is, as a matter of fact, considered as a stage of the photosynthetic process itself. To be sure, this possibility may not be denied, but that it should be obvious is not at all evident. In fact it is possible, that internal processes determine the intensity of the assimilation, and that the accelerated action after an increase of temperature is merely based on the acceleration of these internal processes (alterations of the system in which the reaction occurs). In this case some „branch-chain” of the process will be studied, and not a part of the reaction: $6 \text{ CO}_2 + 6 \text{ H}_2\text{O} \rightarrow \text{C}_6 \text{ H}_{12}\text{O}_6 + 6 \text{ O}_2$.

Quite another conception than the first mentioned, which we may call „chemical” (on account of the internal regulatory processes being neglected) we find in the work

¹⁾ I do not mean the „adaptation” to the factors of the milieu after a long time, which is certainly acknowledged by these investigators, but an immediate reaction of the plasma during the experiment.

of Spoehr and Mac Gee (1923), Kostytschew and his collaborators (1926—31), Harder (1930), and Arnold (1931). The first mentioned investigators assume a dependence of the assimilation on the respiration. They observed that after a long period of darkness, the assimilatory intensity ran parallel to that of the respiration and to the increase in sugar-content. From this they concluded that active products of carbohydrate-metabolism influenced assimilation. The experimental foundation of the theory is weak, but still the assimilation is looked upon as a physiological process being dependent on internal processes.

Kostytschew and collaborators published a series of investigations by which it was shown that assimilation is only slightly influenced by the prevailing external factors. Fluctuations occurred, without there being a noticeable external cause, so that they came to the theory (Kostytschew 1931), that assimilation is dependent on internal factors and on the after-effect of external factors („höchst komplizierte Reizketten"). The irregular assimilation they found appears partly to be under the influence of movements of the stomata, but can also appear when the stomata are quite open.

It was even shown that under very favourable circumstances an evolution of CO_2 may take place. Boonstra (1930), however, could not succeed in confirming in Holland this phenomenon, which has been observed in four different climates in Russia.

Kostytschew was also criticized by Boysen Jensen and Müller (1929), who observed a more regular course of the assimilation. According to them Kostytschew's results are probably a consequence of experimental errors. We wonder indeed how it is possible that rather regular curves, as obtained by Matthaei (1904), Willstätter and Stoll (1918), Lundegårdh (1921), would have been possible if the assimilation would always be subject to

such large fluctuations. As a result of his opinion Kostytschew denies any value of experiments of short duration, in which external factors are varied.

Harder (1930) found that the intensity of assimilation in aquatics changes with time. He found no fluctuations as Kostytschew a.o. did, but a gradual increase of the assimilation in light, followed by a decrease. He explains his results by distinguishing a number of reaction swchich must all be of internal nature, called by him „Aktivierung“, „Gegenreaktion“, „Ermüdung“ and „Anpassung“.

Arnold (1931), who almost at the same time arrived at similar results with *Elodea*, agrees with this opinion.

Other investigators, who did not find a constant course of the assimilation-intensity, are Ewart (1897, 98), Pantanelli (1904), Lubimenko (1905), Willstätter and Stoll (1918), Johansson (1923, 26), Plantefol (1927), Montfort and Neydel (1928).

Willstätter and Stoll ascribed this fact to an accumulation of products of assimilation; the explanation is altogether „chemical“.

Johansson explains the decrease in intensity by the closing of the stomata, so his explanation does not call real internal factors into play either.

Of great importance is, therefore, the work of Montfort and Neydel (1928), who observed the same phenomena with leaves of *Hydrophyllacea*, in which stomata are wanting. They could show that neither accumulation of products of photosynthesis (a decrease began after a short illumination), nor desiccation (the leaves were in water), nor alteration of the position of the chloroplasts (after change in position more light gave less assimilation) caused the decrease, so that it is obvious to account for these facts by means of internal reactions, the more so as the preliminary treatment also proved to have influenced the reaction. With strong light these investigators, too, did

find evolution of CO_2 , however, not alternating with periods of strong assimilation, as Kostytschew found.

Very large fluctuations within the lapse of an extremely short time were found by Maximow and Krassnoselsky—Maximow (1928). They do not know, whether the cause is internal or a result of the movement of the stomata.

Beljakoff (1929), however, did not find these large „jumps”, and ascribes them to experimental errors. Moreover he found that the stomata do not react rapidly on light, temperature and moisture, in contrast to the above-mentioned investigators.

Unusual assimilation-curves, which also point to a very complicated composition of the assimilatory-process were found by Lundegårdh and his collaborators Stocker (1927), Walther (1927), Yoshii (1928). Their curves are characterised by several optima and minima.

Van den Honert (1928, p. 9; 1930, p. 158), however, severely criticizes the apparatus of Lundegårdh and remarks that inasmuch the measurements are made immediately after the illumination, the absorption of CO_2 cannot be constant in the first minutes. To this I would add that for another reason it is undesirable to measure in the first minutes. The opinion that the assimilation is connected with very complicated processes, demands that the organism is allowed some time for the creation of a state of equilibrium. Though the truth of this assertion may be doubted, yet the possibility must be taken into account.

In connection with this last point the work of Tsi-Tung Li (1929) should also be mentioned, who found an immediate considerable increase of assimilation after altering the intensity of the light or its composition (red-white). Tsi-Tung Li explains this by considering the available surface to be a limiting factor. When taken out of the darkness into light there will be a large surface available in the beginning, which will diminish later on.

So he explains the phenomenon by an internal limiting factor, though not an *active* internal one. The case may be more complicated, but it clearly demonstrates the danger of premature measurements.

In the preceding pages we have mentioned a number of modern investigations, where there was more or less reason to consider the assimilation connected with other vital functions. The study of this correlation has become the principal object of the present paper.

In this the conception of Spoehr and Mc Gee have served as a working-hypothesis. I have always asked myself how both assimilation and respiration behave under different conditions. I was led to their hypothesis, not because a direct relation between photosynthesis and respiration seemed likely *a priori*, but because of the fact that, if photosynthesis should be connected with protoplasmic activity, the oxygen consumption of the protoplasm could be taken as a fair measure of this activity.

For this reason the influence of temperature on photosynthesis is compared with the influence of this factor on respiration. (Chapter VI). In itself the determination of the influence of temperature on assimilation is also of importance, because the literature shows, in addition to the none too reliable determinations with leaves of higher plants, only the data of Emerson (1929), who worked with *Chlorella pyrenoides*. All other determinations are more or less fragmentary.

The influence of four different substances on respiration and assimilation is investigated in the Chapters VIII—XI. Since photosynthesis is considered to be a chain process by many investigators (a.o. Briggs, Warburg, van den Honert) and since different reactions possibly control the rate of photosynthesis under different conditions, the influence of these chemicals is determined both in light of low and of high intensity.

Finally, in Chapter XII, I have tried to find an answer to the question whether any processes precede the assimilation. If the assimilation is really dependent on internal processes we may expect that the equilibrium will not be immediately established. We know already the so-called „photochemical induction” of assimilation since the inquiry of Warburg (1920), but it is doubtful whether we have really to do here with a photochemical phenomenon or with physiological reactions which must precede the assimilation. It has been tried to determine the nature of this „preceding” reaction.

In the course of the investigations the necessity became apparent to study respiration as well. Its magnitude during photosynthesis should be established. Fluctuations in the intensity of respiration might be the cause of fluctuations in our analytical results. The study of respiration becomes a necessity when a causal relation between this form of metabolism and assimilation is claimed (Spoehr and Mc Gee). This problem will be treated in Chapter III.

With the exception of a few, which are mentioned in Chapter V, and which were made with the apparatus of van den Honert (1928, 30), the experiments were made in a new constructed, very simple, but at the same time very accurate apparatus, according to the manometrical method.

The advantage of van den Honert's method, the experimentation with a film of *Hormidium* of one cell in thickness was maintained in it. The assimilation of this film is, according to van den Honert very constant during a long time. Kostytschew's objection against the determination of the influence of external factors, which would be impossible on account of the inconstant course, does not hold here.

The apparatus will be described in the following chapter.

The method is based on the measurement of the evolution of O_2 .

In order to check my results with those of van den Honert (analyses of CO_2 concentration) the assimilatory quotient was determined (Chapter IV). This was all the more necessary, because van den Honert had found deviations of this quotient from unity in a few determinations.

CHAPTER II.

Experimental Method.

A. *Material.*

A strain of the filamentous alga *Hormidium flaccidum* served as experimental object. The same strain was used by van den Honert in his investigations. He determined the strain as *H. flaccidum*, which species is subdivided in different sub-species in Pascher's Flora. This strain forms, when cultivated in a suitable way, a film of only a single cell layer on the nutrient solution and is, in this state, very suitable for the investigation of photosynthesis. I obtained another strain, the sub-species *Hormidium nitens* from the collection of Dr. E. G. Pringsheim at Prague. This latter strain was free from bacteria, which is not the case with the former. With the latter strain it appeared more difficult to obtain „films”. Moreover the film is less close, which causes loss of material when removed to other liquid culture media (which was necessary in the experiments of Ch. VIII, IX, XI). For these reasons I mostly used van den Honert's strain; the quantity of bacteria is, moreover, small in a well growing culture. For the sake of comparison other organisms (*Stichococcus bacillaris*, *Oocystis spec.*) were tested occasionally.

The cultivation succeeds very well in Erlenmeyer flasks (contents 150 cm³), which, closed by a cotton-wool

stopper, are placed in a cool room before a north-window; the conditions of cultivation were therefore not constant. In summer the light must be dimmed a little with white paper. Very good cultures were obtained at a temperature of 15°—18° C. A lower temperature is not directly injurious but the growth becomes very slow. At higher temperatures Pringsheim's strain soon forms less fine films; van den Honert's strain may keep well, provided large quantities are constantly transferred. If this is not done, the alga gets overgrown with bacteria. At least once a week the cultures must be transferred; in winter this may be done less frequently. In the summer of 1930 I grew the algae in a Hearson's cool biological incubator, in which the temperature was kept at 15°—16°. The flasks are filled with 20—30 cm³ nutrient solution of the following composition, given by van den Honert; water distilled from glass into glass, containing:

Fe SO ₄	0.001 %
Mg SO ₄	0.01 %
Ca SO ₄	0.004 %
K NO ₃	0.1 %
K H ₂ PO ₄	0.02 %

In this medium also other algae, such as *Stichococcus bacillaris*, *S. minor* and *Oocystis spec.* appeared to grow well.

Besides on the windowsill, I also cultivated the algae with a constant source of light. For this purpose Erlenmeyer flasks (contents 1 litre) were used, shut at the top by a rubber stopper containing two passages. With the aid of an aquarium-pump indoorair continually bubbles through the liquid. The flasks were placed round a 75 Watt milkglass Philips lamp burning day and night and surrounded by a watercooler. After three or four weeks the liquid becomes darkgreen. *Hormidium* is in this case usually unicellular and forms no film.

B. The Apparatus of Van den Honert Used in my First Experiments.

A few experiments (those mentioned in Chapter V) were made with the apparatus of van den Honert. According to his method a current of air containing CO_2 is led at a great speed over a film of *Hormidium*. The used air is not directly removed, but is again led over the alga by means of a small pump. The supply of air containing CO_2 and the removal of air samples for the analysis takes place very slowly in comparison to the speed of the circulating current of air. After some time a state of equilibrium is attained and the tension of CO_2 of the circulating air becomes rather constant. With the aid of a gasanalysis apparatus of Krogh the tension of CO_2 is determined accurately within 0.001 %. The same had been done before with the supplied air. From the quantity of air removed per unit of time and the difference in CO_2 contents between supplied and removed air the assimilation of carbon dioxide may be calculated.

We refrain here from an extensive description. The reader can find it in the publications of van den Honert (1928, 30).

The method is very accurate. An objection, however, is the long duration of the experiment; a disturbance in the intricate apparatus may easily set in. The greatest difficulty is the handling of the Krogh apparatus. This apparatus suffers greatly from leakages, which often occur, especially (as in my case), when it has already been used before, and the stopcocks become older and less true. Moreover the making of a great number of analyses is very trying work. Another drawback is that the apparatus soon gets dirty; the regular cleaning takes up much time. A last objection of the method is, in my opinion, that it may be called accurate, but it is not supple. The course of the process is difficult to follow. When the air is removed, one never knows if the equilibrium is already attained.

For all these reasons I have given up working with this apparatus, and constructed a new one, according to the manometrical method, which fitted better to the purpose of the investigation.

For the present a few more remarks about the apparatus of van den Honert. Some minor improvements have been introduced. The glass tube, numbered 23 in fig. 8 of van den Honert (1928, 130) has been replaced by an unbreakable brass one. The valves, shown in fig. 9 are constructed more simply. Instead of string and a melted wax and resin mixture, a piece of rubber tube is used. This is put onto tube 1, after which tube 5 is passed round it, so that a watertight closure is effected.

C. *The Apparatus Used in my Further Experiments.*

1. *Description of the apparatus.* This apparatus (cf. fig. 1) is an application of the microrespirometer of Krogh for the assimilation of carbon dioxide. Two vessels (*a* and *b*) are connected by means of a manometer (*c*), one of these vessels (*b*) serves as control, and contains only air and water. It served only to enable us to compare the varying pressure of the air with the constant pressure in the other vessel, and to preclude the effect of slight fluctuations in temperature.

The principle of the method is to allow the alga to assimilate or respire in a space of constant CO_2 tension, so that only the oxygen tension changes. This change is indicated by the rise or fall of the liquid in the manometer. By adding mercury out of a calibrated capillary tube (*d*), the original pressure can be restored and the quantity of O_2 evolved or absorbed, which is equal to the volume of mercury displaced can be read. The constant tension of CO_2 is reached with a so-called buffer-solution of Na_2CO_3 and NaHCO_3 . So the method, contrary to that of van den Honert, is based on the measurement of alterations in the

tension of oxygen and not on the determination of the carbon dioxide concentration.

The assimilation- and the control-vessel are very shallow and made of nickel plated brass (cf. fig. 1 A). The inner diameter is 106 millimeter, and the outer 124, the height only 4 mm. They are covered with a glass plate (f) 3 mm thick (a cover of a glass box). Between vessel and plate is a greased rubber ring (e). The plate is clasped on the




Fig. 1. Sketch of the apparatus. For description see text.

vessel with the aid of a brass ring (h) and five clamping-screws (j). In order to prevent cracking of the glass there is also a rubber ring (g) between ring and glass.

On the bottom of the vessels we put 20 cubic centimetres of the buffer mixture. There is an opening in it leading through a twice bent glass tube to the manometer. To prevent the liquid from entering this opening a short tube 4 mm long (k) has been soldered on it. Besides the assimilation-vessel contains an opening (l) communicating with the calibrated capillary tube (d). This connection is esta-

blished by the twice bent tube (*m*), in which there is a small bulge (*n*), which serves as a mercury reservoir. The capillary tube (*d*) is a pipette of one cubic centimetre.




Fig. 1A. Bottom of the assimilatory vessel (*a*) with glass-cover (*f*), metal ring (*h*), rubber rings (*e* and *g*), clamping-screw (*j*). For further description see text.

It is partly filled with water, partly with mercury. It is calibrated to 0.01 cm³ and can be read with an accuracy of 0.001 cm.³ The displacement of the mercury is effected by raising and lowering a level-glass with mercury (*o*), which glass communicates with the capillary tube.

The bent tube, which connects vessel and calibrated tube, contains also a three-way stopcock (*p*) connected with a side-tube (*q*). This tube (*q*) serves to drive gas through it, but was only

used in the experiments for the determination of the assimilatory-quotient and is further unnecessary.

The vessels are put on tripods (*r*), so that the water of the thermostat basin, in which the apparatus is put may freely circulate around them.

The manometer (*c*) is a U-shaped capillary tube, 0.5 mm wide, open at the top, and has a capillary connection (*s*) with the vessels. The opening at the top can be closed with rubber tubing and a clamping-screw (*t*).

As a manometer liquid paraffin oil coloured red with Sudan III is used. Behind the tube is a scale division on millimeter-paper.

2. Fixation of the algal film on the glass cover. The

algal film is fixed on the inside of the glass cover (*f*) in the following way: A plate altogether free from grease is laid in a somewhat larger glass basin. This basin is filled with nutrient solution, so that the plate is a few millimeters below the surface of the liquid. On the water floats a paraffined paper ring (diam. \pm 8 cm), within which a quantity of algae is transferred. The basin is put on the window-sill under a glass cover. After two or three days a round homogeneous film has formed itself, which is suitable for the experiment. The liquid is sucked away with a pipette, the film is pushed towards the middle of the plate, where it stays as a humid layer. The last liquid is removed by means of filter-paper and the paper ring taken away. The plate is placed upside down on the assimilation-vessel, after which this is shut. The film should not be too humid, as drops would form making contact with the buffer-mixture, neither may it be too dry lest the film should dry up. After some practice this is easily performed. The vessel must also be placed strictly horizontal with the aid of a water-level in order to prevent the liquid from sinking to one side.

When unicellular algae serve as object, the cell-suspension is first centrifuged for some time; the clear liquid is poured off and the remaining thick suspension is removed to a clean glass plate with a brush.

3. *The regulation of temperature.* The contents of the waterbasin, in which the apparatus stands, must be kept at a very constant temperature, as the method is highly sensitive to fluctuations in temperature, on account of the great volume of gas of the vessels. A toluol thermostat is practically not sufficient. I preferred to regulate the temperature by hand, as a reading had to take place every five minutes, and I had an opportunity to control the temperature at that moment. Under the water-basin there is a

micro-burner (*u*) for heating purposes, while, in the basin, bent glass tubes (*v*), through which tapwater or ice-water streams, effect the cooling. By regulating the flame and the speed of the stream of cooling-water the temperature of the basin can be controlled very accurately. Slight fluctuations were compensated by keeping at hand little basins of warm and cold water or pieces of ice. The temperature is read from a thermometer with a scale division to 0.1° . The capacity of the basin is 70 litres. This regulation of temperature, which may seem to be primitive, allows an accuracy up to about 0.01° without any trouble, as within a short time a great skill is obtained. In the basin is a stirring-apparatus (*w*) driven by an electro-motor.

4. *The regulation of CO_2 -concentration.* In most cases a solution of the following composition was used as a CO_2 -buffer:

Na HCO_3	0.375 mol.
$\text{Na}_2 \text{CO}_3$	0.125 mol.

Such a solution is at 16° balanced with air containing 0.7 % CO_2 , at 23° with 1.1 %, which was determined by means of the CO_2 generator of van den Honert and the gas analysis apparatus of Krogh. With this concentration CO_2 is no more limiting; alteration of the CO_2 concentration from 0.2 % to 1.0 % did not give noticeable alteration of the CO_2 assimilation under the conditions of my experiments, as appeared from preliminary experiments.

5. *Illumination.* The light source was an ordinary 150 Watt Philips metal filament lamp (*x*), when experiments were made with the light-factor in maximum. At about 20° an alteration in the distance of the glowing filament of the lamp from 15 to 12 centimeters has no result at all. As a rule the distance was 13 or 14 cm. The lamp hangs halfway in the water; so between lamp and algae there is nothing else than a layer of water from 8 to 9 cm thickness.

When the influence of the light-factor was examined, a different arrangement was used. As a lightsource a Leybold projector lantern was used, in which, instead of carbon-points a Philips 1000 W. cinema-lamp was placed. The cone of light is vertically thrown on the assimilation-vessel with the aid of a 45° slanting mirror, while care should be taken that the axis of the cone falls on the centre of the vessel or close to it, so that the whole algal film is equally illuminated. In order to secure a uniform dimming of all wave-lengths the light of the lantern is weakened by means of so-called neutral light-filters (Plotnikov). These filters are made of fine mesh brass wire supplied to us by the Newark Wire Cloth Company, Newark, U.S.A. Other screens sold by the firm of F. Köhler, Leipzig, proved to be useless for my purposes.

The screens were coppered. Then they were hung on a copper wire in a 5 % solution of NaOH, to which 1 % potassiumpersulphate powder was added. After about 5 minutes the surface was dull black from the CuO formed. The screens were washed in water, dried with a soft piece of cloth and clasped between metal rings. The diameter of the screens is 10 cm. The following gauges were used:

mesh	diam. of wire (inches)	size of opening
24 × 24	0.0100	0.0317
30 × 30	0.0100	0.0233
35 × 35	0.0080	0.0206
40 × 40	0.0090	0.0160
45 × 45	0.0090	0.0132
50 × 50	0.0090	0.0110
55 × 55	0.0090	0.0092
60 × 60	0.0090	0.0077

The quantity of light which these screens allow to pass was determined by means of a Moll-thermo-element connected with a Moll-galvanometer and illuminating the element alternatively with or without a filter put before it. The relation of the galvanometer deflections gave the

percentage of light transmitted (with the filters used it varied between 32 and 61 %).

By combining two filters a still lower intensity of light can be obtained. Here great attention should be paid that the screens are parallel to one another, and that their wires do not run parallel, but form a certain angle (Plotnikov). If this precaution is not taken, moiré-phenomena may be obtained and consequently a very irregular illumination. It appeared for this reason that the screens could be placed in one place only, viz. between the two sets of lenses of the Leybold projector lantern, in close proximity of the front lens. Images or shadows of the screens appeared when the screen was mounted in any other place.

A plane parallel cuvet filled with water is placed between the two lenses, 10 cm in diameter measuring.

The vessels are submerged in the water basin to a depth of 6 cm below the surface (in the experiments with the Leybold lantern).

The distance of the top of the light cone, which falls between the two lenses, to the alga is ± 95 cm.

The voltage fluctuated during my experiments between 218 and 225 V.; very often it was much more constant. The inconstant voltage is without any importance, when the work is done with a strong intensity of light; in case of weak intensity a correction must be made (cf. the experiments on the influence of light intensity, Ch. VI).

When measuring the respiration all light was shut off by means of an opaque round plate covering the glass plate.

D. Course of an Experiment and Corrections.

After closing the vessels, filling the water-basin, turning on the screw clamp above the manometer, the experiment can begin. We must wait some time till the equilibrium has been obtained. Any air-bubbles which there might

be under the vessels are removed with a bent pipette. It takes rather a long time before equilibrium has become established, at least 30 minutes. This is probably caused by the changes of temperature and by the changes of CO_2 - and water-vapour tension. The absorption of light heats the vessel a little. The heating effect is about $\frac{1}{6}^\circ$, calculated from the expansion of the gas, when the lamp is at 13 cm distance.

Now the manometer is put at 0 and the level of the mercury in the calibrated tube is read. As a rule both manipulations are repeated every five minutes. We wait till a row of from 5 to 7 practically equal values have been read; the evolution of O_2 is apparently constant now, and the assimilation per hour can be calculated. In case of low intensity of assimilation or when measuring respiration, the experiment lasts longer in order to decrease the chance of a faulty reading.

The quantity measured does not precisely correspond to the volume of gas evolved, as part of the oxygen dissolves in the buffer-liquid, and nitrogen escapes from it. Moreover, the newly formed oxygen mixes with escaping carbon dioxide and water vapour. The increase of the oxygen tension is not quite proportional to the formation of oxygen, because the total volume of gas becomes larger. This may practically be neglected, since the oxygen tension increases only by a few percents during the experiment. The dissolved quantity may therefore be taken as proportional to the volume formed. Since the absolute value of assimilatory CO_2 is of secondary importance to us, a correction is not necessary. The same may be said of the carbon dioxide and the water vapour. Consequently the measured amount of assimilation is somewhat too small, the measured amount of respiration too large.

The above remarks about the corrections do not pertain, when the investigations are made at various temperatures.

The calculation of the corrections in that case was done as follows.

The vessel, in which the first experiments were made (table 6), contained 37 cm³ of gas and 104 cm³ of liquid. When 370 mm² of gas had been formed, the tension of the oxygen had increased by 1 %, that of the nitrogen decreased by 1 %. Now oxygen dissolves in the buffer-mixture, and a little nitrogen escapes. (We consider the solubility of the gases in buffer-mixture to be equal to the solubility of gases in pure water). The solubility of oxygen is greater, consequently more gas dissolves than is given off. The higher the temperature the smaller this difference will be. At 10° the solubility of the two gases is 0.03802 and 0.01857, the difference 0.01945; at 20°: 0.03102 and 0.01542, the difference 0.01560. The second difference is 0.00385 smaller than the first, i.e. when the volume of gas increases by 1 %, per cubic centimetre of liquid 0.0385 mm³ less gas dissolves at 20° than at 10°. The total liquid contains, consequently, $104 \times 0.0385 = 4.004$ mm³, less at 20°, or in percent. $\frac{4.00}{3.70} = 1.1$ %. So, relatively, the assimilation at 20° has been determined too high, and this amount must be deducted.

CO₂ and water vapour mix with this newly formed volume of oxygen. The tension of both gases is greater at the highest temperature. At 16°C the tension of CO₂ appeared to be 0.7 %, at 23°: 1.1 %. The difference in tension between 10° and 20° is 0.5 %; roughly estimated. The tension of water vapour is 9.1 mm and 17.4 mm at both temperatures; a difference of 8.3 mm or in per cent of the total tension $\frac{8.3}{7.6} = 1.1$ %. These amounts too must be deducted from the values found at 20°, so in total 1.1 + 0.5 + 1.1 = 2.7 %. The Q₁₀ of photosynthesis found (2.30, cf table 7) will be corrected 2.30 — 0.06 = 2.24.

In case of the respiration this correction has to be applied in opposite direction. The same amount must be added to the value found. But these determinations (table 9; and those of assimilation in May 1931, table 8) were made in smaller vessels, containing 20 cm³ of liquid and 32 cm³ of gas. The correction for the solubility of the gases is, therefore, about five times smaller, or 0.2 %. The correction for the CO₂ tension was 0.5 %, roughly estimated. In some experiments this tension was zero. Therefore, we put this correction at 0.3 % on an average. Then the total correction becomes 0.2 + 0.3 + 1.1 =

1.6 %. The Q_{10} of respiration found (2.06, cf table 10) will be corrected $2.06 + 0.03 = 2.09$.

These calculations make clear that the corrections are of no great importance.

Another correction to be made, when the influence of temperature is studied, is the one for the expansion of the gas in case of a rise of temperature. The volume is reduced to the lowest temperature of the experimental series, by deducing $\frac{1}{273}$ of this volume per degree of rise.

No further corrections of the apparatus seemed necessary.

According to van den Honert a correction must be made for the increase of the material, a „growth-correction”, which should be about 1 % per hour at 20°. I found a small increase indeed, which however, as van den Honert himself already stated was rather subject to variations. It appeared e.g. to be a little stronger in weak light, and so it seems that it is not exclusively founded on growth. Moreover I was faced by the difficulty, how to execute the correction at other temperatures. Since the duration of my experiments was on the whole shorter as those of van den Honert, it seemed warranted to drop the whole correction, which is moreover of little importance.

Operating with the apparatus as described above is very simple and not at all tiring for the investigator. An experiment hardly ever fails on account of technical difficulties. The accuracy of the method is, moreover, very great, the determination possible with an accuracy within a few cubic millimeters. The course of the process can always be followed, so that the course of an experiment may be changed if it seems advisable. Moreover, incidental phenomena can readily be recognized as such. Besides the advantage of the method of van den Honert, the feature of the thin film of algae, has been preserved.

Changes in the solution of the cell-layer may be easily effected in the following simple manner: the glass plate is again placed in a basin, a paraffined paper ring is laid round the film and liquid is added (containing the factor studied e.g. a narcotic). The films are so tight that this manipulation can take place without loss of material.

CHAPTER III.

The Respiration During the Assimilation.

A. Introduction. On good grounds it has always been supposed that respiration goes on during assimilation. It appears that different processes as e.g. protoplasmic current keep going on. These processes cannot be imagined without respiration.

Whenever the amount of the assimilation is determined, the amount of the respiration must be added to the value measured, in order to evaluate the exact amount of photosynthesis. For want of a better standard the respiration is here determined in the dark before or after the determination of assimilation, under the supposition that the difference of respiratory activity in the light and in the dark would be slight.

Yet there are other possibilities, viz. that the respiration would have been changed by the supply of products of photosynthesis or by direct action of light on the protoplasm, or by an increase in temperature. The first possibilities were mentioned by Meijer and Deleano (1911, 1913) and called the „ergastogeneus” action, and the „plasmo-geneus” action of the light. The third may especially be of interest in experiments made with leaves. Blackman and Miss Matthaei (1905) found in sunlight temperature increases from 4° to 12°, which large differences, however, were later contradicted by Willstätter and Stoll (1918), Lundegårdh (1924) and Johansson (1926), who could

find but small differences of a few degrees at most, when the light passed water, and the transpiration of the leaves was not hindered.

That an „ergastogeneus” influence of light exists, is fairly certain. Borodin ascertained (1876, 81) that the respiration is stronger after an exposure to light than before. Aereboe (1893) confirmed this. These investigators put this down to an increase of products of photosynthesis. An increase after an exposure, following a long period of darkness, was further observed by Matthaei (1904), Meijer and Deleano (1911, 13), Kniep (1914), Pantanelli (1915), Harder (1915), Plaetzer (1917), Warburg and Negelein (1922), Spoehr and Mac Gee (1923). The last-mentioned authors were able to show that the increase of sugar content runs parallel to this. Warburg proved that an addition of glucose stimulates the respiration, which fact could be confirmed by Emerson (1927), Genevois (1927) and by myself (Chapter X).

Meyer and Deleano accentuated that these experiments do not prove that at the same time light could not exercise a plasmogeneus influence. Experiments made in air free from CO_2 , which had to solve this point, gave doubtful results. On the other hand these investigators did succeed in finding a plasmogeneus action of light on the respiration of another nature. By means of alternate exposures (in the day, time light, at night dark) a rhythmic respiration was obtained. The rhythm continued when the leaf remained in the dark afterwards.

Miss Plaetzer noticed that these rhythmic phenomena did not appear in *Cladophora*. With *Spirogyra* on the other hand, she found a rise at night, probably in connection with nuclear division.

The experiments on the influence of illumination on the respiration of non-chlorophyllaceous parts of plants give little reason to believe in an important „plasmogeneus” influence

of light in case of green plants. The numerous investigations yielded few results, sometimes an increase, but mostly a slight decrease or no change was found. A slight increase may, moreover, have been caused by a rise of temperature (Löwischin, 1908). A discussion of the literature on this subject is to be found with de Boer (1928).

A better insight into this problem is the more desirable if one considers that sometimes the assimilation is relatively slight and only little larger than the respiration. Here the ignorance about the exact amount of respiration might be a dangerous source of errors.

B. Experiments. In the first place the intensity of respiration was determined before and after a period of illumination in which assimilation took place. The result




Fig. 2. Rate of respiration determined before and after a period of assimilation. The full lines represent the experimental values, the broken line the probable course of respiration.

of this experiment, graphically represented in fig. 2, shows clearly that the respiration has increased markedly (from 13.5 mm³ to 27 per hour), to return to the old value after some time. Several experiments yielded the same result as the experiment re-

presented in the figure. More examples of this phenomenon are the experiments made in June 1931; see table 22 in Chapter XII. The respiration is very high after a long period of illumination by daylight; it keeps decreasing during the determinations of respiration in darkness.

The result of the experiment, mentioned above, indicates that the respiration during the exposure to light had been at least twice as great as in the period before it. It stands to reason that this experiment does not show, whether this increase is of an ergastogeneus or plasmogeneus nature, according to the conception of Meyer and Deleano.




Fig. 3. Increase of respiration in a CO_2 -free vessel after exposure to light. Full lines respiration in the dark (experimental values); ——— lines respiration in the light diminished with the assimilation of respiratory CO_2 (exp. values); the broken line represents the probable course of respiration before, during, and after exposure. In spite of the lack of CO_2 (no assimilation!) there is a remarkable increase of respiration during and after the exposure.

of the influence of exposure (lamp 150 W. at 13 cm). An increase of respiration may no doubt be concluded.

This experiment shows furthermore the remarkable feature that the respiration could also be determined during the exposure. The amount that is determined now is of course not the pure respiration, for part of the carbon dioxide formed by the respiration (x) will be assimilated once more, after which O_2 evolves. The amount determined is, therefore, the respiration — x , the real respiration in this experiment amounted consequently to $51 + x$ per hour. This is already higher than the rate of respiration

nature, according to the conception of Meyer and Deleano.

In order to establish this the respiration was examined in a CO_2 -free vessel over a solution of 1 % KOH, instead of over a buffer-mixture containing CO_2 . The following experiment made with *Oocystis spec.*, graphically represented in fig. 3, gives an impression

before in the dark (39). For the rest this experiment shows the phenomenon much less beautifully than I always found it with *Hormidium*. I chose this experiment as an example, as the increase after the exposure is shown here very markedly.




Fig. 4. Rate of respiration before, during, and after exposure to light in a CO_2 -free vessel. Full lines respiration in the dark (exp. values); ——— lines respiration in the light diminished with the assimilation of respiratory CO_2 (exp. values); the broken line shows the probable course of respiration.

The amount of respiratory CO_2 which is assimilated again is unknown. Therefore the intensity of respiration in the light must be higher than is shown by the experimental values (———).

A very beautiful example is given by e.g. experiment 5, made with a *Hormidium* strain of Pringsheim grown in constant light (fig. 4).

During the exposure the respiration rises from 39.5 to $87 + x$, so the increase is at least more than 100 % here. For the sake of control, the alteration in volume was also measured, after the equilibrium of temperature being re-established in the dark. This determination expresses the respiration during the time of exposure followed by 10 minutes of darkness; this value, too, is very high ($80.5 + y$). The rise after exposure is small, but visible. Again the experiment was repeated, first in light of lower intensity, then in the same. In the light of low intensity the rise is much smaller, in strong light it was almost as large as before.

Other experiments yielded similar results.

The rise also takes place with the algae, cultivated on the windowsill; both with Pringsheim's and van den Honert's strain. A positive result was also obtained with a culture in constant light of *Stichococcus bacillaris*; increase from 40 to $48 + x$.

These experiments distinctly show that the respiration during the illumination (assimilation) may be much greater than one would expect from a determination after or before the measuring of the assimilation. When we can make it clear, that the increase is not due to a rise of temperature, then the plasmogeneous action of light has been proved. In the first place, it be once more mentioned here, it can be ascertained from the expansion of the gas, that the assimilation-vessel becomes $\pm \frac{1}{6}^{\circ}$ warmer. The general heating can consequently be neglected. Also the film, attached to a thin glass plate, can hardly be warmer.

There may exist an unequal temperature in the different parts of the cells, but the probability of this supposition is not very great. The increase of respiration was sometimes twice or three times as much, or more. As we shall see in Chapter VII the Q_{10} of respiration at higher temperatures is a little lower than 2. In order to explain such an increase

we should have to accept an average rise of temperature of 10° to 20° at the least. Now I noticed that owing to a short stay of 10 minutes in the dark at 41° the assimilation had afterwards considerably decreased at lower temperature. In the face of this, an almost constant assimilation at 34° , as we found in Chapter VII, would hardly be possible, if the alga had a temperature of from 44° to 54° on the average.

A second argument against a strong rise of temperature we meet in the light intensity-assimilation curve (Chapter VI). In strong light this curve runs parallel to the abscissa. Yet the assimilation in strong light is very sensitive to temperature (Chapter VII). It may be objected that the light intensity-assimilation curve in itself is possibly an optimum curve and that the fall may accidentally have been counterbalanced by the action of the temperature. This does not seem to be very likely, and so this parallel course proves (when this subtle counterargument will be relinquished), that no heating (worth mentioning) takes place in case of an increase of light intensity.

In the third place the influence of red light on the respiration has been examined in experiment 9, where a material increase of respiration from 12 to $33 + x$ was met with. The Wratten filter 71 A β was put on the vessel; this filter allowed about 64 % of the energy and only radiation $> 600 \text{ m}\mu$ to pass (determined with thermopile). Yet the respiration was only $15.5 + x'$, and, therefore, as x' will be about equal to x (because CO_2 concentration is the limiting factor), much less than in white light. This seems to disprove the influence of temperature.

In the last place the after-effect of light on respiration, which is distinctly ascertainable even one hour after the exposure (fig. 3 and 4), seems to disprove a rise of temperature.

By these considerations the plasmogenuous influence of light on the examined algae is made very plausible.

The objection may be made that the respiration is measured in air, free from CO_2 , and the rise is wholly or partly due to the lack of the CO_2 factor ¹⁾. For this reason two experiments mentioned in Chapter VI (exp. 29 and 30) are of great importance. In a vessel containing CO_2 there appeared to be no assimilation in strong light, probably because the assimilation of the cells (cultivated in weak light) was hindered by very strong light. During the exposure the respiration proved to be 2—4 times as great as in the dark.

I have also tried to ascertain whether light may cause an increase of respiration in white plants, because an influence of temperature will be smaller here, though the literature gave me little hope. Yet in experiment 10, a suspension of *Saccharomyces Vordermannii* appeared to increase the respiration from 152 to 183. White leaves of *Oplismenus* from the hothouse, stuck on the glass with strips of paper, gave the following results:

Exp. 11, 19—3—31; 5 leaves; temp. 20.00°.

Rate of respiration, in the dark: 49.5, 45;

” ” ” , in the light: 63 + x;

” ” ” , in the dark: 46 per hour.

Exp. 12, 7—7—31; many leaves; temp. 24.00°.

Rate of respiration, in the dark: 76, 62, 56;

” ” ” , in the light: 61 + x;

” ” ” , in the dark: 49, 48, per hour.

It is not certain that these leaves do not contain any chlorophyll. In that case the result can at most be unfa-

¹⁾ Noack (1925) stated an increase of O_2 absorption from leaves illuminated in absence of CO_2 after 84 hours, caused by photo-oxidative processes. Simultaneously the leaves grew yellow. These processes probably play no part in my experiments, as the cells proved to be quite normal after the experiment, and were able to assimilate normally.

vourably influenced. So these results also speak against heating.

When we closely consider the figures of the increase of respiration in glucose and fructose (Chapter X, table 20), it strikes us that this increase is only rather slight, usually no more than 50 %, often much less. So it is no exaggeration to say that the plasmogeneous action of light exceeds the influence of the ergastogeneous action by far.

The knowledge that there is a plasmogeneous action of light is, besides being important in itself, also significant for us in connection with the inquiry about the assimilation of CO_2 . In what follows we shall many times have to deal with the question, whether light (or other external factors) controls the intensity of assimilation directly as a factor in the photosynthetic process itself, or whether the action is indirect and is caused by the intermediary of the protoplasm. It goes without saying that the latter conception becomes more likely, now that we have ascertained a very strong influence of light on protoplasm.

The fact that the respiration has increased during the assimilation of CO_2 is of practical importance, especially for the method of determining the assimilation. A minimum value has only been ascertained; an exact quantitative determination will very likely be among the impossibilities. When the rate of assimilation is very high the respiration may be neglected, without a serious objection. Matters change, however, when the rate of assimilation is low. Especially in strong light the error will be very large, and may give rise to altogether wrong conclusions, so that this knowledge warns us at least to be cautious when we interpret the results. In the literature it is easy to find examples which are, for this reason, of doubtful value. One example I will mention.

Lundegårdh (1924) has found very interesting optimum curves. He found e.g. an optimum curve when

studying the influence of light intensity in case of low CO_2 concentration. In these experiments the assimilation was rather small. The decrease of assimilation in strong light may now well be explained by a large increase of respiration. I do not mean to say that Lundegårdh cannot be right, but I will only point out how cautious we must be in the explanation of such a case.

Notwithstanding the fact that there is an increase of respiration in light, yet in my experiments I have stuck to the oldfashioned way of correcting the respiration. We had perhaps better multiply the values found by 1.5 or 2. But how does the respiration behave during an exposure with other intensities of light? We saw that, in strong light, there is a large increase, but how is the quantitative relation? I always determined the respiration after an exposure, except in the experiments in Chapter V, when I did not yet know this influence.

One remark should be made. The results recorded in the tables 15—23 of van den Honert and in table 2 of myself (experiments made with the van den Honert apparatus) seem at first to be little in conformity with those described above. A lower concentration of CO_2 is continually found in light than in darkness. The difference between the method of van den Honert and my own method can account for this discrepancy. In the apparatus of van den Honert a molecule CO_2 formed by respiration, and afterwards evolved, is repeatedly offered to the cells and will have a great chance of being assimilated again; in other words: x will be very great. In my apparatus, as used in the above experiments, it will directly be bound by the alkali. The unknown x in my results will possibly be not very great. Moreover the light intensity in the experiments of van den Honert is considerably lower, and consequently the increase of respiration rather slight (cf. fig. 4).

CHAPTER IV.

The Assimilatory Quotient.

By assimilatory quotient I mean, with Stiles (1925), the quotient of the volume O_2 evolved and the absorbed CO_2 , in contrast with Willstätter and Stoll (1918) who define the notion reversely. Stiles is quite right, when he observes that the reverse of a quotient, which is historically always used in this form, makes no sense.

The knowledge of this quotient was for me of importance for two reasons. In the first place because van den Honert (1928, 30) contrary to Maquenne and Demoussy (1913), Willstätter and Stoll (1918), and Johansson (1926), who had always found that the quotient approximates to unity, had noticed a deviating quotient of 1.1 with *Hormidium*. The second reason is that I would investigate to what extent my method, in which the assimilation of carbon dioxide is stated by measuring of the O_2 evolved, is in conformity with the method of van den Honert, based on CO_2 determinations.

In order to determine the quotient the assimilation vessel is thoroughly cleaned from CO_2 buffer. In the vessel there is only air containing carbon dioxide and the algal film. The CO_2 concentration is increased by expired air blown, after deep breathing, through the tube *q* (Fig. 1) which has been inserted only for this purpose in the apparatus. Then follows an exposure to light. If the volumes of gas evolved and absorbed should be equal, the manometer liquid will have to remain at rest. If, on the other hand CO_2 absorption or O_2 evolution predominates, a pressure difference will arise. When this difference has been determined, the vessel is opened and CO_2 buffer is added, after which the assimilation can be measured, i.e. the development of O_2 . We now add to this

value the difference stated before and find the CO_2 absorption. In this way the quotient has been determined four times, cf. table 1. The respiratory quotient, which is perhaps very deviating from unity is not taken into account, which is admissible as the respiration is slight in proportion to the assimilation.

TABLE 1.
Determination of the assimilatory quotient.

Number of experiment	Date	Strain	Temperature	O_2 evolution per hour	Difference between O_2 evolution and CO_2 absorption per hour	Assimilatory Quotient O_2/CO_2
13	11-9-30	Pringsheim's	22.00	192	— 18	0.92
14	16-9-30	"	22.00	1068	+ 66	1.07
15	19-9-30	"	22.00	957	— 6	0.99
16	10-7-31	v. d. Honert's	21.50	610	+ 28	1.05

The quotient found deviates only little from unity. Van den Honert found 1.09 and 1.13, amounts that are not very unlikely. Yet there seems to me to be no reason for the opinion of van den Honert, that there should be a connection between the high assimilatory quotient and the synthesis of oil as first product of assimilation. Moreover, I never succeeded in demonstrating fat in the *Hormidium* cell either with osmic acid or Sudan III. Pascher (1914) who describes the formation of reserve material in detail, tells nothing about such an oil synthesis.

The second conclusion that may be derived from these experiments, is that it is immaterial, whether the assimilation of CO_2 is determined by measuring the CO_2 -absorption or the O_2 -evolution.

CHAPTER V.

The Influence of Carbon Dioxide Concentration on Assimilation ¹⁾.A. The CO_2 Concentration-Assimilation Curve.

Van den Honert (1928, 30) investigated the influence of CO_2 concentration on assimilation. His results deviated remarkably from those of other investigators. When the CO_2 concentration was low the assimilation rose directly proportionally to the tension of CO_2 , when it was high it was altogether independent of it. Between the two regions




Fig. 5. Relation between CO_2 concentration and photosynthesis in Hormidium.

there was a small transition. The curve of van den Honert therefore approximates the Blackman-scheme.

I have repeated this experiment with the pure Hormidium strain of E. G. Pringsheim. All corrections were carried

¹⁾ The experiments of this chapter were all made with the van den Honert apparatus.

out as by van den Honert. The constant rate of assimilation at high CO_2 concentration has been put at 100. The illumination was effected by 6 lamps as used by van den Honert in a row, with a reflector of white paper behind. The latter was absent in van den Honert's experiments; the intensity of light was therefore somewhat higher.

Figure 5 gives the graphical representation of the experimental result. The curve perfectly agrees with the one found by van den Honert.

Van den Honert has been able to make it probable that the direct proportionality in low CO_2 concentration arises here, because a diffusion process controls the whole photosynthetic process, and that in case of high CO_2 concentration a chemical process is limiting. In the latter case a rise of temperature gives a considerable acceleration, in the former it remains without any influence. This is, as van den Honert explained, in conformity with the properties of the (purely physical) CO_2 diffusion, taking into account the solubility of the gas at different temperatures.

According to van den Honert the sharp break in the curve is caused by a rapid fixation of the CO_2 by the organism.

The curve found with *Hormidium* deviates altogether from those, determined by other investigators, e.g. from Warburg's curve, representing the relation between CO_2 concentration and assimilation in *Chlorella*. This may be caused by the fact that, in the experiments of Warburg, the diffusion of CO_2 was no limiting factor. At least he found a great influence of temperature in low CO_2 concentrations.

Finally it may be observed here, that it also appeared from the preliminary experiments with the manometric method that in case of high CO_2 concentration (0.1—1.0 %)

the assimilation is quite independent of the concentration. No optimum curve has been found, as has been observed by Lundegårdh (1921, 24) for higher plants.

B. The Influence of Carbon Dioxide Concentration when the Assimilation has been Retarded by Narcotics.

It is a well-known fact that the assimilation may be depressed by the addition of narcotics. We may ask whether, by means of a suchlike retardation, we cannot eliminate the interference of the diffusion-process, and in this way measure the influence of CO_2 concentration on the assimilatory process itself.

For this purpose I chose a concentration of phenylurethane, which according to Warburg retards the assimilation: $2.0-2.4 \times 10^{-4}$ mol. Moreover I made some experiments with antipyrin, examined by Ewart (1896) and Jacobi (1899) for its retarding action; I experimented with a weak concentration, viz. ± 0.001 mol.

TABLE 2.

The influence of CO_2 concentration on photosynthesis after the addition of a narcotic.

Exp. 18, 19-3-30. Pringsheim's strain. phenylurethane conc. 2.2×10^{-4} mol.

Time	Light	Assimilation in mm^3 per hour	Corrected CO_2 conc. in 0.001 %	Corrected assimilation
11.50	darkness	-29	—	—
14.25	strong light	95	135	92
16.00	"	105	173	100
17.10	"	76.5	50.5	72
19.00	"	64	37	59
20.45	"	75	40.5	68
22.00	"	114	168	102

Exp. 19, 27-3-30. Pringsheim's strain. Phenylurethane conc.

 2.0×10^{-4} mol.

Time	Light	Assimilation in mm ³ per hour	Corrected CO ₂ conc. in 0.001 %	Corrected assimilation
10.45	darkness	-20.5	—	—
13.45	strong light	108	100	—
15.10	"	146.5	136	90
16.15	"	121	55	74
17.20	"	97.5	19	59
19.15	"	65	10.5	39
20.50	"	167.5	367	100
22.00	"	152	132	90

Exp. 20, 3-4-30. Pringsheim's strain. Phenylurethane conc.

 2.2×10^{-4} mol.

10.30	darkness	-27	—	—
15.40	strong light	109	16.5	64
16.40	"	135	51	79
19.30	"	182	328	100
21.00	"	131	26	73
22.45	"	86	12	47

Exp. 21, 24-5-30. Pringsheim's strain. Antipyrin conc.

0.001 mol.

11.40	darkness	-33.5	—	—
12.55	strong light	188	368	100
14.10	"	149	145	80
15.10	"	122	64	65
16.20	"	110	22	58
19.45	"	200	374	100
20.50	darkness	-40	—	—
22.30	strong light	85	11	43

Exp. 22, 3-6-30. Van den Honert's strain. Antipyrin conc.

 $\frac{1}{1300}$ mol.

12.20	darkness	-30	—	—
13.45	strong light	87	76.5	73
15.00	"	125	414	100
16.20	"	70	40	58

Table 2 shows the results of these experiments. It strikes us that the second observation in experiment 19 has yielded a rather low value. This is in accordance with the further experiments made with phenylurethane (Chapter IX), where it has been shown that the retardation is stronger at the outset, and diminishes in course of time. The rate of assimilation becomes rather constant some hours after the addition of the narcotic. All other observations fall in this constant region. The rate of assimilation in high concentration of CO_2 is again arbitrarily put as 100. Here we must, however, not forget that this 100 cannot be compared with the 100 of normal assimilation, as in higher CO_2 concentration the assimilation is retarded as well as in low concentration.




Fig. 6. Relation between CO_2 concentration and photosynthesis in *Hormidium* when phenylurethane has been added to the algal film. The „Blackman-type” of the curve shown in fig. 5 has entirely disappeared. There is a striking resemblance between this curve and the curve published by Warburg, showing the relation between CO_2 concentration and photosynthesis in normal *Chlorella* (see fig. 7).

When we represent the result of exp. 19 graphically (fig. 6), a curve is obtained which deviates very considerably from the CO_2 concentration-assimilation curve found before. The „Blackman-type” has disappeared, the transition is very gradual. The rate of assimilation increases in case of high CO_2 concentrations where normally no increase takes place.

The curve very strongly calls to mind the curve of Warburg, represented in fig. 7. In order to make com-

parison easier the concentration of CO_2 is also given in 0.001 %. It seems not unlikely that the same part of the process has been studied in both cases, possibly the chemical binding of the CO_2 .




Fig. 7. Relation between CO_2 concentration (expressed in 0.001 %) and photosynthesis in *Chlorella*, according to Warburg. The maximum rate of assimilation has been put at 100.

Van den Honert is of opinion that a curve as I obtained in fig. 6 (represented in his publications in fig. 17), represents the quantity of the CO_2 -assimilatory agent (probably CO_2 -chlorophyll) at different CO_2 concentrations, and at the same time the rate of the limiting process. It is true that he speaks only of the photochemical part of the process, but from what follows it appears that this is presumably a slight error, and that the limiting process is meant, which may be chemical as well as photochemical. He thinks that the determination of this curve will enable us to find the sequence of the different parts of the chain process. If, he argues, a lowering of the CO_2 concentration should cause a retardation, when light were the limiting factor, then CO_2 would be a factor reacting in the photochemical process, whereas, should this lowering be of influence, when temperature were limiting, CO_2 would be a factor in the chemical process.

It will be seen below that the reaction is very sensitive to temperature. A lowering of the CO_2 tension acts, therefore, retarding, when a chemical process is limiting. When we apply van den Honert's argument to my result, CO_2 would be a factor in the chemical process and this consequently would precede the photochemical one. Now, this would refute the theory of Willstätter and Stoll and agree with a theory as formulated by Warburg.

In my opinion this conclusion goes too far. It is certain that the chemical reaction is retarded by a lowering of the CO_2 concentration. But from this it does not follow that CO_2 itself takes part in this reaction. Let us imagine that a photochemical conversion of CO_2 precedes. This reaction is not limiting, (in the case considered here) goes „faster” than the other processes, and the product of this reaction will be dependent on the quantity of the CO_2 only (it being indifferent for my considerations if this CO_2 is free or bound to another substance, for instance to chlorophyll). The concentration of the latter is dependent on the CO_2 -supply, so that the same will, therefore, hold good for the quantity of substance that has been converted photochemically. In this way the chemical reaction could consequently be dependent on the CO_2 concentration, even if CO_2 itself does not react in this chemical reaction.

For that reason the problem of the sequence of the processes must as yet in my opinion remain unsolved.

I will also point out that it is not necessary that the chemical reaction, which comes to the foreground here, is the same as the chemical reaction that is limiting in a normal organism when temperature is the limiting factor. Different parts of the chemical reaction complex may be unequally sensitive to these chemicals.

C. *The Influence of Temperature on the CO_2 Concentration-Assimilation Curve when Phenylurethane has been Added.*

In order to answer the question whether the reaction which controls the process after the addition of narcotics is sensitive to temperature, I have determined this CO_2 concentration-assimilation curve at two temperatures. Now that we know the general trend of the curve the fixing of a few points suffices.

The results of both experiments have been recorded in table 3. In exp. 23 the temperatures 19.2° and 22.8° have been compared and in exp. 24 the temperatures 19.2° and 22.3° .

The few figures point to a strong increase at higher temperature. In later experiments (Chapter IX) it has been shown that the retardation, strong at the beginning, becomes less in course of time. Owing to this the values at the lower temperature have been found a little too low. Yet the third determination in exp. 23 falls already $5\frac{1}{2}$ hours after the beginning of the narcosis, the much higher 7th value must, therefore, for the greater part be explained by the rise in temperature, as the retardation is fairly well constant after 5 hours (cf. exp. 71 and 72).

We may, therefore, conclude that temperature is of great influence.

TABLE 3.

The influence of the CO_2 concentration after the addition of phenylurethane at different temperatures.

Exp. 23, 14-4-1930. Van den Honert's strain. Phenylurethane-concentration 2.2×10^{-4} mol.

Time	Temperature	Light	CO_2 concentration	Assimilation
11.05	19.2°	darkness	—	-43
12.20	"	strong light	425	124
15.10	"	"	90	87
16.50	22.8°	darkness	—	-62
18.00	"	strong light	397	193
20.10	"	"	154	159
21.15	"	"	71	137

Exp. 24, 14-5-1930. Van den Honert's strain. Phenylurethane-concentration 2.2×10^{-4} mol.

11.40	19.2	darkness	—	-29
14.50	"	strong light	174	90
16.15	22.3	darkness	—	-34
17.20	"	strong light	158	125
19.25	"	"	405	147

CHAPTER VI.

The Influence of Light Intensity on Assimilation.

A. Introduction.

Already many investigators have examined the influence of light intensity on CO_2 assimilation. This relation, represented graphically, usually yielded a curve of logarithmic type. Higher land plants and aquatics are unsuitable for this work as the light intensity will rapidly diminish in the leaf. Such experiments do not prove that, if the illumination had been ideal, no curve agreeing with the Blackman-scheme, would have appeared. Boysen Jensen and Müller (1929) lately drew the attention once more to this fact. This objection also perhaps pertains

to Harder's experiments (1921) with *Fontinalis*. The same criticism was made by van den Honert (1928, 30) against the experiments of Warburg (1919), in which thin cell-suspensions were employed, weakening the light only 10–20 %, but in which some cells would, however, be much less illuminated. The objection is perhaps not a serious one as the average diminution of the light is only 0 to about 15 %. Moreover the cells that receive less light, will be relatively small in number and exercise little influence on the total result. Van den Honert has avoided this objection by using an algal film of one cell thickness where only the unequal illumination of light-and shade-side remains. Essentially the curve of van den Honert does not deviate from that of other investigators. However I thought it desirable to reinvestigate the problem, as an objection may be raised against the way in which van den Honert illuminated the algae.

He placed 1–6 incandescent lamps in a row. Using the same arrangement I determined the distribution of light intensity in the assimilation vessel of the apparatus of van den Honert (with the aid of a thermopile). Taking 10 for the light intensity in the middle of the vessel, and placing at first one lamp before the middle of that vessel, the intensity at 8 cm distance from the middle appears to be only 5. This is somewhat more unfavourable than in the experiments of van den Honert, in which, only a central part of ± 11 cm in length is used. The distribution of light in the vessel shows to be unequal. Now when we put 6 lamps in a row the intensity in the middle of the vessel is about 40, and at 8 cm distance from the centre only 30.

For these reasons the illumination used proved to be improper.

In my experiments I used the arrangement described in Chapter II.

Equal distribution of the light intensity is attained by throwing the centre of the light cone of a Leybold projector lantern on the algal film. The variation of light intensity is obtained by placing wire cloth screens in the

bundle of light. The light intensity of the undimmed projection lamp is arbitrarily put as 1000. As a very strong light I used an ordinary 150 W. Philipslamp; the intensity of it at a distance of 14 cm agrees with about 2300 (1000 ± 3000 Lux, roughly determined with a photometer). The slight fluctuations in the voltage were measured with a voltmeter. According to Holst (1920) a correction of 3.5 % for the light intensity must be made for each per cent variation of voltage.

B. Experiments on the Influence of Light Intensity on the Assimilation of Hormidium Cultivated in Daylight.

The experimental results are recorded in table 4.

We find the result of exp. 27 graphically represented in fig. 8. Evidently a curve of a logarithmic type has been obtained, notwithstanding the improved arrangement of

TABLE 4.

The influence of light intensity on photosynthesis.

Exp. 25, 10-3-31. Van den Honert's strain. Temp. 22.00°.

Time	Light intensity			Assimilation
	Transmitted	Voltage	Corrected	
11.30	1000	219½	952	171
12.45	611	222¼	609	136
14.15	415	222¼	413	102
15.15	254	222½	254	66
16.20	2300	—	2300	211
17.05	1000	222¾	1004	171
18.20	0	—	—	-35

Exp. 26, 13-3-31. Pringsheim's strain. Temp. 20.00°.

10.25	1000	221¼	988	493
11.05	611	222	611	351
11.50	415	222	415	271
12.55	254	222½	256	181
14.00	174	222	174	134
15.30	0	—	0	-35

Exp. 27, 16-3-31, (fig. 8). Pringsheim's strain. Temp. 18.00°.

Time	Light intensity			Assimilation
	Transmitted	Voltage	Corrected	
11.00	611	220 $\frac{1}{4}$	613	350
11.40	335	220	335	213
12.35	174	220 $\frac{1}{2}$	175	120
13.40	1000	221 $\frac{1}{2}$	1024	465
14.25	2300	—	2300	539
15.30	0	—	0	-26

the experiment. The shape of the curve corresponds to those of Warburg (1919), Harder (1921), Bose (1924) a.o. An approximation of the Blackman-scheme is apparently out of the question.




Fig. 8. Relation between light intensity and rate of assimilation in an algal film cultivated in daylight. Light intensity „1000” = \pm 3000 Lux. Though the experiment was carried out under improved experimental conditions, the curve does not show the „Blackman-type”.

C. Experiments on the Influence of Light Intensity on the Assimilation of *Hormidium* Cultivated in Weak Artificial Light.

The way of cultivation is of great influence on the shape of the curve. Experiment 28 has been made with a cell-

suspension, cultivated near a lamp of low intensity of light, burning continually. The distinguishing feature of this culture appeared to be that the assimilation optimum was reached at a much lower light intensity (611), which appears very distinctly from fig. 9. The alga has adapted itself to weak light, it has become a „shade plant”. The curve shows a weak „optimum type”. It is however not impossible that this is caused by an increase of respiration in the light of high intensity (cf. Chapter III).




Fig. 9. Relation between light intensity and photosynthesis in a suspension of *Hormidium* cells cultivated in artificial light of low intensity. The curve shows the „shade-plant type”. The optimum of photosynthesis is reached at a low light intensity („600”—„700” = ± 2000 Lux).

Two remarkable results, which also demonstrate the influence of the preliminary treatment, were obtained with an algal culture, cultivated in the same way as mentioned above, a young cell-suspension, 17 days old. In light of the 150 W. lamp at 15 cm distance not the slightest assimilation appeared to take place; on the contrary the O_2 absorption was greatly enhanced. This experiment was repeated with the same result on the same day.

On the next day the experiment was repeated with another cell-suspension and with a similar result. I verified the apparatus with an algal film from the windowsill, which proved to assimilate in a normal way. What was the cause?

I supposed that the high intensity of light had been injurious to the cells, and an assimilation optimum would exist at very low light intensity. In order to settle this point exp. 30 was continued and a weaker light applied (lamp at 40 cm distance). It was remarkable that the great O_2 absorption (-47) had totally disappeared and had passed into a very slight evolution of O_2 ($+8$). So the supposition proved to be correct.

This experiment is moreover a remarkable example of an increase of respiration in light, demonstrated in Chapter III. The absorption of oxygen had become at least from 2 to 4 times as great in the light.

I intended to determine the entire light-curve of this most interesting phenomenon. When, however, I wanted to repeat the experiment a month later, I could not get another result than that of figure 9, in spite of several attempts with old and young cultures. Without any doubt this diminishes the value of the discovery, and an explanation must be given with some restriction, though a mistake in the determinations is to be considered as out of the question.

The phenomenon calls to our mind the evolution of CO_2 so often found by Kostytschew and other Russian investigators, with favourable illumination, granting that the evolution or absorption of O_2 is determined in my experiments.

The above instance remained exceptional. In all other experiments I found a strong assimilation which continued to exist for hours at a stretch without any variation worth mentioning.

D. The Influence of Light Intensity on Assimilation at Different Temperatures.

Harder (1921) investigated the behaviour of the assimilation in case of a simultaneous change of the factors

light and carbon dioxide. He found that an increase of CO_2 also has got of influence when the light intensity is low and vice versa. A similar result was obtained by Lundegårdh (1921, 22).

We make a similar experiment for the factors light and temperature, when we determine the light intensity-curve at two different temperatures. The result, which we find in the graphical representation in fig. 10, is that in case of low light intensity, the influence of temperature is only slight. Another experiment gave a similar result.

It is certain that the curve never makes a much larger angle with the abscissa, when temperature is increased, as Harder had observed by varying the factors light and CO_2 . In this respect there is a greater agreement with the opinion of Blackman.

From both experiments it does not appear convincingly, whether there is still a small increase, or that it is totally absent. The question is for me of special interest, as it will appear from other experiments (Chapter VIII and Chapter X), that a stimulation of the assimilation in low light intensity is very well possible. Possibly the rate of assimilation is not in the first place determined by the quantity of light energy added to the assimilatory apparatus, but internal factors may also control the rate of photosynthesis, also when light is the external limiting factor. Seeing that




Fig. 10. Relation between light intensity and photosynthesis at different temperatures.

temperature will influence these internal protoplasmic processes, it is of importance to look for a possible influence of temperature in case of weak light.

Van den Honert has already made statements about this question. He found, however, a somewhat lower assimilation at higher temperatures.

Van den Honert thought he could prove by this, that the Q_{10} of the photochemical part of photosynthesis is ± 1 (Warburg is of the same opinion), in my opinion, however on insufficient grounds. If the velocity of the process should really be determined by the velocity of a photochemical reaction (let us suppose this to be the case), yet the sensitivity to temperature could not exercise any influence. At both temperatures an equal quantity of light energy is given, and in case all the absorbed light is photochemically at work, the coefficient of temperature found cannot be anything else but the quotient of the quantities of light supplied in both cases. In our instance this quotient will be 1, because the light intensities are equal.

Photochemistry is repeatedly faced by the same difficulty. Q_{10} must always be ascertained when the factor light is in maximum. It is then often an accompanying chemical process of which the Q_{10} is determined, but this difficulty cannot be overcome and the problem remains unsolved as yet.

As we stated before, it will appear in the following chapters that even when light intensity is the limiting factor, the intensity of assimilation may be varied through changes in the internal factors.

The supposition that *the photochemical stage of photosynthesis controls the entire process exclusively*, is therefore unwarranted.

Table 5 shows the results of a series of experiments, in which the problem is treated more accurately.

The voltage fluctuated a little. Therefore, the assimilation velocity is reduced to equal light intensity. The rate of assimilation is supposed to be proportional to the light intensity.

Experiment 33 seems to show a distinct increase at higher temperature. However, results of the experiments 34 and 35 are not so evident. But for a difference between the first and the second observation the assimilation is distinctly

constant. It rather seems that a kind of adaptation to the weak light takes place, than that temperature is of influence.

TABLE 5.

The influence of temperature on photosynthesis in light of low intensity.

Exp. 33, 22-5-31. Van den Honert's strain. Light intensity „153”.

Time	Temperature	Voltage	Apparent assimilation + respiration	Assimilation reduced to equal light intensity
11.35	10.00	220	93 + 17	110
13.55	16.00	221 ³ / ₄	95 + 30	122
16.15	22.00	220 ¹ / ₂	97 + 43	139
18.25	28.00	223 ¹ / ₂	95 + 59	146

Exp. 34, 27-5-31. Van den Honert's strain.
Light intensity „140”.

10.45	28.00	219	141 + 42	186
12.40	22.00	223 ¹ / ₄	178 + 32	200
14.50	16.00	221 ¹ / ₄	179 + 21	196
17.20	10.00	224	195 + 13 ¹⁾	195

Exp. 35, 3-6-31. Van den Honert's strain.
Light intensity „140”.

10.25	14.00	218 ³ / ₄	132 + 24	159
12.30	20.00	221	134 + 40	171
14.40	26.00	218 ³ / ₄	112 + 56	172
16.30	20.00	218 ¹ / ₄	128 + 40 ¹⁾	173
17.30	14.00	222 ¹ / ₄	166 + 19	179

Exp. 36, 2-6-31. Van den Honert's strain.
Light intensity „140”.

11.25	20.00	220 ¹ / ₂	94 + 28	121
13.25	20.00	221	106 + 28	132
15.45	20.00	221	106 + 28 ¹⁾	132

¹⁾ Estimated.

A similar little jump is also to be seen in experiment 36, where I experimented under constant temperature. Experiment 33 is distinguished by a repetition of this jump. Adaptation and temperature are working here in the same direction.

At all events the influence of temperature at low light intensity is small. Yet the fact that adaptation and temperature together cause a rather considerable effect, proves that the assimilation is not exclusively dependent on the external factor light, but that the extent to which the light energy is used, is also under the influence of internal factors.

CHAPTER VII.

The Influence of Temperature on Assimilation and Respiration.

A. *Introduction.*

The influence of the factor temperature on the assimilation, when neither light nor CO_2 are limiting, has also been repeatedly investigated. The result is often expressed in the coefficient Q_{10} . Some American investigators as Crozier, Navez, Emerson (their works are especially to be found in *The Journal of General Physiology*) violently oppose the use of this coefficient, which they wish to replace by the temperature-characteristic μ , the energy of activation. It is true that Q_{10} is not a constant; $\log Q_{10}$ is inversely proportionate to the product of the absolute temperatures. This seems no objection to me, as in biology only a rather limited range of temperature is studied, and a constant Q_{10} , therefore, fairly well corresponds with a constant μ . The conclusion they try to get from the study of these coefficients, viz. whether one or more processes control the rate of the total process studied, is therefore equally possible whether the results are expressed in Q_{10} or in μ . So there is no objection to

use Q_{10} , which remains a convenient symbol. A conversion in μ is moreover readily performed.

A danger of working with Q_{10} is that the temperature interval of 10° is rather great, and it does not state deviations, e. g. optima and minima. For these reason Lundegårdh (1924) uses the coefficient Q_1 . Only an investigator who experiments with an utterly refined method, has a right to measure by this standard. Otherwise he runs the risk of determining apparent variations occasioned by small experimental errors. For instance, the Q_1 1.08 changes with an error of 5 % already in 1.13 or 1.03, which seems to be a very great difference. This criticism applies to Lundegårdh himself; with detached leaves it is impossible to experiment accurately. Besides, L. measures already after a preliminary heating of 5 minutes. The question remains, whether the assimilation has already become constant.

The values Lundegårdh finds for Q_{10} with cucumber, potato and tomato leaves are from $5-15^\circ$: 5.0, 5.6 and 6.4; $20-30^\circ$: 1.6, 2.1 and 1.6; so Q_{10} falls considerably.

Blackman and Matthaei (1905), and Blackman and Smith found temperature coefficients between 2.0 and 2.5.

Miss van Amstel (1916) found a lower Q_{10} with *Elodea*, in which case presumably the diffusion of carbon dioxide acted disturbingly, as she supposed herself.

A lower Q_{10} of 1.5 was also found by Willstätter and Stoll (1918) with green leaves, while yellow leaves gave a still lower Q_{10} .

Osterhout and Haas (1919) found from $17-27^\circ$ a $Q_{01} = 1.81$ with *Ulva*.

Warburg (1919) found with *Chlorella* a high coefficient between 5° and 10° of 4.5—5. It fell regularly at higher temperatures.

The determinations of Yabusoe (1924) suffer from too great an interval of temperature. Emerson (1929) could prove that the direct proportionality between temperature and assimilation, as claimed by Yabusoe, is a result of this large interval and is therefore only imaginary. In reality the curve has a sigmoid shape.

The experiments of Emerson in this matter are the most important, as he determined complete temperature curves with an accurate method (of Warburg) with a lower plant (*Chlorella*), in which case the other factors may easily be brought in maximum. Emerson's result is in principle in accordance with that of Warburg, the increase of the assimilation diminishes with the rise of temperature, in other words μ , in which Emerson expresses his result, is not constant and falls gradually.

Finally van den Honert (1928, 30) investigated the influence of temperature with *Hormidium flaccidum*. He found from 12° to 20° a Q_{10} of 1.87.

Ecological investigations about the influence of temperature under natural circumstances cannot of course be put on the same level with investigations where all factors are in maximum and will not be discussed here.

It has been my intention to study this temperature influence on a larger range with *Hormidium*.

Further I felt interested in the influence of temperature on respiration. It will appear later on (Ch. VIII—XI) that there are reasons to assume a connection between assimilation of carbon dioxide and respiration.

On the one hand the possibility exists that in case temperature is „limiting factor”, in reality a part of the assimilatory chain process is investigated, viz. a dark chemical part of it (an opinion held by many investigators).

On the other hand the possibility must not be excluded, that, while the influence of temperature on assimilation

is studied, yet nothing else is done than to determine the influence on the total life-intensity of the organism as a whole, namely when the vitality of the organism would primarily determine the intensity of assimilation. The intensity of respiration would in that case be the direct criterion of this influence of temperature on the organism, the intensity of assimilation on the other hand only an indirect result. Temperature would in this case act as an indirect factor only, viz. by means of accelerating protoplasmic processes, other than photosynthesis *sensu stricto*. This acceleration of the protoplasmic processes might then be the cause of the acceleration of photosynthesis.

It is quite imaginable that protoplasm produces an essential substance for the photosynthetic process, for instance an enzyme, and that it is the rate of this process that controls the total rate of assimilation, but there are other possibilities as well (cf. Spoehr and Mac Gee).

The conception given above would at once be refuted if the influence of temperature on assimilation and respiration should prove to be quite different, but it would be supported, though not proved, when both processes were influenced consistently. For these reasons an inquiry about the influence of temperature on respiration became necessary, in addition to that on assimilation.

A simultaneous investigation of the influence of temperature on assimilation and respiration has been carried out by Lundegårdh (1924). In his results nothing can be discovered of a parallelism between the two processes. The determination of the temperature coefficient of respiration is, however, rather difficult in case of detached leaves. Respiration decreases in the dark, whereas, according to Meyer and Deleano (1911, 13), cutting causes a temporary rising, and a sudden increase of temperature would temporarily give too high a respiration as well.

One reason more made the determination of the coefficient

of temperature desirable. With each measurement of the assimilation a correction for the respiration must be made. Virtually a determination of respiration should be made after each measurement of assimilation. If the Q_{10} of the respiration is known one determination is sufficient. Moreover an unavoidable error is made, because it has been shown in Chapter III that respiration strongly increases in light. This correction would, however, be quite superfluous if the Q_{10} of both processes should prove to be equal. Anticipating my result, I may say already now that this is, approximately, the case. This conveys, therefore, a great simplification. It is sufficient to determine the influence of temperature on the apparent assimilation, which will be of the same order of magnitude as the influence on the real assimilation.

My method deviates from those of some other investigators. While Warburg, Lundegårdh, Emerson a. o. start their measurement, as soon as the temperature equilibrium has set in, I waited till the assimilation appeared to be constant, in other words, until the establishment of a *physiological* equilibrium.

According to Blackman the determination should have to be made in an infinitely short time. If the assimilation could be measured in this way, the temperature curve at injurious temperature, too, should be quite normal, according to Blackman's conception, and would not show any optimum-type. The injurious action of the high temperatures could be eliminated in the results in this way. The supposition underlying this would be right, if indeed we studied a purely chemical reaction. If, however, Harder (1930) and Kostytschew (1931) are right and „höchst komplizierte Reizketten" are connected with the assimilation, we may not at all expect that the final rate of assimilation would immediately set in. The alterations in the protoplasm, which would determine the

final rate of assimilation, would precede the establishment of the equilibrium. In order not to neglect this possibility, it is better always to check the constancy of the reaction.

B. The Influence of Temperature on Assimilation.

The range of temperature investigated lies between 4° and 34°. Above 30° it is already difficult to observe a constant assimilation. It may be that secondary factors, as desiccation of the algal film are also responsible here.

The 150 W. lamp at 13 cm distance was always used for illumination. My results are to be found in table 6.

In a few series of experiments the variations are rather great, much greater at any rate than might be expected with this accurate method. A possible explanation of these variations will follow later (Chapter XIII § E).

TABLE 6.

The influence of temperature on CO₂ assimilation (in November).
Van den Honert's strain.

Exp. 37, 7-11-30. Conversion factor $\frac{100}{420}$

Time	Temperature	Assimilation	Assimilation converted
15.55	20.00	420	100
16.45	24.00	542	129
17.25	28.00	736	175
19.50	31.00	928	221
20.40	34.00	1025	244

Exp. 38, 11-11-30. Conversion factor $\frac{100}{465}$

10.30	17.00	390	84
11.30	21.00	483	104
14.00	25.00	646	139
14.55	29.00	818	176
16.35	21.00	499	107

Exp. 39, 13-11-30. Conversion factor $\frac{100}{360}$

Time	Temperature	Assimilation	Assimilation converted
10.20	30.00	563	156
12.10	26.00	493	135
14.20	22.00	422	117
15.25	18.00	296	82
16.35	14.00	240	67
17.35	22.00	447	123
20.05	30.00	615	171

Exp. 40, 17-11-30. Conversion factor $\frac{100}{470}$

10.50	29.00	572	122
11.50	24.00	544	116
14.25	19.00	447	95
15.30	14.00	307	65
16.55	9.00	194	41
18.25	4.00	115	24
20.25	14.00	294	63

Exp. 41, 25-11-30. Conversion factor $\frac{100}{245}$

10.45	14.00	137	56
11.20	18.00	167	68
14.15	22.00	287	117
15.20	26.00	354	144
16.15	30.00	434	177
17.00	22.00	285	116
18.10	18.00	232	95
20.30	14.00	152	62

Exp. 42, 26-11-30. Conversion factor $\frac{100}{340}$

11.20	15.50	242	70
12.10	19.50	317	92
14.30	23.50	513	149
15.25	27.50	621	180
16.05	31.50	678	197
17.05	23.50	449	130
17.55	19.50	312	90

Exp. 43, 28-11-30. Conversion factor $\frac{100}{455}$

Time	Temperature	Assimilation	Assimilation converted
10.40	16.00	360	79
11.55	12.00	233	51
13.50	12.00	226	50
15.05	8.00	154	34
16.00	4.50	± 84	± 18
16.55	12.50	260	57
17.55	17.50	384	84
20.00	22.50	531	117

I have represented this experimental series graphically and drawn a probable curve between the determined points. In this way the assimilation could be read off at 20°. In order to compare all experimental series with one other, the assimilation at 20° was put as 100. The rate of assimilation at 20° is a physiological measure for the quantity of material used. This measure was also taken by van den Honert. All values must consequently be multiplied by a conversion-factor (4th column of the table). The values calculated have been recorded in fig. 11. It appears that in spite of the individual variations a smooth curve can freely be drawn between the points. The increase of the rate of assimilation with the temperature is, consequently, very regular. Pronounced optima, as found by Lundegårdh, are altogether wanting.

Q_{10} can be calculated from the curve. These values have been combined in table 7. In the last column the determined Q_{10} has been corrected in the way as discussed in the second chapter. At low temperatures Q_{10} is largest. Apart from this it is much more uniform than Emerson found with *Chlorella*. Above 10° it is fairly well constant. Since $\log Q_{10}$ is inversely proportional to the product of the absolute temperatures ($\log Q_{10} =$

$\frac{10 A}{T \times (T + 10)}$, if $A = R \mu$, $R = \text{gas constant}$, $\mu = \text{energy of activation}$), the characteristic μ is also nearly constant. μ calculated from this formula, usually put in the form:




Fig. 11. The influence of temperature on photosynthesis in Hormidium (rate of assimilation at 20° put at 100).

$$\frac{R_2}{R_1} = e^{\frac{\mu}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)} \text{ or } Q_{10} = e^{\frac{\mu}{R} \left(\frac{1}{T} - \frac{1}{T+10} \right)} \text{ is 11400 from } 15^\circ - 25^\circ.$$

TABLE 7.

Temperature-coefficient of photosynthesis.

Temperature	Q_{10} determined	Q_{10} corrected
5—15	2.84	2.77
10—20	2.30	2.24
15—25	2.02	1.96
20—30	1.93	1.87

Q_{10} being constant shows that the total chemical reaction complex is possibly determined by one part of the process.

These experiments were all made in the month of November 1930. A few experiments performed in May 1931 gave similar results, cf. table 8.

TABLE 8.

The influence of temperature on CO_2 assimilation in May.

Van den Honert's strain.

Exp. 44, 16—5—31.

Time	Temp.	Assim.	Q_{10} determ.
10.10	13.40	292	} 2.18 2.10
11.20	18.40	404	
13.15	23.40	636	
14.20	28.40	848	

Exp. 45, 19—5—31.

10.55	10.00	356	} 2.25 2.07 1.85
13.45	15.00	568	
14.50	20.00	811	
16.05	25.00	1177	
17.20	30.00	1504	

Exp. 46, 21-5-31.

Time	Temp.	Assim.	Q_{10} determ.
10.45	11.00	302	
12.05	16.00	474	2.51
14.15	21.00	757	2.30
15.30	26.00	1089	
16.55	16.00	484	2.25

It is of importance to know this, because it appears, that, in spite of the different properties which the way of cultivation can create (Chapters VI, VIII), the temperature coefficient of the reaction remains unchanged.

The experiments mentioned here have all been made with the *Hormidium* strain of van den Honert. A similar result was obtained with Pringsheim's strain.

It is remarkable that my results give a higher Q_{10} than van den Honert found with the same alga, though the difference is not very large. He found from 12—20° 1.87, whereas I find 2.15. Deviating values were sometimes stated, so that this want of uniformity is apparently inherent in the material used.

C. *The Influence of Temperature on Respiration.*

The measuring of Q_{10} of the respiration presents some difficulties. A falling respiration will be found when algae are taken from the window and placed in darkness. To prevent this the culture was placed in the dark on the preceding evening.

Good results were also obtained by the experiments 33—35 in Chapter VI, in which periods of darkness alternated with weak illumination.

In table 9 the results of the experiments are combined. Only with one experiment (50) the results deviate and a high Q_{10} was found at a rather high temperature. In the other experiments the determined value was reduced

TABLE 9.

The influence of temperature on respiration.

Van den Honert's strain.

Exp. 48, 20-11-30. Conversion factor $\frac{100}{138}$

Time	Temperature	Respiration	Respir. converted
11.25	8.00	57	41
13.30	13.00	96	70
15.50	18.00	117	85

Exp. 49, 18-2-31. Conversion factor $\frac{100}{52}$

10.50	9.00	24	46
12.10	14.00	37	71
13.30	19.00	40.5	78
14.50	24.00	67	129
16.05	29.00	92	177
17.10	34.00	125	240
18.30	19.00	40	77

Exp. 50, 5-3-31.

11.10	33.50	92	—
12.40	28.50	43	—
14.20	23.50	28.5	—
15.30	18.50	18	—
16.50	28.50	57	—

Exp. 51, 16-5-31. Conversion factor $\frac{100}{38}$

10.10	13.40	25	66
11.25	18.40	33	87
12.20	23.40	48	126
14.20	28.40	71	187

Exp. 52, 19-5-31. Conversion factor $\frac{100}{51}$

11.15	10.00	26.5	52
13.35	15.00	32.5	64
14.55	20.00	53	104
16.15	25.00	68	133
17.25	30.00	81	159

Exp. 53, 21-5-31. Conversion factor $\frac{100}{34}$

Time	Temperature	Respiration	Respir. converted
10.50	11.00	22.5	66
12.40	16.00	26.5	78
14.15	21.00	35.5	104
15.30	26.00	48	141
17.00	16.00	27.5	81

Exp. 33, 22-5-31 (table 5). Conversion factor $\frac{100}{38}$

11.35	10.00	17	45
13.55	16.00	30	79
16.15	22.00	43	113
18.25	28.00	59	155

Exp. 34, 27-5-31 (table 5). Conversion factor $\frac{100}{28}$

10.45	28.00	42	150
12.40	22.00	32	114
14.50	16.00	21	75

Exp. 35, 3-6-31 (table 5). Conversion factor $\frac{100}{38}$

10.25	14.00	24	63
12.30	20.00	40	105
14.40	26.00	56	147
16.30	20.00	40	105
17.30	14.00	19	50

again to a respiration of 100 at 20°. I thought it better to exclude from this the deviating result of exp. 50, which was possibly caused by too high heating. The values calculated are graphically represented in fig. 12.

The individual deviations are also very large here. The respiration is rather small, and the experimental error for that reason larger. Yet the curve can be pretty accurately determined. The great resemblance with the temperature

curve of the assimilation is very striking. Q_{10} has been determined from the curve, the values are to be found in table 10.




Fig. 12. The influence of temperature on respiration in Hor-midium (rate of respiration at 20° put at 100).

Q_{10} is but a little lower than the coefficient of the assimilation. This agreement is of course *no proof* that the assimilation proceeds under the control of other processes, connected with the respiration, but seeing that we shall

observe other points of agreement between the two processes later on, it is certainly important that the temperature coefficient does not speak against the idea of an indirect action of temperature on the assimilation.

It is interesting that with *Hormidium* the respiratory Q_{10} is even a little more constant than the Q_{10} of assimilation. This may point to the fact that the whole respiratory reaction complex at all temperatures is possibly controlled by one part of the process.

TABLE 10.

Temperature-coefficient of respiration, compared with the coefficient of assimilation.

Temperature	Q_{10} respiration determined	Q_{10} respiration corrected	Q_{10} assimilation corrected
10—20	2.06	2.09	2.24
15—25	1.87	1.91	1.96
20—30	1.80	1.85	1.87

The temperature characteristic μ is from 15° to 25° calculated at 11000.

CHAPTER VIII.

The Influence of Potassium Cyanide on Assimilation and Respiration.

A. Introduction.

Warburg investigated the influence of prussic acid on assimilation. He stated that prussic acid has, in a weak concentration, a retarding influence on assimilation, when temperature is the limiting external factor, that this retardation, however, does not appear when light is the limiting factor. From these facts Warburg concluded that photosynthesis is at least composed of two stages, one of which, the Blackman reaction (the chemical part of the assimilation), would be sensitive to HCN

only, and that this process would be brought about by a heavy metal catalysis. Another fact found by Warburg is that assimilation, in case of increasing concentration of the cyanide, cannot be retarded beyond a certain limit. This limit is the compensation point of carbon dioxide assimilation and respiration. Warburg is of opinion that a high prussic acid concentration completely retards the splitting off of the O_2 from the CO_2 , but not of the O_2 from the respiratory products. The photochemical mechanism remains, therefore, unimpaired, but only the dark reaction in which an active derivate of CO_2 is produced, is inhibited.

Warburg could not find a similar action on respiration. On the contrary, respiration appeared stimulated by concentrations where the assimilation became greatly retarded. Retardation does not set in until in a concentration of $\frac{1}{10}$ mol. (Warburg 1921).

Another investigator who made assimilation experiments with prussic acid was Emerson (1929). He found that *Chlorella*, grown in a nutrient solution deficient in Fe, are poor in chlorophyll, and more sensitive to HCN than normal cells. Therefore Emerson is of opinion that chlorophyll not only takes part in the photochemical reaction, but also in the chemical one. I would, however, observe that the chlorophyll content of cells grown in a solution deficient in Fe, is certainly the only *visible* point of difference, but that it is also to be expected that it is the Fe containing enzyme that will have been diminished by this treatment.

I studied the influence of potassium cyanide on respiration and assimilation with different light intensities again for *Hormidium* (van den Honert's strain).

B. *Experiments with Low Concentrations of KCN, Showing a Stimulating Influence.*

$\frac{1}{10000}$ mol. KCN is added to the algal film in experiment

54. This concentration will probably soon diminish by evaporation. Table 26 gives the result, which appears even more clearly from fig. 13. It shows that the assimilation is increased by the addition of KCN. It is important that this does not occur only, when temperature is the limiting factor, but that the whole light intensity-assimilation curve rises to a higher level. In strong light the increase is 27 %, in weak light 18 %.

TABLE 11 (fig. 13).

Stimulating influence of a treatment with $\frac{1}{10000}$ mol. KCN at different light intensities.

Exp. 54, 26-3-31. Van den Honert's strain. Temp. 18.00°.

Solution	Time	Light intensity			Assimilation	Increase in per cent
		transmitted	voltage	corrected		
normal	10.35	2300	—	(2300)	413	
"	11.15	611	220	611	295	
"	12.05	205	221½	210	130	
"	13.00	0	—	0	-29	
+ $\frac{1}{10^4}$ mol. KCN	14.25	1000	—	(1040)	488	+ 30
"	15.15	205	223	215	158	+ 18
"	16.05	455	223	477	316	+ 25
"	16.50	1000	223	1048	478	+ 27
"	17.25	2300	—	(2400)	526	+ 27
"	18.30	0	—	0	-38.5	+ 33

The respiration has also increased; the increase is 33 %.

In the following experiments, recorded in table 12, a similar stimulation, both of assimilation and respiration is found.

Experiments 55 and 60 were carried out in the same way as exp. 54. The evaporation of prussic acid was prevented

in the other experiments by adding KCN in an equal concentration to the buffer mixture. In some experiments the increase is rather considerable. Both in experiment




Fig. 13. Stimulating influence of the addition of a low concentration of KCN on photosynthesis in *Hormidium*. The assimilation is increased in low and in high intensity of light; increase in respiration 33%.

TABLE 12.

The influence of low KCN concentrations, not added to the buffersolution.

Exp. 55, 30–3–31. Temp. 19.00°. KCN $\frac{1}{2000}$ mol.

Solution	Time	Light intensity			Assimilation	Increase in per cent
		transmitted	voltage	corrected		
normal	11.15	2300	—	(2300)	472	
"	12.00	611	222½	606	278	
"	12.55	205	221¾	201	121	
"	13.45	0	—	0	-38	
+ $\frac{1}{2000}$ mol. KCN	15.20	1000	223	1000	397	+ 6
"	16.20	205	222	202	135	+ 11
"	17.10	455	223	455	254	+ 14
"	18.20	0	—	0	-39	+ 3
"	19.30	2300	—	2300	533	+ 13

Exp. 56, 25-4-31. Temp. 19.00°. KCN $\frac{1}{50000}$ mol.

Solution	Time	Light intensity			Assimilation	Increase in per cent
		transmitted	voltage	corrected		
normal	10.15	2300	—	2300	504	
"	11.00	611	222½	606	382	
"	11.45	181	223	181	154	
"	12.40	0	—	0	-42	
+ $\frac{1}{50000}$ mol. KCN	14.00	2300	—	2300	541	+ 7
"	14.45	611	224	621	408	+ 3
"	15.30	181	224½	185	165	+ 6
"	16.25	0	—	0	-45.5	+ 8

Exp. 57, 30-4-31. Temp. 19.00°. KCN $\frac{1}{20000}$ mol.

normal	10.25	2300	—	(2300)	595	
"	11.05	611	224	611	410	
"	11.55	181	224¾	182	139	
"	13.00	0	—	0	-50	
+ $\frac{1}{20000}$ mol. KCN	14.40	2300	—	(2300)	751	+ 26
"	15.20	611	224½	616	455	+ 11
"	16.10	181	224½	182	155	+ 12
"	17.10	0	—	0	-55	+ 10

Exp. 58, 11-5-31. Temp. 19.00°. KCN $\frac{1}{10000}$ mol.

normal	15.20	2300	—	—	766	
"	16.10	0	—	—	-63.5	
+ $\frac{1}{10000}$ mol. KCN	17.30	2300	—	—	659	- 14
"	18.30	0	—	—	-57	- 10
"	19.30	2300	—	—	777	+ 1
"	20.20	0	—	—	-61	- 4

Exp. 59, 13-5-31. Temp. 19.00°. KCN $\frac{1}{40000}$ mol.

Solution	Time	Light intensity			Assimilation	Increase in per cent
		transmitted	voltage	corrected		
normal	12.05	2300	—	—	1132	
"	13.00	0	—	—	-43	
+ $\frac{1}{40000}$ mol. KCN	15.05	2300	—	—	1167	+ 3
"	16.05	0	—	—	-57	+ 33
"	17.05	2300	—	—	1140	+ 1
"	18.10	0	—	—	-54	+ 26

Exp. 60, 15-5-31. Temp. 19.00°. KCN $\frac{1}{10000}$ mol., not

added to the buffer solution.						
normal	11.30	2300	—	—	876	
"	12.30	0	—	—	-46.5	
+ $\frac{1}{10000}$ mol. KCN	14.50	2300	—	—	919	+ 5
"	15.55	0	—	—	-49	+ 5
"	16.55	2300	—	—	969	+ 11
"	17.50	0	—	—	-45	- 3

55 and in 57 the assimilation rises with the same amount in strong and weak light. In exp. 57 an initial strong increase seems gradually to become a little smaller. The fact, found in experiment 54, viz. that the respiration behaves like the assimilation, is generally confirmed. That the respiration cannot be determined with great precision must not be overlooked, however.

It appears in experiment 58, contrary to what was found in exp. 54, that the concentration $\frac{1}{10000}$ mol. already retards the assimilation. After the lapse of a few hours, however, a recovery sets in and the retardation passes into an acceleration. Later on we shall repeatedly meet with this phenomenon. Also the respiration recovers a little.

C. Experiments with Higher Concentrations of KCN, Showing a Retarding Influence.

In these experiments KCN was added, besides to the alga, also to the buffer-mixture.

In the first place I call the attention to exp. 61, which shows the same type as 58, described above, and even more distinctly. The addition of $\frac{1}{1000}$ mol. KCN causes an initial fall of 13 %, which, after some hours has changed into an increase of 21 %.

TABLE 13.

Retarding influence of KCN, changing in course of time into an increasing one. Van den Honert's strain.

Exp. 61, 5-5-31. Temp. 19.00°. KCN $\frac{1}{5000}$ mol.

Solution	Time	Light intensity			Assimilation	Increase in per cent
		transmitted	voltage	corrected		
normal	10.20	2300	—	(2300)	360	
"	11.05	611	221	604	260	
"	11.55	205	221 $\frac{3}{4}$	205	121	
"	13.00	0	—	0	-28	
+ $\frac{1}{5000}$ mol. KCN	14.30	2300	—	(2300)	313	- 13
"	15.15	611	221 $\frac{3}{4}$	611	248	- 5
"	16.05	205	222	206	128	+ 5
"	17.15	2300	—	(2300)	405	+ 13
"	18.10	0	—	0	-30	+ 7
"	19.15	2300	—	(2300)	434	+ 21

Most remarkable is the result of the experiments 62 and 63, which were made two days later, in quite the same way as exp. 61 (cf. table 14).

The retardation is extremely strong in exp. 62, somewhat less in 63. In both the assimilation does not show any inclination to recover; it has on the contrary, diminished

again after three hours in exp. 63. Respiration is little affected, in exp. 62 a slight decrease, in exp. 63 even an increase.

TABLE 14.

Strong retardation of photosynthesis after the addition

of $\frac{1}{5000}$ mol. KCN.

Exp. 62, 7-5-31. Temp. 19.00°. KCN $\frac{1}{5000}$ mol. (addition at 13.03).

Solution	Time	Light intensity	Assimilation per hour	Increase or decrease in per cent
normal	11.35	2300	941	
"	12.30	0	— 51	
+ $\frac{1}{5000}$ mol. KCN	13.40	2300	± 512	
"	13.50	"	± 344	
"	14.00	"	± 230	
"	14.10	"	± 170	
"	14.20	"	± 152	— 84
"	14.55	0	— 44	— 14

Exp. 63, 7-5-31. Temp. 19.00°. KCN $\frac{1}{5000}$ mol. (addition at 17.25).

normal	16.10	2300	856	
"	17.00	0	—46.5	
+ $\frac{1}{5000}$ mol. KCN	18.07	2300	438	— 49
"	18.20	"	365	— 57
"	19.15	0	—49.5	+ 7
"	20.15	"	—53	+ 14
"	21.10	2300	311	— 64

The momentary disposition of the alga is apparently of great influence on the rate of retardation. In those days warmer weather had set in. This remarkable behaviour

TABLE 15.

Retardation of assimilation and respiration by the addition of higher KCN concentrations.

Exp. 64, 23-4-31, (fig. 14). Temp. 19.00°. KCN $\frac{1}{1000}$ mol.

Solution	Time	Light intensity			Assimilation	Increase or decrease in per cent
		transmitted	voltage	corrected		
normal	10.25	2300	—	(2300)	826	
"	11.35	611	221	621	506	
"	12.20	181	223	190	201	
"	13.20	0	—	0	-50	
+ $\frac{1}{1000}$ mol. KCN	14.50	1000	223½	1056	315	— 62
"	15.45	0	—	0	-18	— 64
"	16.35	181	223½	191	100	— 51
"	17.15	2300	—	(2400)	230	— 72
normal	19.55	2300	—	(2400)	476	— 42
"	20.40	0	—	0	-55.5	+ 11
"	34.00	2300	—	—	827	+ 0
"	35.05	0	—	—	-55	+ 10

Exp. 65, 10-4-31. Temp. 19.00°. KCN $\frac{1}{400}$ mol.

normal	10.55	2300	—	(2300)	581	
"	11.40	611	221	621	401	
"	12.35	205	—	(205)	160	
"	13.30	0	—	0	-43	
+ $\frac{1}{400}$ mol. KCN	15.00	1000	218	968	111	— 78
"	15.50	205	219	202	69	— 56
"	16.35	455	220	455	106	— 67
"	17.15	1000	223	1048	98	— 81
"	17.55	2300	—	(2400)	78	— 87
"	18.50	611	223	640	88.5	— 79
"	19.50	0	—	0	-36	— 16

is all the more striking, when we compare it with exp. 64 and 65 of an earlier date. The concentration of KCN was much stronger, $\frac{1}{1000}$ and $\frac{1}{400}$ mol., the retardation not stronger than in exp. 62.




Fig. 14. Retarding influence of the addition of KCN on photosynthesis in *Hormidium*. Contrary to Warburg's results with *Chlorella*, the assimilation in *Hormidium* is retarded in high as well as in low light intensity. Decrease of respiration 64 %. The „optimum type” of the KCN-curve is possibly incidental.

- Rate of assimilation 2 hours after the expulsion of KCN.
- Rate of assimilation 16 hours after the expulsion of KCN.

Results of experiment 64 are given in table 15. Contrary to Warburg's result with *Chlorella*, we find that the assimilation is retarded in weak as well as in strong light, although to a somewhat smaller measure in weak light (fig. 14). Exp. 64 is the only example where a corresponding strong retardation of the respiration is found. In very strong light the assimilation seems to decrease. It is not settled yet, if this really points to the existence of a light-

assimilation optimum; perhaps the retardation had become a little stronger after some time.

It has been investigated whether the retardation is reversible. The respiration recovered immediately indeed, the assimilation increased, though a considerable retardation remained. The next morning, however, it appeared that also the assimilation had regained its old value.

In exp. 65 (table 15) the retardation is stronger, and, besides, increases a little in course of time. In weak light the retardation is distinct, though again somewhat smaller than in strong light. There is again a flat assimilation-optimum. We may perhaps explain this by an increase of respiration in the stronger light.

D. Experiments, Showing a Complete Inhibition of Assimilation after the Addition of KCN.

Finally, in two experiments, a complete stopping of the carbon dioxide assimilation was observed (table 16).

The concentrations $\frac{1}{2000}$ and $\frac{1}{1000}$ mol., which were used, show how this phenomenon of retardation is independent of the concentration used, but seems to depend on the individual disposition of the alga. In the first experiment the respiration had not been changed; in the second it had been retarded, though not altogether wanting. In exp. 66 there is still at first some assimilation, which disappears gradually; in exp. 67 the assimilation is directly stopped altogether.

Both experiments are contradictory to the results of Warburg, as according to him the retardation would stop at the compensation point of assimilation and respiration. An explanation of this deviating result of Warburg may perhaps be attempted, when we observe that the assimilation under the influence of KCN was not directly brought to a standstill. When the assimilation is determined directly after the adding of KCN, as

TABLE 16.

Complete inhibition of photosynthesis after the addition of KCN.

Exp. 66, 13-5-31. Temp. 19.00°. KCN $\frac{1}{2000}$ mol.

Solution	Time	Light intensity	Assimilation per hour	Increase or decrease in per cent
normal	12.05	2300	769	
"	13.00	0	-32	
+ $\frac{1}{2000}$ mol. KCN	14.45	2300	120	- 84
"	15.05	"	75	- 90
"	16.10	0	-33	+ 3
"	17.10	2300	7	- 99
"	18.10	0	-31	- 3

Exp. 67, 15-5-31. Temp. 19.00°. KCN $\frac{1}{1000}$ mol.

normal	11.25	2300	844	
"	12.30	0	-34	
+ $\frac{1}{1000}$ mol. KCN	14.15	2300	81	- 90
"	14.40	"	35	- 96
"	15.00	"	- 4.5 ¹⁾	- 100
"	15.55	0	-21	- 42
"	16.55	2300	- 6 ¹⁾	- 100
"	17.50	0	-21	- 42

Warburg did, we may continually find a slight assimilation. As for the rest it is not excluded that *Chlorella* behaves otherwise than *Hormidium*.

¹⁾ Apparent assimilation (-25.5) + respiration in the dark (21) = - 4.5. The respiration in light is apparently stronger than in darkness.

E. Discussion of the Experimental Results.

Let us combine the results of the above experiments in a few words. It appeared that Warburg's results could not be confirmed. The addition of potassium cyanide caused a retardation not only when temperature, but also when light was the limiting factor. So on this ground it is not possible to distinguish two processes, one of which would be sensitive to K C N („Blackman reaction”).

It is, moreover, important that this retardation may disappear in course of time, and pass into an increase. This speaks strongly against Warburg's conception. It seems to me that the action of the K C N is much more complicated.

The fact that the same concentration of K C N on different days, can influence *Hormidium* (cultivated under rather constant conditions before a north-window) very differently, seems unaccountable in a simple way. It cannot be assumed that the quantity of heavy metal, probably Fe, which is, no doubt, present in unlimiting amount, would be so highly subjected to fluctuations in the cells.

This is also a reason to doubt Emerson's results. Even small differences in the way of cultivation cause a much changed sensitivity. This may also be the cause of the different behaviour of normal-*Chlorella* and *Chlorella* cultivated in a sugar solution deficient in Fe.

Another fact on which Warburg has built up his theory, the impossibility to retard the assimilation below the compensation-point of assimilation and respiration, could not be confirmed with *Hormidium*.

It stands to reason that it is not excluded that *Chlorella* and *Hormidium* behave quite differently in this respect. At any rate it appears, however, that no general validity may be placed on these facts and the theory built up on them.

Moreover it is an important fact that the assimilation can be stimulated by a small dose of KCN and that to the same extent, no matter whether either temperature or light is the external factor which limits the process. This fact, added to the observation that a uniform retardation also enters under both conditions, speaks strongly against the conception that in both cases different processes govern the assimilation. On the contrary, the uniformity of the controlling process is made plausible by these facts.

The stimulation of the assimilation in weak light is also important for another reason, because it appears from it that it is not the available light energy as such that limits the rate of assimilation. If this were the case, a stimulation would be impossible. That this stimulation is possible shows us that the assimilation, also when light is limiting, is not exclusively dependent on the quantity of light energy added, but at the same time on internal processes. (It may be possible that they themselves are connected again with the light intensity). Consequently it is not true either that the photochemical part of the process exclusively controls the entire reaction-chain under these circumstances, as is now generally accepted.

Other investigators who found a stimulation of the assimilation, are Bose (1924), who obtained stimulation with much more diluted solutions, and Schmucker (1928), who discovered an increase of the assimilation of *Cabomba aquatica* in diluted ether of 0.125 % and alcohol 0.3—1 percent. Sabalitschka and Weidling (1926) found a 100 % increase in acetaldehyd; Schmucker could not confirm this statement (cf. further Ch. X).

We have seen that assimilation can be uniformly retarded and uniformly stimulated under different circumstances. In case of retardation, we might think of an injury of the assimilatory apparatus, in other words of direct

action of the poison on the assimilatory process. The fact that also stimulation has been found, and even a retardation may pass into a stimulation, points to the action being indirect, and that the assimilation is affected „from the interior”. Therefore it is obvious that we must inquire into the other metabolic processes of the organism. The extent of the respiration can give us a hint on this metabolic state. We find indeed that a stimulation of the assimilation goes parallel with a stimulation of the respiration. When the assimilation was slightly retarded, the respiration too appeared to have decreased. These facts speak for the idea that the increase or decrease of assimilation is a result of the behaviour of other vital processes.

A strong retardation of the respiration accompanied a corresponding decrease of the assimilation only in one case (exp. 64). In the other experiments the contrary proved to be the case. This fact which is in accordance with the results of older investigations on the action of KCN and narcotics makes the above idea, which lays a connection between assimilation and other vital processes uncertain again. It does not imply a refutation, however. In order to escape this difficulty we need only take that in case of highly injurious stimuli a strong retardation arises through the direct action of the poison on the assimilatory apparatus. Another possibility is that a stronger poisoning calls into existence an increase in respiration, in the same way as e.g. wounding may act as a respiratory stimulus.

CHAPTER IX.

The Influence of Phenylurethane on Assimilation and Respiration.

It was Claude Bernard who discovered the retarding action of narcotics on the assimilation of carbon dioxide (1878). After him others, among whom Bonnier and

Mangin (1885), Ewart (1896, 97, 98), Jacobi (1899), Irving (1911), Thoday (1913), also dealt with this subject. As a result of these investigations it appeared a.o., that photosynthesis is retarded by concentrations that have no influence on respiration.

The question has been thoroughly reinvestigated by Warburg (1919, 20), who stated that the assimilation is retarded in about the same degree, whether light, CO_2 or temperature is the limiting factor of the process. From this Warburg concluded that photosynthesis is a surface-phenomenon, just like respiration. The action increases in a homologous series with the adsorbability. Warburg confirmed that higher concentrations are necessary to retard the respiratory process.

In Chapter V I discussed already some experiments in which the assimilation of CO_2 was investigated when phenylurethane or antipyrin had been added. Both substances appeared to retard the assimilation.

Though Warburg already showed that the assimilation altogether recovers after removal of the narcotic, I performed a few experiments on the matter.

The assimilation was measured before, during and after the narcosis. The narcotic was removed by transferring the alga on its old nutrient medium, part of which had been reserved for this purpose. This is not an absolute removal, but it is certainly a very strong dilution. The experiments showed that the recovery is perfect. There may even be a slight increase after the removal. This manifests itself distinctly in experiment 69. The increase amounts to 25 %, notwithstanding a slight loss of material during the transfer. This is much more than could be caused by „growth“. According to van den Honert the correction for growth should amount to 5.7 per cent in this experiment. We see the phenomenon is more complicated than a simple physical removal of the narcotic from the surfaces.

I thought it interesting to determine the action of phenylurethane on the whole light intensity-assimilation curve, just as was done when studying the influence of potassium cyanide. An example of such an experiment we find in table 17, graphically represented in fig. 15. In this figure the experimental points are numbered, the numbers indicating the time at which the determinations were made. It directly appears, in this and all following experiments, that the assimilation recovers from the first decrease, as

TABLE 17.

The influence of phenylurethane on the assimilation at different light intensities.

Exp. 71, 23-3-31, (fig. 15). Temp. 18.00°. Phenylurethane 2.4×10^{-4} mol.

No.	Solution	Time	Light intensity			Assimilation	Increase or decrease in per cent
			transmitted	voltage	corrected		
1	normal	11.30	2300	—	2300	593	
2	"	12.35	205	222½	207	158	
3	"	13.35	455	222½	459	331	
4	"	14.15	1000	—	1000	492	
—	"	15.10	0	—	0	-31	
5	+ phenylurethane	16.15	1000	222	1000	246	— 50
6	"	17.00	455	222	455	165	— 51
7	"	18.20	205	—	(205)	110	— 30
8	"	19.30	2300	—	(2300)	428	— 28
9	"	20.10	1000	224	1032	346	— 31
—	"	21.00	0	—	0	-27	— 13

will be clear from a comparison of the points 5 and 9. A curve has been drawn through the points 7, 8 and 9. The decrease appears to be uniform over the whole region, which is in accordance with Warburg's results with urethane, *but also with my own results with potassium cyanide*. Respiration has also diminished by 13 per cent.

Experiment 72 (table 18) gives the same results. Recovery is even more apparent. A comparison between the numbers 4, 7 and 9 shows this, but at the same time it is clear that the difference between 7 and 9 is not great, so that after some time a constant rate of assimilation seems to be reached. Retardation of respiration is as large as of assimilation. This points again to the possibility of a connection between the two processes, of which examples were also found in the preceding Chapters (VII and VIII).




Fig. 15. Retardation of photosynthesis in *Hormidium* after the addition of phenylurethane. The figures 1—9 show the successive experimental values. In agreement with Warburg's results, the assimilation is retarded in high and in low intensity of light. There is a recovery of the assimilation as is shown by the difference of the values 5 and 9. Decrease of respiration 13 %.

One may ask whether also respiration shows an initial strong retardation that decreases later on. An answer to this question is sought in experiments 73—76 recorded in table 19. In these experiments a recovery of respiration always occurred, though the results are not altogether convincing. The finest example is supplied by exp. 76. Retardation decreases uniformly, and the depression of the respiration is in an ideal way in accordance with that of the

assimilation. Whatever may be the cause of this recovery, it might be based on the urethane being oxidized, at any rate it is of importance, as the uniform retardation of assimilation and respiration is here illustrated in an unambiguous way.

This supposition of the oxidizing of the urethane is, however, unlikely on other grounds. In the first place it

TABLE 18.

The influence of phenylurethane on assimilation.

Exp. 72, 24-3-31. Temp. 18.00°. Phenylurethane 2.4×10^{-4} mol.

No.	Solution	Time	Light intensity			Assimilation	Increase or decrease in per cent
			transmitted	voltage	corrected		
1	normal	10.30	2300	—	—	718	
2	"	11.10	611	220	611	430	
3	"	11.55	205	221½	210	173	
—	"	12.55	0	—	—	-26.5	
4	+ phenylurethane	14.55	1000	221¼	1020	322	— 45
5	"	15.40	455	221	462	206	— 41
6	"	16.25	205	—	—	127	— 26
7	"	17.20	1000	—	—	483	— 18
—	"	18.20	0	—	—	-22	— 17
8	"	19.25	205	224	218	169	— 6
9	"	20.15	1000	—	—	522	— 12
10	"	21.05	2300	—	—	584	— 19

appears that the retardation has not quite disappeared even after several hours, but has only diminished. Also the experiments in Chapter V seem to contradict it. How could I have found the deviating result, represented in fig. 6, if the urethane had lost its efficacy? And finally there is the analogy with the action of KCN and Ba Cl (Chapter VIII and XI), where oxidation is out of the question.

TABLE 19.

The influence of phenylurethane on assimilation and respiration in course of time.

Exp. 73, 22-4-31. Temp. 19.00°. Phenylurethane 2.9×10^{-4} mol.

normal			with phenylurethane			
Time	Assimilation	Respiration	Time	Assimilation	Respiration	Decrease in per cent
10.25	519	54	13.10	412	48	— 11
11.30			14.15		52	— 21
			14.45		52	— 4
			16.35		51	— 4
			17.40		51	— 6
Exp. 74, 27-4-31. Temp. 19.00°. Phenylurethane 2.8×10^{-4} mol.						
11.00	580	29.5	14.15	257	20	— 56
12.10			15.15		22	— 33
			16.35		22	— 32
			17.30		22	— 25
Exp. 75, 4-5-31. Temp. 19.00°. Phenylurethane 3.0×10^{-4} mol.						
11.30	619	26.5	13.50	318	21	— 49
12.30			14.40		22	— 17
			15.55		21	— 21
			16.30		21	— 21
			17.25		24.5	— 15
			18.25		24.5	— 8
Exp. 76, 18-5-31. Temp. 20.00°. Phenylurethane 2.7×10^{-4} mol.						
11.30	709	45	14.25	617	35	— 22
12.35			15.30		41	— 13
			16.30		41	— 9
			17.30		43	— 7
			18.30		43	— 4

I consider this analogy of great importance. The assimilation reacts in all cases in a similar way, it does not matter which poison is the cause. The conclusion is evident that the initial large retardation is caused by a temporary lowering of the vital intensity of the protoplasm, which slowly recovers from the „shock”. The decrease, and recovery

afterwards, of the respiration also points in this direction. So it is not a mere matter of course that the urethane exclusively acts by diminishing the available surface of the assimilatory apparatus. On the contrary, it appears that something may be said in favour of an indirect influence of the injurious substance. Also the above-mentioned fact that the assimilation has been accelerated after the removal of the narcotic, and not simply recovers its former rapidity, is an argument in favour of this conception.

I made another attempt to prove, in a more direct way, that the urethane is not oxidized by the cells (exp. 77). I applied to a recovering algal film the same dose of urethane. In case the assimilation did not fall again, I should have proved that the urethane was not oxidized. However, the reverse appeared to be true. The retardation of 26 % had fallen to 7 %, a new dose enhanced it to 30 %, after some time it was 15 %. Though the result is in accordance with the supposition of oxidation of urethane, it has not conclusively proved it. It is not at all unlikely, that, owing to the absorption of the alga, the concentration of urethane in the liquid of the film had quickly diminished, and that, therefore, the addition of fresh urethane had caused the new decrease of assimilation.

Experiments 78 and 79 give us an insight into the influence of other concentrations.

The very slight concentration in exp. 78 (2.4×10^{-5} mol.) causes a small decrease (ass. 5—14 %, resp. 10 %), after which no noticeable recovery sets in. Stimulation does not occur here.

Exp. 79 is made with a stronger solution (1.2×10^{-3} mol.) The retardation is considerable, but diminishes gradually (from 77 % to 50 %). Respiration has been retarded less strongly, but recovers in the same way (from 15 % to 5 %). This, too, is in accordance with the results with potassium cyanide; assimilation is more sensitive to high concentrations of the injurious substance than respiration.

CHAPTER X.

The Influence of Glucose and Fructose on Assimilation and Respiration.

The preceding three chapters more or less all point to a connection between photosynthesis and respiration. It was a matter of course that an investigation of the assimilation under circumstances under which respiration increases was now to be taken in hand.

A great many investigators, of whom Borodin (1876) was the first (further: Aereboe (1893) Meyer and Deleano (1911, 13), Kniep (1914), Pantanelli (1914), Harder (1915), Plaetzer (1917) have shown the increase of respiration after illumination. This increase was ascribed to the production of oxidizable materials. Warburg (1922), Emerson (1927) and Genevois (1927) directly showed an increase of respiration by adding sugars to the algal suspension. In a preliminary experiment I could confirm this with *Hormidium* (a pure culture; increase 33 % in 0.7 % fructose). If I should also succeed in stimulating the assimilation under these conditions, then the relation between assimilation and respiration, or, at least, between assimilation and other protoplasmic processes connected with respiration, would be very convincingly obvious, because from a chemical point of view a retardation of assimilation should be expected rather than an acceleration, as sugars are indeed the final products of assimilation.

Spoehr and Mc Gee (1923) have observed that the sugars disappear out of the leaves in darkness, and that after a long stay in the dark the rate of assimilation has diminished. After continued exposure to light, both the rates of assimilation (cf. also Osterhout and Haas, 1918; Harder, 1930; Arnold 1931) and of respiration increase with time as the sugar content increases. They

consider this increase of assimilation as a result of the increase of sugar-content and respiration.

So the connection between assimilation and respiration, according to Spoehr and Mac Gee, consists in a dependence of the former on the latter. It would appear, however, even more convincingly, when the supply of sugar took place from the exterior, because in that case it would be proved that the increase of sugar content was really the cause of this phenomenon.

Treboux (1903) examined the action of sugars, and found that they lower assimilation in *Elodea*. The concentrations studied were high, and the lowering was probably caused by osmotic action, as Treboux thought.

I added the sugar by dropping a number of drops of a sugar solution on the algal film. Beforehand I removed the nutrient solution for the greater part by means of filter-paper, so that the final solution, was only a little more diluted than the solution added.

The sugars investigated were glucose and fructose. The action was investigated both in strong and in weak light. The *Hormidium* strains of Pringsheim and van den Honert were both used. The results of the experiments are to be found in table 20.

From these experiments an increase of assimilation appears, when a sugar solution of a concentration of 0.7—1.0 % was added, both with the pure culture of Pringsheim and van den Honert's strain. In a 2 % solution Pringsheim's strain gave a slight lowering of CO₂ assimilation. In strong light the increase is, to be sure, not exceedingly large, the values found are 5, 12, 9, 12, 5 and 6 %; but an increase of 655—730 in exp. 82 and another of 1097—1199 and 1233 in exp. 83 far exceed any experimental error. I consider the increase in weak light of great importance. This result is a confirmation of the results of experiments communicated in Chapter VIII,

in which it has been proved that in low light intensity assimilation is stimulated by the addition of a small dose of KCN.

TABLE 20.

The influence of glucose and fructose on assimilation and respiration.

Exp. 80, 8-5-31. Pringsheim's strain. Temp. 19.00°. Glucose 1 %.

Solution	Time	Light intensity			Assimilation	Increase or decrease in per cent
		transmitted	voltage	corrected		
normal	10.45	2300	—	—	422	
"	11.30	0	—	—	-29	
+ glucose	12.30	2300	—	—	444	+ 5
"	13.25	0	—	—	-44	+ 52

Exp. 81, 8-5-31. Pringsheim's strain. Temp. 19.00°. Glucose 2 %.

normal	15.00	2300	—	—	396	
"	15.45	0	—	—	-24	
+ glucose	16.50	2300	—	—	372	— 6
"	17.30	0	—	—	50	+ 108

Exp. 82, 9-5-31. Van den Honert's strain. Temp. 19.00°. Glucose 0.8 %.

normal	10.20	2300	—	—	655	
"	11.05	0	—	—	-56	
+ glucose	12.10	2300	—	—	730	+ 12
"	12.50	0	—	—	-52	— 7

Exp. 83, 12-5-31. Van den Honert's strain. Temp. 20.00°. Glucose 1 %.

normal	11.40	2300	—	—	1097	
"	12.35	0	—	—	-74	
+ glucose	13.50	2300	—	—	1199	+ 9
"	14.45	0	—	—	-63	— 17
"	15.50	2300	—	—	1233	+ 12
normal	17.15	2300	—	—	1277	+ 16
"	18.20	0	—	—	-78	+ 5

Exp. 84, 23-5-21. Pringsheim's strain. Temp. 20.00°. Glucose 0.7 %.

Solution	Time	Light intensity			Assimilation	Increase or decrease in per cent
		transmitted	voltage	corrected		
normal	10.20	2300	—	—	570	
"	11.10	181	221 $\frac{1}{4}$	181	190	
"	11.55	0	—	—	-49	
+ glucose	13.05	181	223	186	226	+ 15
"	13.55	2300	—	—	599	+ 5
"	14.30	0	—	—	-50	+ 2

Exp. 85, 25-5-31. Van den Honert's strain. Temp. 20.00°. Fructose 0.7 %.

normal	15.15	2300	—	—	860	
"	16.05	140	218 $\frac{1}{2}$	137	181	
"	17.00	0	—	—	-42	
+ fructose	18.45	140	223 $\frac{1}{2}$	148	233	+ 18
"	19.55	0	—	—	-62	+ 47
"	21.00	2300	—	—	914	+ 6

Exp. 86, 29-5-31. Van den Honert's strain. Temp. 20.00°. Fructose 0.7 %.

normal	14.50	140	220 $\frac{3}{4}$	140	198	
"	15.35	0	—	—	-32	
+ fructose	16.40	140	220 $\frac{3}{4}$	140	221	+ 12
"	17.10	0	—	—	-37	+ 15

It is peculiar that the respiration shows a decrease in two experiments. It is more or less clear that respiration decreases when the alga is taken away from daylight (Chapter III), but it is strange that an addition of the sugar does not counterbalance this effect.

Exp. 83, where I met with this phenomenon, has been continued. The sugar was removed and assimilation and respiration were again determined. The removal appeared to result in a new small increase. The assimilation rose from 1233 to 1277, and was now 16 % higher than in the beginning. Further the increase of respiration was noteworthy, which recovered from 63 to 78, and was now

even 5 % higher than at the outset. This may point to the action of the sugar being complicated, too. It may be that the stimulating action is checked by an injurious, possibly an osmotic, action, and that this accounts for the divergence of the results.

One more experiment was made with a culture (van den Honert's strain) which had stood in the dark continually for 62 hours (exp. 87). The concentration added was 0.7 %. Contrary to the expectation, that now the influence would be great, it appeared not to be so. Respiration really increased by 90 %, but assimilation proved to diminish by 24 per cent.

These few deviating results may, however, not turn our attention from the important fact, that it has become evident in this chapter, that an addition of sugars, which in most cases brings about an increase of respiration, causes, at the same time, an increase of assimilation. This fact points to the assimilation being affected by the protoplasm. According to its being provided with more sugar, the assimilation increases, altogether in accordance with the conception of Spoehr and Mac Gee.

The increase also takes place in weak light and this speaks in favour of the unity of the assimilatory process which in its entirety is controlled by internal processes, and not exclusively by external factors.

Now that it appears that the external factor light (when it is limiting factor) alone does not determine the intensity of the assimilation, but that the effect is also dependent on internal factors, we have to consider the possibility, that the increase of assimilation in stronger light is not exclusively brought about by more energy available for the photosynthetic process, but that the protoplasm, too, is affected by the light intensity, and that the effect of stronger light on the protoplasm promotes the assimilation again. The protoplasm therefore may ultimately determine the rate of the assimilation; the action of the external factor

is possibly by means of it; its action may be indirect.

This opinion is supported by the fact that light exerts a strong influence on the protoplasm (Chapter III). It appears that the organic system itself does not remain unaltered, that among the factors in the photosynthetic process, light is probably not the only variable.

In connection with the described increase of the assimilation by the addition of respiratory material, it should be noted here that Sabalitschka and Weidling (1926) found a stimulation of the assimilation of as much as 100 % by the addition of acetaldehyde, which can also serve as respiratory material. The concentration was small (0.016—0.032 %) and the increase is, according to them, effected by a stimulating action of the substance. This result could not be confirmed by Schmucker (1928).

Finally it may be mentioned that attempts have been made to stimulate respiration and assimilation by putting the algal film in water distilled from glass into glass. After remaining there for some time, this water was again replaced by fresh. For Shibata (1929) has found with *Chlorella*, that washing with distilled water enhances respiration. In my experiment neither respiration nor assimilation was affected by this treatment.

CHAPTER XI.

The Influence of $BaCl_2$ on Assimilation and Respiration.

While in the preceding chapter the question was treated, how carbon dioxide assimilation behaves under circumstances in which respiration is promoted, we shall now put the question how assimilation behaves if respiration has undergone a strong decrease. The reply to the first question appeared to be that under these conditions assimilation

was promoted as well. Will this parallelism also repeat itself in case of retardation?

In the preceding chapters we have found instances of a strong retardation of photosynthesis, but a great decrease of respiration is still wanting (except exp. 64 with KCN, where both processes appeared to be greatly retarded). We find the best example in the chapter on the influence of urethane, where both respiration and assimilation were slightly retarded.

The question was now to find a substance that has a retarding influence on the respiration. I found this substance in the paper of Shibata on the antagonistic action of electrolytes on respiration (1929). According to this research the respiration of *Chlorella* in a solution of one mol. Ba Cl_2 is diminished by 80 per cent. This substance seemed suitable to me for that reason, and also, because the depressed respiration, according to the statement of Shibata, remains rather constant, contrary to the result with (likewise strongly retarding) Ca-salts. The concentration I used in my experiments amounts to 0.5 mol.

In Chapter III the decrease of respiration in the dark has been stated. In order to prevent that a decrease of respiration could partly be caused by the influence of darkness (this influence being high in the month of June, as a result of the high light intensity), the algae were placed in the dark for some hours before the experiment.

The nutrient solution was removed by once washing the algal film with distilled water before the addition of Ba Cl_2 solution. In table 21 the results of the experiments have been given.

Exp. 89 and 90 have been made in strong light, exp. 91 in weak light. In the former experiments, both respiration and assimilation appear to be strongly diminished, the former (40 and 56 %) even a little more than the latter (25 and 44 %), but the agreement is nevertheless very sufficient.

TABLE 21.

The influence of 0.5 mol. Ba Cl₂ on assimilation and respiration.

Van den Honert's strain.

Exp. 89, 13-6-31. Temp. 21.00°.

Solution	Time	Light intensity			Assimilation	Increase or decrease in per cent
		transmitted	voltage	corrected		
normal	10.10	2300	—	—	555	
"	10.55	0	—	—	-48	
+ 0.5 mol. Ba Cl ₂	12.15	2300	—	—	416	- 25
"	12.55	0	—	—	-29	-40
"	14.00	2300	—	—	478	- 14
"	14.40	0	—	—	-32	-33

Exp. 90, 23-6-31. Temp. 21.00°.

normal	17.35	2300	—	—	1015	
"	18.25	0	—	—	-66	
+ 0.5 mol. Ba Cl ₂	20.00	2300	—	—	± 569	- 44
"	20.45	0	—	—	-29	- 56
aqua dest.	22.15	2300	—	—	640	- 37
"	23.00	0	—	—	-70	+ 6

Exp. 91, 25-6-31. Temp. 21.00°.

normal	17.15	139	221¼	139	158	
"	18.25	0	—	0	-66	
0.5 mol. Ba Cl ₂	20.05	139	223¼	143	127	- 22
"	21.10	0	—	0	-29.5	- 48

Exp. 89 gives retardation as a function of time. Altogether in accordance with the results of the experiments with potassium cyanide and urethane, we see the recovery of both assimilation and respiration. So it seems that here we have a very general reaction before us, of respiration as well as of assimilation. In the uniform retardation of the two processes and the simultaneous recovery, I see an indication of a connection between both vital processes.

It is important that Shibata, too, regularly finds a recovery after an initially strong decrease, and that after the addition of very different electrolytes. From Shibata's figures this is to be seen at a glance. Ca only excites a continual decrease in high concentrations. The activity of this substance is analogous to the action of higher KCN concentrations on the assimilation.

Ba Cl₂ has been replaced again by distilled water in exp. 90. Respiration recovers altogether, assimilation only partly. We have already an example of this phenomenon in exp. 64, where the removal of KCN had a similar effect. We must admit that this certainly does not point to a direct connection between the processes of respiration and assimilation.

Exp. 91 has been carried out in weak light. The retardation of the assimilation seems to be a little smaller (analogy with action of KCN); against a retardation of respiration of 48 %, there is a decrease of 22 % only.

In the last experiment (92) the influence of 1 mol. Ba Cl₂ was investigated. Here it appeared that the assimilation was totally, or almost totally inhibited (decrease 574 to 2). On the other hand the respiration still amounted to 23, against 48 before. To strong stimuli, therefore, assimilation is more sensitive than respiration. This also appeared to us when studying the influence of higher concentrations of KCN and of phenylurethane.

CHAPTER XII.

The Time Course of Photosynthesis.

A. *The Induction-Time of Photosynthesis.*

1. *Introduction.* In the preceding chapters we have more than once been led to the conclusion that the influence of the external factors is not direct, but that the rate of the assimilation is determined by the inter-

mediary of the protoplasm. Even when light is limiting and the increase directly proportional to its intensity, it appeared that the assimilation could be stimulated by factors that may cause a similar increase in respiration. When this conception of the indirect action of external factors, is right, then it is to be expected that the assimilation will not reach its full rate directly after the beginning of the exposure to light, but that some time will pass before the protoplasm has adapted itself to the new conditions and the assimilation has come to a constant intensity.

Warburg (1920) already established that in *Chlorella* the assimilation is indeed smaller in the first minutes, and within a short time rises from nothing to the final value.

Quite in accordance with his physico-chemical conception of the assimilatory process Warburg is of opinion that this increase in assimilatory activity is due to the existence of a photochemical induction-time in photosynthesis.

He came to this discovery, because he began intentionally to investigate, whether photosynthesis is characterized by a photochemical induction period, as has been proved to be the case in other photochemical processes. Therefore, we can readily understand that, when he stated that photosynthesis, too, showed an induction period, he did not think of any other way of explanation of the induction phenomenon, though he remarked that the induction of photosynthesis is not quite identical with the induction of the photochemical reaction of hydrogen and chlorine.

Possibly the influence of light on assimilation is more or less indirect; a change of the protoplasmic structure might be caused by illumination, and of this protoplasmic change the start of the assimilatory process might partly, or totally, be the result.

Certainty is altogether wanting, and calling the phenomenon "photochemical induction" is consequently hypo-

thetical. An explanation as proposed here, by which the induction is explained by means of preceding protoplasmic processes that are induced by light, seems equally probable.

It seemed interesting to investigate whether this induction time could also be observed with *Hormidium*, in order to settle which of the two hypotheses was true. This could be done by investigating the influence of temperature on the phenomenon. If the process were of a physiological nature then temperature had to influence it, just as temperature affects other physiological processes, where light plays a part (e.g. phototropical phenomena). On the other hand no distinct influence of temperature is to be expected from a phenomenon that is merely photochemical.

2. *Experiments on the Induction of Photosynthesis at Different Temperatures.* The presence of the phenomenon was demonstrated at 20°. When this appeared to be entirely in accordance with Warburg's result, the experiment was repeated at 14° and 26°, in order to answer the second question. The experiment was made as follows. The algal film was repeatedly illuminated for 0.5, 1, 1.5, 2 or 3 minutes. Between the exposures the culture was in the dark for a much longer period. After due correction for respiration the assimilation during a period of exposure could be determined. The respiration was not constant but continually falling, and had to be determined several times during the experiment. The short exposures had no observable influence on respiration (determined in the dark), so that an error was possibly not made in the respiration-correction (see Ch. III; yet an increase of respiration during the short exposure, without an after-effect on the respiration in darkness, is equally possible).

The illumination was effected with the projector lantern at full capacity. A new lamp with a little more candle power had been placed in it. In spite of this fact the

light was no longer quite in maximum at 26° and the assimilation about 25 % lower than it could have been. Yet I gave the preference to this illumination, as in this case the light passed a layer of water of 16 cm thickness, by which desiccation of the algal film which can easily occur in a prolonged experiment at this temperature, could be prevented.

TABLE 22, (fig. 16).

The assimilation in the first minutes after the beginning of the illumination. Van den Honert's strain.

Exp. 93, 5-6-31. Temp. 20.00°. Assim. at 10.35 : 795; per minute 13.25, reduced 10. Respir. at 11.20 : 73, at 12.50 : 54, at 15.47 : 50, at 18.35 : 41.5.

Time	Duration in minutes	Exposure in minutes	Assimil- ation	Assim. in x minutes	Assim. reduced
13.55	66	6 × 1	24	4.0	3.0
14.55	60	5 × 2	81	16.2	12.2
16.35	56	4 × 3	115.5	28.9	21.8
17.35	56	8 × 0.5	11	1.4	1.1
19.55	66	6 × 1.5	59.5	9.9	7.5

Exp. 94, 8-6-31. Temp. 14.00°. Assim. at 11.25 : 549; per minute 9.15, reduced 6. Respir. at 12.55 : 46, at 15.50 : 38, at 18.35 : 32.

14.00	60	8 × 1	16	2.0	1.4
15.00	54	5 × 2	46	9.2	6.0
16.45	54	10 × 0.5	4.5	0.5	0.3
17.35	48	4 × 3	67	16.8	11.0
19.40	48	5 × 1.5	24	4.8	3.1

Exp. 95, 9-6-31. Temp. 26.00°. Assim. at 10.50 : 972; per minute 16.2, reduced 13. Respir. at 11.40 : 72, at 12.31 : 50, at 14.50 : 47, at 18.20 : 42.5

13.25	43	4 × 2	85.5	21.4	17.2
14.05	46	6 × 1	57.5	9.6	7.8
15.40	48	8 × 0.5	29.5	3.7	3.0
16.30	50	4 × 3	174	43.5	34.9
17.20	48	5 × 1.5	93	18.6	14.9

Before the periodical exposure the assimilation in constant light was determined. This is the optimum velocity which the assimilation can reach at alternative exposure.

In order to facilitate a comparison the optimum rate at 20° was put at 10 per minute. The optimum rate at 14° and 26° reduced to this value, can be calculated from fig. 11 in chapter VII. We find 5.75 and 14.8. Now that the light is not quite in maximum, the increase will be a little slighter, especially at high temperature. It seems to me better to round the numbers to 6, 10 and 13. The results of the three experiments are to be found in table 22.

The great accordance with the results of Warburg is evident from these data. This is distinctly shown by fig. 16. It appears that the final optimum rate is fairly well reached after two minutes. What strikes the eye most is the great difference between the three curves. At 26° already 1 mm^3 has been assimilated after 12 seconds, at 14° this takes 55 seconds. This is certainly not a right measure for the induction process. If the possibility for assimilation is given as a result of the running down of the induction period, a larger quantity will be assimilated at higher temperatures, because the assimilation does proceed more rapidly at higher temperature. But this needs not point to a faster rate of the processes contributing the induction-chain.

In order to settle this point we have to see whether the optimum speed of the process is sooner reached at higher temperature than at lower. In practice it is difficult to state where the transition-point between curved and straight line is situated. It is much easier to fix another point; viz. the point at which the speed is one half of the final speed. A cross indicates this point in the figure.

This point is reached at 14° after 64, at 20° after 54, and at 26° after 24 seconds. A considerable difference is apparent. In spite of the fact that the optimum rate at 26° is much greater than at 14° , this higher rate of assi-

milation is reached much quicker than the lower at 14°. Therefore, the induction is greatly influenced by temperature, so that a chemical protoplasmic process plays a




Fig. 16. Assimilation in the first three minutes after the beginning of exposure to light at three different temperatures. The dots represent the rate of assimilation in 0.5, 1, 1.5, 2, and 3 minutes. The rate of assimilation rises to its full value (6, 10 and 13 per minute, represented by the lines in the right corner) in a few minutes. The crosses indicate the moment at which the speed of the process is half of the final speed. It takes a longer time before this point is reached at the lower temperature, which proves the sensitivity to temperature of the induction process.

induction is a process of a protoplasmic nature.

It has been established, therefore, that, before assimilation reaches its full intensity, a reaction takes place which is sensitive to temperature and caused by light. This points to a new indirect action of both factors in the photosynthetic process. A chemical reaction must occur in the protoplasm under the influence of both factors, before the proper photosynthetic activity may begin.

part here; Warburg's idea of photochemical induction must be rejected.

We can also get an impression to what extent temperature influences the induction, when we state the temperature coefficient from the values found at 14° and 26°. Q_{10} appears to be 2.2 and is strikingly in accordance with the temperature-coefficients of respiration and photosynthesis. This affords a new argument for the conception that

B. The Course of Photosynthesis after a Long Period of Darkness.

The increase of assimilation, described above, must be distinguished from that which Osterhout and Haas (1918), Spoehr and Mac Gee (1923), Harder (1930) and Arnold (1931) observed during the course of the experiment.

Harder could demonstrate that the assimilation of *Cladophora* and *Fontinalis* sometimes kept increasing the whole day. This rise was not interrupted by periods of darkness, on the contrary this caused an increase of even longer duration. Finally an optimum was reached, and a decrease set in. Thus Harder showed that the assimilation under constant conditions is subject to alteration.

Osterhout and Haas found a similar phenomenon with *Ulva* thallus. They stated, however, that the assimilation was already constant after 1½ hours.

Finally, Spoehr and Mac Gee observed this increase with leaves which had been in the dark for a long space of time. As said before, this increase paralleled an increase in sugar content and respiration, and they pointed to the latter factors as the cause of the increase in assimilation.

It appeared already to van den Honert (1928, 30) that an increase during the whole day, as was found by Harder, does not occur in *Hormidium*, and that the assimilation remains fairly constant. Only a slight increase of about 1 % an hour at 20° was stated. This result was regularly confirmed in my experiments. Van den Honert thought that he could explain this by an increase of the living material.

However, when I look through my protocols, it is evident that there exists an initial increase, after a long stay in the dark and also in the dark wintermonths. A few examples will illustrate this:

Exp. 37, 7th Nov. 1930. First determination after 15 min.

exposure. Ass. at 20°, strong light: 22 in 5 min., 54 in 10, 29 in 5, 31, 32, 35, 34, 35, 37, 35, 34. The last 6 values are fairly constant; so the assimilation increased during 45 minutes.

Exp. 19th Nov. '30. Ass. per 5 min.: 28, 39, 44, 45, 43, 41, 42, 42. First determination after 10 minutes' exposure. After 20 min. the assimilation is constant.

Exp. 22nd Dec. '30. Exposure from 1.26 to 2.35; 1.40—1.50 : 23; then 28 in 10 min., 17 in 5, 82.5 in 30. Constant after 34 min. exposure.

Exp. 24th April '31. After 16 hours' darkness. First determination after 20 min. exposure. Ass. 56 in 5 min., 62 in 5, 62 in 5, 132 in 10, 127 in 10, 127 in 10; constant after ± 25 minutes.

The above examples, to which I could add a few more, give an impression of the assimilation during the first hour. There is, apparently also in *Hormidium*, an increase of assimilation, as Osterhout and Haas found in *Ulva*. The only difference is that the constant rate is reached earlier.

In the preceding chapters we have seen that, all other conditions being constant the assimilation can be stimulated by the addition of certain substances such as KCN, narcotics and sugars; so it is not at all impossible that substances formed during the assimilation, stimulate the assimilation, as Spoehr and Mac Gee think. I do not mean by this to give an explanation of the increase as found by Harder. It would seem to me very unlikely that this increase of long duration would entirely be a result of improved nutritional conditions. I only want to point here to one of the possible causes of this phenomenon.

It also appeared that the respiration increased during the exposure by purely plasmogeneous action of the light (Ch. III). If it is true that the assimilation can be influenced by other vital processes, the increase of assimilation may also be explained by this plasmogeneous influence of light.

CHAPTER XIII.

Discussion.

It was mentioned before that some of Warburg's results are contradictory to mine. Of one of them we could say how Warburg might have arrived at his results, but not with any amount of certainty. It seems much more probable that the differences are due to the experimental material, and that *Hormidium* and *Chlorella* behave differently. ¹⁾ Only a thorough, comparative study of the assimilation of these and other organisms, which I consider of the greatest importance, can throw light upon this subject.

Should my opinion be right, it seems better to base my considerations on my own results. It is of more importance that we understand the assimilation in one organism than that we attach too much importance to differences, possibly based on individual peculiarities. Especially if actually assimilation is controlled by other vital functions, there is no reason to be astonished at a different behaviour. There would be more reason for such astonishment if we had always studied the "real" photosynthesis, and the plasmatic factor had showed itself to be negligible.

We need not, therefore, to worry about the apparant discrepancy of our results and those of Warburg. It seems rather a support for the idea that photosynthesis is controlled by the organism which has an individuality of its own.

A. *The Assimilation Reacts to the Addition of Uncommon Substances in a Similar Way under Different Conditions.*

The following grounds may be given for this statement:

I. The assimilation, when injurious materials are

¹⁾ The methods, too, are in principle the same.

added, is as much retarded in strong as in weak light (temperature or light intensity as limiting factors). We meet this in case of retardation owing to three altogether different causes, e.g. addition of *a.* phenylurethane, *b.* KCN, *c.* Ba Cl₂. The recovery from this injury acts similarly, both in high and low light intensity.

II. The assimilation may be stimulated, and again the increase is of the same extent under both conditions, whether this stimulation is caused by *a.* diluted KCN, or *b.* glucose and fructose.

Therefore we may conclude that the assimilatory process behaves as a unity; it does not matter whether either light or temperature is the limiting factor.

In relation to this conclusion the work of Briggs may be mentioned here. Briggs (1922) found that the photosynthetic activity of plants grown in culture solutions devoid of one of the elements necessary for normal growth was less than that of normal plants. This depression resulted whether light or temperature was the limiting factor. However, Briggs' results were contradicted by Gregory and Richards (1929). These investigators stated a slight depression of photosynthesis in light of low intensity, whereas the depression proved to be very marked in high light intensity. These very interesting results, however, may not be compared with the results of my experiments mentioned above, in which the immediate reaction of photosynthesis to the addition of unusual substances has been studied.

B. The Dependence of Photosynthesis of Internal Factors.

A first argument for a dependence of the assimilatory process on internal factors is the inconstancy of the assimilation while the external factors are kept constant.

Osterhout and Haas, Spoehr and Mac Gee, but especially Harder and Arnold have described this

phenomenon with other organisms. In Harder's experiments the assimilation appeared to increase during the whole day. The decrease, which followed subsequently, and also the decrease that Ewart, Pantanelli, Montfort and Neydel, and Arnold could demonstrate, shows a similar dependence on internal factors. The results of Ewart and Pantanelli are partly based on injury and accumulation of chloroplasts. An increase as described above also proved to exist in my experiments with *Hormidium*. The assimilation was inconstant after a long period of darkness. However, a constant rate of assimilation was reached after a rather short time ($\pm \frac{1}{2}$ hour).

Another reason for the dependence of the assimilation on internal factors is the parallelism with the respiration (with *Hormidium*), which occurs in many cases.

The first striking parallelism between the two processes is that both assimilation and respiration may be stimulated with the same amount of greatly diluted KCN . Moreover, the assimilation is stimulated by the addition of sugars which in the first place may be looked upon as respiratory material, and usually stimulate the respiration. This seems to be important because it refutes a naive application of the mass-law to the process, according to which law a reaction product (sugar) should retard the reaction (photosynthesis) instead of stimulating it.

This parallelism is the more important as an increase in the rate of assimilation results, no matter whether the external conditions are such that light or temperature is limiting.

Other examples of a parallelism between the two processes are instances of a slight retardation of photosynthesis (in weak and strong light!), which are always accompanied by a small decrease of respiration. This was found in case of retardation by KCN , phenylurethane and $BaCl_2$. The recovery of a slight retardation, which always occurs in course of time, proceeds in a similar way in both processes.

The assimilation is more considerable in strong than in weak light. Nevertheless we have seen that it can increase in weak light under changed internal conditions (addition of sugars). So we supposed that the increased assimilation in strong light, besides being a result of the greater quantity of light that can do photosynthetic work (direct action), is also a result of internal factors, changed under the influence of light (indirect action). There is all the more reason for this opinion, as we were able to show that the protoplasm is strongly affected by light. Respiration increased considerably, which is, to be sure, caused by the formation of sugars (ergastogeneous action; at the same time a cause for greater assimilation), but for the greater part light affects the protoplasm directly (plasmogeneous action). The actively assimilating organism also respire actively.

Temperature affects both processes in a similar way. The idea of a possible connection between them would be rather untenable, if this had not been the case. Heating causes a very regular acceleration of the two processes, so that the types of the resulting curves are altogether the same. The coefficients of temperature differ only very little; from 15°—25° Q_{10} is 1.96 for the assimilation, 1.91 for the respiration. The resemblance is so great that it could be accounted for by a full dependence of assimilation on respiration (Spoehr and Mac Gee).

The optimum rate of photosynthesis does not set in immediately after the beginning of the exposure to light. Temperature is of great influence on the duration of this induction-period, by which it is proved that the phenomenon is not founded on photochemical induction (Warburg). So an internal chemical reaction (or a chain of reactions) must precede assimilation. The temperature coefficient of this induction time resembles the coefficient of respiration, though a very accurate determination seems impossible. From 14°—26° Q_{10} amounts to about 2.2.

In addition to our statements of a correlation between assimilation and respiration in *Hormidium*, it may be mentioned here that Plester (1912), Boysen Jensen (1918), Henrici (1918), Spoeher and Mac Gee (1923), Gregory and Richards (1929), found a correlation between the intensity of assimilation and the intensity of respiration in other organisms.

These were the arguments in favour of a relation between assimilation and respiration, now a few will follow that seem to point in another direction. KCN , phenylurethane or BaCl_2 in high concentrations, causing a strong decrease of assimilation, affect the respiration in quite another manner. An entire cessation of the first process may be accompanied by a rather strong or slight retardation, or even by a slight rise of the respiration. This result is in accordance with the results of other investigators (Jacobi, Irving, Thoday, Warburg).

Another discrepancy arose after removal of KCN or BaCl_2 in my experiments. The respiration recovered in this case at once to the initial value, the assimilation recovered incompletely. In case of KCN the assimilation also managed to recover after a longer period, but at all events there was a difference in the time in which a complete recovery set in.

A few more discrepancies were found in the experiments with sugars.

The parallelism of the two processes was to us a ground for the dependence of the assimilation on internal factors, and the few cases in which this parallelism did not hold, certainly cannot invalidate the occurrence of this phenomenon. It is plausible that, when an organism is injured in the first place the normal vital functions as metabolism, growth, propagation are stopped, without death following inevitably. On the contrary; the organism will defend itself against the injurious influences and

consequently may manifest an increased oxygen consumption. It is a well-known fact that e.g. wounding stimulates the consumption of oxygen.

In our case a high concentration of injurious material retards the normal function of the organism, the assimilatory apparatus (or perhaps also an internal regulatory process), suffers direct injury. But the organism is not yet dead, it can defy much larger doses of poison before respiration will at last come to a standstill.

If this opinion is correct the quantitative determination of respiration in these stronger concentrations has lost its importance for the study of photosynthesis.

In exp. 63 I even stated an increase of respiration of 14 % in high concentration of KCN. Irving (1911) and Warburg (1921) found a strong stimulation in high concentrations, in which assimilation is greatly retarded. They found, however, a gradual decrease of respiration after exposure to very high concentrations.

This may show the cause for the difference in Warburg's and my own results. The „pathological” rise of respiration may have started a little earlier in *Chlorella*, and may also be a little stronger than in *Hormidium*, so that a parallelism might have become totally hidden.

So the non-appearance of parallelism in case of strong retardation need not be contradictory to the parallelism observed in case of moderate retardation.

The other discrepancies are no more of great importance. The assimilation, after the poison being removed, did not recover directly, though the respiration had recovered. But ultimately we may also take this as an argument in favour of a connection with internal processes (recovery from injury, which takes time).

The discrepancy mentioned in the chapter on the influence of sugars — a retardation of respiration was

found twice, though the assimilation had increased — is neither very convincing. After the removal of the sugar the respiration appeared to increase, so that the presence of sugar, though beneficial to some processes, appeared to be injurious to others (perhaps osmotic action). So the influence is probably more complicated than one would superficially think (see also Genevois, 1927) and the deviation found is, therefore, no more an important argument.

C. The Explanation of the Parallelism of Respiration and Photosynthesis.

We now come to discuss the question in which way we must explain the parallelism of assimilation and respiration.

1. Spoehr and Mac Gee think that the assimilation is directly dependent on the respiration, and that very active splitting products of carbohydrates, which arise by respiration, play a part in the assimilatory process. We shall see if there is any reason to accept such a direct connection.

2. One may also have the idea that assimilation does not depend directly on the respiratory processes themselves, but on some protoplasmic functions in general, and that, consequently, there is a certain parallelism with the consumption of oxygen, which is so to say a resultant of all protoplasmic functions.

3. Finally we may suppose that this parallelism is not based on a causal dependence of the assimilation on other processes, but that it is based on a joint cause, which acts on all processes in a similar way. We might for instance imagine that the colloidal structure of the plasmatic proteins changes, and consequently the processes which depend on such a structure.

Let us first consider the third case. We can easily see that a retardation of both processes may be caused by a

structural change. The surfaces where the vital processes occur may be destroyed, or taken up by strange substances by which essential substances are expelled.

In this connection it may be mentioned here that Briggs (1922) holds the opinion that the weaker assimilation of plants grown in insufficient nutrient solution, would be caused by a retarded development of the reactive surface.

The above theory could not, however, explain the *recovery* of a function. If this recovery had been found only once, another explanation would certainly have been possible. The phenylurethane might have been oxidized, the KCN dissociated otherwise under the influence of a changed pH (a result of paralysis of function), etc. The very fact that the recovery is the same under all influences (probably also after extreme temperature (cf. p. 612—613), addition of chloroform (Irving), or electrolytes (Shibata) — points, however, to an *active* interference of the organism.

That the alteration of the assimilation under uncommon influences would be simply caused by a passive structural change seems, therefore, very improbable. The organism must always assist with an action of its own, either passively (e.g. more sugars, more oxidation), or actively, which will be the case in a recovery.

So the first mentioned two possibilities remain. Does the respiration supply the energy necessary for the assimilatory process directly ¹⁾ (1), or is the assimilation controlled by definite vital functions, which in their activity are more or less dependent on the total vitality, consequently also on the oxygen consumption, which we may take as a criterion of the vitality (2).

The temperature coefficients of assimilation and respiration do not conflict with the first opinion, as they are

¹⁾ Except, of course, the light energy which is necessary in the first place.

almost equal. The equality offers even a somewhat greater support for this opinion. Dependence on other vital functions might be in accordance with different Q_{10} . It is inessential and improbable that all vital functions of the same organism are affected by temperature in the same way.

Neither does the dissimilar retardation in high concentrations of injurious substances speak against a direct relation, if we assume that the superficial functions are injured, and the increase of respiration has a quite different cause; it might be the result of the active reaction against the injurious influence.

Other facts, however, seem to contradict such a dependence. We found that light causes respiration to increase. However, in the experiments in Chapter XII the agreement is less satisfactory. In those experiments we started to determine both assimilation and respiration. The latter proved to be very high, which seems plausible, since in the month of June the alga got daylight already for a great many hours before the experiment began and the nights were short. It appeared now that respiration fell considerably in the course of the day, in spite of repeated illuminations of short duration. The assimilation, however, was able to rise nearly to the first stated level within a few minutes, without the respiration being interrupted appreciably in its fall. The possibility remains, however, that the respiration did indeed increase during the short period of illumination, but that this short exposure had no after-effect. But even if this is not the case, the hypothesis need not be abandoned yet, though it cannot be maintained unaltered. One may claim that a certain amount of respiration is necessary, but that, as soon as it has been reached, a further increase can only be of little influence. Respiration is here considered to be a factor that can be limiting, or not.

The experiments on the influence of sugars, however, do not agree very well with this amended hypothesis.

Parallelism after a treatment with sugars was not clearly expressed. It is true that respiration increased in general, but this increase did not agree much with that of assimilation. However, it seems to me too early to reject the interesting hypothesis of Spoehr and Mc Gee in the face of my experiments.

Personally I am of the opinion that the remaining second case explains the facts in the most satisfactory way. There is no direct causality between assimilation and respiration, but a relation between assimilation and some protoplasmic functions, which are, however, (like all vital functions) closely connected with respiration. The connection between the two processes of assimilation and respiration may be less close now. We need not be surprised when, after KCN or Ba Cl₂ being removed, respiration recovers more quickly than assimilation, while the somewhat irregular behaviour in sugar-solutions is no more disquieting.

Whatever the explanation of the parallelism of respiration and assimilation may be, we must claim a connection between photosynthesis and other protoplasmic processes.

- a. The equality of the temperature coefficients;
 - b. the parallel stimulation or slight retardation after the addition of chemicals;
 - c. the parallel and simultaneous recovery after retardation;
 - d. a chemical process preceding the assimilation of which Q_{10} is almost equal to the coefficient of respiration;
- are all grounds for the assumption of a close connection between assimilation and other protoplasmic functions.

D. The Reaction of Photosynthesis to Injurious Influences.

We have mentioned repeatedly the reaction to abnormal influences. It may be well to discuss this again at the hand of the diagram represented in fig. 17. The lines 1 and 2 show the cessation of the assimilation after more

or less time. This happened in exp. 66 and 69 after the addition of KCN in exp. 92 after the addition of Ba Cl_2 . A quick, and a successive slow, decrease (line 3) was also found with KCN. Type 4 was very frequently found, a rather strong retardation, followed by recovery. It manifested itself in experiments with the three substances investigated. Type 5, retardation, passing into an increase, resulted in experiments with KCN. Type 6 and type 7




Fig. 17. Diagram to illustrate the various types of reaction of the assimilatory and respiratory processes upon the addition of unusual substances. For explanation see text.

give the experimental results with diluted KCN and sugars. The results obtained all show the gradual transitions between very strong- and almost immediate-retardation and stimulation.

The types 4—7 were also determined for the respiration. In case of a stronger injury of the assimilation, respiration behaves differently.

Another interesting fact was that in the experiments with KCN the effect did not depend on the concentration of this substance. A moderate retardation was obtained with a strong concentration, and a strong retardation another day with a weaker one. Consequently, the incidental disposition of the alga determines whether the effect will be strong or weak.

I had already drawn this diagram, when I was struck by the great resemblance with a figure in Miss Irving's work (1911; fig. 7, p. 1083). This diagram bears on the influence of chloroform on respiration of cherry-laurel leaves and young barley shoots. My types of curves are to be found back there. The concentrations used are rather high, and retarded the assimilation completely in her

experiments. It is, however, of great interest that we find a similar mode of reaction for the respiration of these plants and for the assimilation of *Hormidium*, while this parallelism partly applies to the respiration of this alga.

Shibata (1929) too, gives curves that are fundamentally the same, and in which all my types can be found back. He investigated the antagonistic action of electrolytes on the respiration of *Chlorella*.

The deviations of some points on my temperature-assimilation curve will also be very plausibly explained by assuming the occurring of reaction-type 4, as a result of the action of high or low temperature (cf. the following paragraph).

So it seems that the diagram indicates a very general reaction of the organism on a factor, uncommon to its ordinary conditions of life.

E. An Attempt to Explain the too Large Deviations of Some Determinations on the Influence of Temperature on Photosynthesis from the Average, and a Few Remarks on Optimum Curves.

We saw in the chapter on the influence of temperature on assimilation (Chapter VII) that the deviations of the individual determinations from the average curve were in many cases larger than might be expected in connection with the accuracy of the method. When discussing this fact it was already pointed out that attempts would be made later on to account for these variations with the aid of other results. These results were obtained in the Chapters VIII, IX and XI. It has been discussed in the former paragraph that assimilation reacts in principle in the same way to injurious influences, however different they may be (addition of KCN, phenylurethane or Ba Cl₂): after the initial strong decrease of assimilation a recovery of the function may start.

Now in the first place it strikes us while studying the results of experiments on the influence of temperature, when no high temperatures (30° and upwards) were investigated in the experimental series, deviations did not appear. This shows very distinctly from the experiments carried out in May 1931 (table 8), in which high temperatures were avoided, but also from exp. 38 and 43 (table 6).

An example of great variations is to be found in exp. 42. At 19.5° the assimilation is 317, at 23.5° it is 513, at 31.5° 678, at 23.5 again 449, as we see much lower, in spite of the difference in time of 2 h. 35 min. The last determination in this experiment, also at 19.5° gives 312, in other words the assimilation is again nearly as high as 5 h. 45 min. ago; the assimilation that had been found much too low at 23.5° , has almost completely recovered at 19.5 . This bears a strong resemblance to the results obtained with poisons.

So I think I am able to explain this variation in the determinations from the normal reaction to the injurious influence which in this case had been the high temperature.

In exp. 41 we observe something similar, though a little less pronounced. After a rise of temperature to 30° the next determination at 22° is not higher than the one at 22° , booked 2 h. 45 min. earlier, while the following values at 18° and 14° were higher than before. (The first determination at 18° is rather a little too low).

In experiments 39 and 40 the influence of a lowering in temperature has been investigated. The first values of these experiments are represented by the evidently lowest points at 30° and 29° in fig. 11. This is easy to understand. From the very cold windowsill the algae were suddenly exposed to these high temperatures, so it is a matter of course that the lowering action makes itself strongly felt now.

In exp. 40 the low temperature at 4° has also been investigated. The next reading at 14° turned out 4% lower than the preceding one, in spite of a 5 hours' difference

in time. So a low temperature has apparently the same influence as higher ones. This is a new base for the generality of this reaction. There are, however, no observations as to the appearance of recovery after low temperatures. So we see that the deviations from the average curve can be explained from the physiological reaction to an injurious influence.

Not only the variations of individual determinations, but also the less steep course of the curve at higher temperature (fall of Q_{10}) can be explained by the physiological reaction to injurious stimulants.

It may be mentioned here that, if my explanation is true, this might be to a certain extent a support for Blackman's classical theory about the optimum.

Blackman (1905) is of opinion that a slighter increase of the rate of reaction at high temperature is a result of the injurious action of this temperature, and that in this way the optimum-curves, so often met with in physiology, might be explained. Even before the optimum were reached, a less strong rise of the curve might be caused by the influence of this injurious action.

Therefore, the lower intensity of assimilation when experimenting again at lower temperature might support Blackman's opinion. But the difference between this conception and my own results is, that in my experiments the injury is followed by a recovery.

Neydel (1930), working with *Cladophora*, noted an immediate recovery of the assimilation after the exposure to high temperature. In my own experiments the recovery takes a rather long time, but I believe this difference is not essential.

A transition between Blackman's theory and the conception of the real physiological optimum is apparent in my results. The reversibility of the optimum curve seems to be determined by the time wanted for the recovery,

which may vary between zero (Neydel) and the infinite (Blackman).

I was not able to state a light-intensity optimum of photosynthesis in intensities up to 8000 Lux with algae cultivated in daylight. This is entirely in accordance with the results of van den Honert. However, algae cultivated in weak artificial light, showed a slight decrease of assimilation in high light intensity (fig. 9). In two other cases the assimilation was even brought to a standstill, and the absorption of oxygen began to predominate.

Finally, carbon dioxide did not show any influence in high concentrations in van den Honert's experiments. I can confirm this altogether. Up to a concentration of 1 % the curve is quite parallel with the abscissa and does not show any tendency to fall.

F. Blackman's Rule of Limiting Factors.

In the fifth chapter, in which the relation between CO_2 concentration and assimilation has been established, the peculiar Blackman curve seemed to disappear altogether, when the reaction was retarded by the addition of narcotics, and to show a logarithmic type, established by other investigators (e.g. Harder). One may think here of a slower chemical combination of the CO_2 (or a product derived from it) with other substances. Complications might also be assumed. The question was put whether the respiration with different CO_2 concentrations had perhaps become deviating, but this proved not to be the case in a testing experiment. Whatever the explanation of the CO_2 concentration-assimilation curve after the addition of urethane or antipyrin must be, the remarkable fact remains that the Blackman curve has altogether disappeared.

The light intensity-assimilation curve of van den Honert and the criticism on his method of illumination led to the opinion that a better approach to the Blackman's

scheme might be possible. It is therefore of interest that this did not prove to be so and the curve kept a same logarithmic type with an improved experimental technique, and did not show any tendency to approach a straight curve with a sharp break.

The preceding considerations, in which the probability of an indirect action of external factors was claimed on more than one ground, makes it very unlikely that Blackman-curves will often be met with. These curves sometimes appear in incidental circumstances, such as the CO_2 concentration-assimilation curve of van den Honert.

Summary of the Principal Results.

A very simple, but at the same time very accurate manometrical apparatus is described. Films of only a single cell layer of the filamentous alga *Horridium flaccidum* or the sub-species *H. nitens*, served as experimental objects.

A parallelism between assimilation and respiration was repeatedly observed. Both processes can be stimulated by the addition of highly diluted KCN or of sugar to about the same degree. Not too strong a retardation of photosynthesis, caused by higher concentrated KCN , by BaCl_2 or phenylurethane is accompanied by a decrease of respiration. This parallelism disappears in case of a strong retardation of assimilation. When the retardation is not too strong a recovery will follow. This occurs simultaneously and in about the same measure for both processes.

Temperature affects them both in a similar way; Q_{10} is at moderate temperatures rather constant: it is from 15° to 25° for the assimilation 1.96, for the respiration 1.91.

The influence of various substances on assimilation is of equal strength, whether light or temperature is the limiting factor, which for KCN is contrary to the results

of Warburg, whose theory, therefore, may not be said to be of general validity.

Another fact found by Warburg in the case of *Chlorella*, viz. the impossibility of retarding the assimilation below the compensation-point of assimilation and respiration by means of KCN, could not be confirmed with *Hormidium*.

It will last a few minutes before the assimilation after a period of darkness reaches a constant final speed. This phenomenon is highly sensitive to temperature, and consequently is not caused by photochemical induction (Warburg). A chemical process must precede the beginning of the assimilation. The Q_{10} of this process is, determined roughly, 2.2, and of about the same magnitude as that of respiration and assimilation.

After a long period of darkness an initial increase of photosynthesis, was found, but after half an hour it proved to be fairly constant.

The facts mentioned led to the conclusion that the assimilation is dependent on an internal factor. If this internal factor is the extent of the available, active surface — a supposition made by Briggs in order to explain subnormal assimilation of young and starving plants — then this factor is nevertheless directly dependent on other active internal factors (as was a.o. proved by the recovery of the function after the addition of toxic chemicals).

Respiration is increased during the exposure to light. It could be made plausible that this increase cannot be due to a rise in temperature. The increase is partially caused by the production of carbohydrates, but for the most important part it is a result of the action of light on the protoplasm.

Light, therefore, causes other protoplasmic reactions in addition to photosynthetic. Consequently light may affect the assimilatory process indirectly.

The light intensity-assimilation curve showed, under improved experimental conditions, the normal image of a logarithmic curve. It does not at all approach the scheme proposed by Blackman.

The Blackman-curve, which was found by van den Honert for the influence of CO_2 , disappears when the assimilation is retarded by phenylurethane or antipyrin.

Finally the assimilatory-quotient of *Hormidium* approximates to unity.

I started this investigation at the Botanical Laboratory of Utrecht University and I should like to express my grateful thanks to Professor F. A. F. C. Went for the undiminished interest he took in my work up till the end.

By far the greater part of the investigation was carried out at the Laboratory of Technical Botany at Delft and I am very much obliged to Professor G. van Iterson for his valuable help, wherefore I am thanking him particularly.

Literature.

- F. Aereboe, *Forsch. a. d. Gebiete d. Agrikult. phys.* 16, 429, 1893.
 J. E. van Amstel, *Rec. trav. bot. néerl.* 13, 1, 1916.
 A. Arnold, *Planta* 13, 529, 1931.
 E. Beljakoff, *Planta* 8, 269, 1929.
 C. Bernard, *Leçons sur les phénomènes de la vie*, etc. Paris, 1878.
 F. F. Blackman, *Ann. of Bot.* 19, 281, 1905.
 ——— and G. L. C. Matthaei, *Proc. Roy. Soc. B*, 76, 402, 1905.
 ——— and A. M. Smith, *Proc. Roy. Soc. B*, 83, 374, 1911.
 S. R. de Boer, *Rec. trav. bot. néerl.* 25, 117, 1927.
 Chr. Bohr, *Ann. Physik u. Chem. N. F.* 62, 644, 1897; *Ann. Phys. u. Chem. N. F.* 68, 500, 1899.
 A. E. H. R. Boonstra, *Planta* 10, 108, 1930.
 J. Borodin, *Just's Bot. Jahresber.* 4, Abt. II, 919, 1876; *Mém. acad. imp. sci. St. Pétersbourg*, VIIe Sér. 28, No. 4, 1, 1881.
 J. C. Bose, *The Physiology of Photosynthesis*. London 1924.
 P. Boysen Jensen, *Bot. Tidskr.* 36, 219, 1918.
 ——— and D. Müller, *Jahrb. f. wiss. Bot.* 70, 493, 1929; *Jahrb. f. wiss. Bot.* 70, 503, 1929.

- G. E. Briggs, *Proc. Roy. Soc. B*, 94, 20, 1922.
 W. H. Brown, *Philippine Journ. Sci. C. Bot.* 13, 345, 1918.
 ——— and G. W. Heise, *Phil. Journ. Sci. C. Bot.* 12, 1, 1917;
 Phil. Journ. Sci. C. Bot. 12, 85, 1917.
 R. Emerson, *Journ. Gen. Physiol.* 10, 469, 1927; *Journ. Gen. Physiol.* 12, 609, 1929; *Journ. Gen. Physiol.* 12, 623, 1929.
 A. J. Ewart, *Journ. Linn. Soc. Bot.* 31, 364, 1896; *Journ. Linn. Soc. Bot.* 31, 554, 1897; *Ann. of Bot.* 12, 415, 1898.
 L. Genevois, *Biochem. Zeitschr.* 186, 461, 1927.
 F. G. Gregory and F. J. Richards, *Ann. of Bot.* 43, 119, 1929.
 R. Harder, *Jahrb. f. wiss. Bot.* 56, 254, 1915; *Jahrb. f. wiss. Bot.* 60, 531, 1921; *Planta* 11, 263, 1930.
 M. Henrici, *Verhandl. nat.-forsch. Ges. Basel*, 30, 43, 1919.
 G. Holst, *Electrische lichtbronnen en haar eigenschappen*. Haarlem, 1920.
 T. H. van den Honert, *Diss. Utrecht*, 1928; *Rec. trav. bot. néerl.* 27, 149, 1930.
 A. A. Irving, *Ann. of Bot.* 25, 1077, 1911.
 B. Jacobi, *Flora* 86, 289, 1899.
 N. Johansson, *Svensk Bot. Tidskr.* 17, 215, 1923; *Svensk Bot. Tidskr.* 20, 107, 1926.
 H. Kniep, *Internat. Rev. Hydrobiol. u. Hydrogr.* 7, 1, 1914.
 S. Kostytschew, *Planta* 13, 778, 1931.
 ——— K. Bazyrina and W. Tschesnokov, *Planta* 5, 696, 1928.
 ——— and V. Berg, *Planta* 11, 144, 1930.
 ——— and H. Kardo—Syssoiewa, *Planta*, 11, 117, 1930.
 ——— M. Kudriavzewa, W. Moissejewa and M. Smirnowa, *Planta* 1, 679, 1926.
 ——— and S. Soldatenkow, *Planta* 2, 1, 1926.
 ——— W. Tschesnokov and K. Bazyrina, *Planta* 11, 160, 1930.
 A. Krogh, *The Respiratory Exchange of Animals and Man*. London 1916; *Biochem. Journ.* 14, 267, 1920.
 Tsi-Tung Li, *Ann. of Bot.* 43, 587, 1929.
 A. Löwshin, *Beih. Bot. Centralbl.* 23, Abt. I, 54, 1908.
 W. Lubimenko, *Comp. rend. acad. sci.* 143, 609, 1906.
 H. Lundegårdh, *Svensk Bot. Tidskr.* 15, 46, 1921; *Der Kreislauf der Kohlensäure in der Natur*. Jena 1924; *Biochem. Zeitschr.* 154, 195, 1924; *Flora* 121, 273, 1927.
 L. Maquenne and E. Demoussy, *Comp. rend. acad. sci.* 156, 506, 1913.
 G. L. C. Matthaei, *Phil. Trans. Roy. Soc. London*, B. 197, 47, 1904.

- N. A. Maximow and T. A. Krasnosselsky—Maximow, Ber. deuts. bot. Ges. 46, 383, 1928.
- A. Meyer and N. T. Deleano, Zeitschr. f. Bot. 3, 657, 1911; Zeitschr. f. Bot. 5, 225, 1913.
- C. Montfort and K. Neydel, Jahrb. f. wiss. Bot. 68, 801, 1928.
- K. Neydel, Biochem. Zeitschr. 228, 451, 1930.
- K. Noack, Naturwiss. 14, 383, 1926.
- W. J. V. Osterhout and A. R. C. Haas, Journ. Gen. Physiol. 1, 1, 1918; Journ. Gen. Physiol. 1, 295, 1918.
- E. Pantanelli, Jahrb. f. wiss. Bot. 39, 167, 1904; Ber. deuts. bot. Ges. 32, 547, 1914.
- A. Pascher, Die Süßwasserflora Deutschlands, Oesterreichs und der Schweiz. H. 6: Chlorophyceae III. Jena, 1914.
- H. Plaetzer, Verhandl. phys. med. Ges. Würzburg, 45, 31, 1917.
- L. Plantefol, Ann. sci. nat. Bot. 10e Sér. 9, 1, 1927.
- W. Plester, Beitr. Biol. Pflanzen, 11, 249, 1912.
- I. Plotnikow, Abderhalden Handb. Biol. Arbeitsmeth. III, A 2, 1653.
- Th. Sabalitschka and H. Weidling, Biochem. Zeitschr. 176, 210, 1926.
- Th. Schmucker, Biochem. Zeitschr. 195, 149, 1928.
- M. Shibata, Sci. Reports Tôhoku Imp. Univ. 4e Ser. 4, 431, 1929.
- H. A. Spoehr, Photosynthesis, New York 1926.
- and J. M. Mc Gee, Carnegie Inst. Washington, Publ. 325, Washington 1923.
- W. Stiles, Photosynthesis, London 1925.
- O. Stocker, Flora 121, 334, 1927.
- D. Thoday, Ann. of Bot. 27, 565, 1913.
- O. Treboux, Flora 92, 49, 1903.
- O. A. Walther, Flora 121, 301, 1927.
- O. Warburg, Biochem. Zeitschr. 100, 230, 1919; Biochem. Zeitschr. 103, 188, 1920; Biochem. Zeitschr. 119, 134, 1921; Naturwissenschaften 9, 354, 1921.
- and E. Negelein, Zeitschr. physik. Chem. 102, 236, 1922.
- and T. Uyesugi, Biochem. Zeitschr. 146, 486, 1924.
- R. Willstätter and A. Stoll, Untersuchungen über die Assimilation der Kohlensäure, Berlin 1918.
- M. Yabusoe, Biochem. Zeitschr. 152, 498, 1924.
- Y. Yoshii, Planta 5, 681, 1928.