

3.2

Soort	Zuid-Holland + IJmuiden			Hondsbossche			TOTAAL
	→ Z	→ N	aantal uren waarin de soort werd gezien + percentage	→ Z	→ N	aantal uren waarin de soort werd gezien + percentage	
Scholekster +	2351	6142	669 = 76 %	1078	2529	608 = 65 %	12100
Kievit	131	67	28 = 3 %	375	40	41 = 4 %	613
Goudplevier	11	44	11 = 1 %	2	2	4 < 1 %	59
Zilverplevier	42	13238	278 = 32 %	11	3788	228 = 25 %	17279
Bontbekplevier	23	5614	169 = 19 %	12	606	63 = 7 %	6255
Strandplevier	-	-	-	1	2	2 < 1 %	3
Kleine plevier	1	7	7 = 1 %	-	5	3 < 1 %	13
Steenloper +	34	333	174 = 20 %	79	320	172 = 19 %	766
Watersnip	4	3	4 < 1 %	-	4	3 < 1 %	11
Houtsnip	-	-	-	1	-	1 < 1 %	1
Bokje	2	-	2 < 1 %	-	-	-	2
Wulp	106	466	171 = 20 %	47	624	129 = 14 %	1243
Regenwulp	32	678	123 = 14 %	23	229	86 = 9 %	962
Grutto	5	114	16 = 2 %	17	60	19 = 2 %	196
Rosse grutto	104	1075	180 = 20 %	73	3979	276 = 30 %	5231
Oeverloper +	-	294	63 = 7 %	-	133	25 = 3 %	427
Witgatje	-	4	3 < 1 %	-	1	1 < 1 %	5
Tureluur	26	3429	249 = 28 %	35	469	115 = 12 %	3959
Zwarte ruiter	-	36	12 = 1 %	-	10	6 < 1 %	46
Groenpootruiter	1	277	75 = 9 %	2	45	25 = 3 %	325
Kanoetstrandl.	120	12844	171 = 20 %	9	13110	196 = 21 %	26083
Paarse strandl.+	2	36	71 = 8 %	-	3	3 < 1 %	41
Bonte strandl.	168	1808	230 = 26 %	42	6292	281 = 30 %	8308
Krombekstrandl.	-	14	5 < 1 %	-	3	4 < 1 %	17
Kleine strandl.	-	12	4 < 1 %	-	-	-	12
Drieteenstrandl.+	142	1825	235 = 27 %	18	1209	128 = 14 %	4194
Kemphaan	-	41	12 = 1 %	-	4	1 < 1 %	45
Kluut	9	340	61 = 7 %	4	94	21 = 2 %	447