

3.2.

Soort	Zuid-Holland + IJmuiden			Hondsbossche Zeewering			TOTAAL
	→ Z	→ N	aantal uren waarin de soort werd gezien + percentage	→ Z	→ N	aantal uren waarin de soort werd gezien + percentage	
Scholekster +	6828	465	526 - 48 %	942	1060	415 - 62%	9.301
Kievit	20251	160	94 - 9 %	10122	53	95 - 14 %	30.586
Goudplevier	1753	14	71 - 6 %	215	7	14 - 2 %	1.989
Zilverplevier	202	27	96 - 9 %	44	20	32 - 5 %	293
Bontbekplevier	594	17	134 - 12 %	61	107	41 - 6 %	779
Strandplevier	2	-	1	-	1	1	3
Kleine plevier	3	-	2	3	-	1	6
Steenloper	159	29	174 - 16 %	320	479	208 - 31 %	987
Watersnip	75	8	26 - 2 %	98	9	17 - 3 %	196
Houtsnip	-	-	1	-	-	-	-
Bokje	6	-	4	-	-	-	6
Wulp	375	47	143 - 13 %	73	33	41 - 6 %	528
Regenwulp	408	19	91 - 8 %	55	33	58 - 9 %	515
Grutto	6	-	4	1	-	1	7
Rosse grutto	222	138	151 - 14 %	124	131	77 - 12 %	615
Oeverloper +	52	5	74 - 7 %	79	49	27 - 4 %	203
Witgatje +	-	-	3	2	-	3	2
Bosruiter	-	1	1	-	1	1	2
Tureluur	466	143	138 - 12 %	63	693	133 - 20 %	1.365
Zwarte ruiter	4	-	4	1	2	3	7
Groenpootruiter	44	2	14 - 1 %	21	6	16 - 2 %	73
Kanoetstrandloper	574	1	48 - 4 %	98	697	84 - 13 %	1.370
Paarse strandl. +	52	1	97 - 9 %	2	2	8 - 1 %	57
Bonte strandloper	1909	79	265 - 24 %	378	439	155 - 23 %	2.805
Krombekstrandl.	2	2	3	1	-	1	5
Kleine strandl.	10	-	8 - 1 %	-	1	1	11
Temminck strandl.	1	-	-	-	-	-	1
Drieteenstrandl.	404	211	132 - 12 %	48	174	65 - 10 %	837
Kemphaan	4	2	5	1	1	4	8
Kluut	53	-	6	3	2	2	58
Grauwe franjepoot	2	-	2	-	-	-	2
Rosse franjep. +	2	-	6	1	-	1	3