

## NIEUW BESCHREVEN CONTINENTALE MOLLUSKENSOORTEN (NEW TAXA: CONTINENTAL MOLLUSCS)

R.A. Bank


Newly described taxa (rubricated at the family level), with reference to the publications in which they were introduced. Ordered per family, families sorted alphabetically.

### ACAVIDAE

- *Cylindroclavator*, *Paraclavator*, *Embertoniphanta*, *Nigriphanta* and *Spiriphanta*. New (sub)genera described by Groh & Poppe (2002, A conchological Iconography) from Madagascar.

### ARIOPHANTIDAE

- Zonitacionini and Acantharionini. New tribes described by A.A. Schileyko (2002, Treatise on Recent terrestrial molluscs, 9: 1167-1710).
- *Celectella*. Nomen novum introduced by A.A. Schileyko (2002, Treatise on Recent terrestrial molluscs, 9: 1167-1710) for *Euplectella* Jousseaume, 1894.



*Ariophanta (Celectella) trimeni* - Schileyko, 2002, Treatise Pulmonate Molluscs, 9: fig. 1688

### CHONDRINIDAE

- *Chondrina falkneri* and *Chondrina gerhardi*. Described by E. Gittenberger (2002, Collectanea Malacologica: 41-44) from the French dépts. Drôme and Isère (Massif de la Chartreuse and Vercors region) and dépt. Hautes-Alpes (Combe de Queyras), respectively.
- *Rupestrella rupestris margritae*, *R. r. carolae*, *R. r. lamellosa*, *R. homala masae* and *R. jaeckeli*. Described by K.-H. Beckmann (2002, Collectanea Malacologica: 49-78) from Sicily (Italy).

### CLAUSILIIDAE

- *Triloba thaumasia talevi*, *Carinigera pelucida* and *Balea (Pseudalinda) nordsiecki*. Described by I. Dedov & E. Neubert (2002, Archiv für Molluskenkunde, 131 (1-2): 201-209) from the Macedonian part of the Shar Mountains.

- *Tropidaucheniini* (new tribe) and *Neniauchenia* (new subgenus). Described by H. Nordsieck (2002, Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie), 640: 1-23) within the subfamily Garnieriinae.
- *Garnieria mouhoti moellendorffi*, *G. saurini*, *Tropidauchenia (Neniauchenia) amoena* and *T. (N.) dautzenbergi decollata*. New taxa described by H. Nordsieck (2002, Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie), 640: 1-23) from Laos.
- *Tropidauchenia (Indonenia) excellens*. Described by H. Nordsieck (2002, Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie), 640: 1-23) from Burma, Carin Mountains, 1200-1300 m.
- *Tropidauchenia (Grandinenia) cinderella*. Described by H. Nordsieck (2002, Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie), 640: 1-23) from China, Guangxi, 8 km SW. Yangshuo.
- *Tropidauchenia (Tropidauchenia) donggiaoensis*, *T. (T.) fischeri fischeri*, *T. (T.) fischeri reticulata*, *T. (T.) giardi citrina* and *T. (T.) giardi brunnea*. Described by H. Nordsieck (2002, Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie), 640: 1-23) from Vietnam.
- *Macroptychia jickelii*, *M. obesa*, *M. producta* and *M. distorta*. Described by E. Neubert (2002, Collectanea Malacologica: 253-275) from Eritrea (area N. of Keren), Somalia (Gan Libah), Ethiopia (Harar, mountain area around Mt. Batu) and Mt. Abune Yoséf (prov. Welo, Ethiopia), respectively.
- Euxinellini and Olympicolini. New tribes of the subfamily Mentissoideinae described by E. Neubert (2002, Collectanea Malacologica: 253-275).
- *Albinaria caerulea pholegandrica* and *A. c. scaletta*. Described by H. Nordsieck (2002, Collectanea Malacologica: 291-299) from the island Folegandros (Cyclades, Greece).
- *Pararillya* and *Rillyella*. New genera described by H. Nordsieck (2002, Collectanea Malacologica: 343-361) for Paleocene taxa of the Paris basin.
- *Rillya uva*, *Rillyella chenayensis* and *Neniopsis cossmanni*. Described by H. Nordsieck (2002, Collectanea Malacologica: 343-361) from Chenay, Sables de Châlons-sur-Vesle (France, Paris basin, Late Paleocene deposits).
- *Macedonica pindica pindica*, *M. p. bellu-*

*la*, *Montenegrina janinensis maasseni*, *M. dennisi dennisi*, *M. d. protruda* and *Idyla castalia yeruni*. - Described by E. Gittenberger (2002, Basteria, 66 (4-6): 129-138) from Greece, Macedonia, Nomos Grevena, west and southwest of the town Grevena.

- *Strumosa strumosa meridiana*. Described by H. Schütt (2002, Malakologische Abhandlungen Staatliches Museum für Tierkunde Dresden, 20 (2, 22): 213-217) from prov. Isparta, Turkey.

### COCHLICELLIDAE


- *Monilearia montigena*. New name introduced by R.A. Bank, K. Groh & Th.E.J. Ripken (2002, Collectanea Malacologica: 89-235) for *Helix lemniscata* Webb & Berthelot, 1833 (Canary Islands, Gran Canaria).

### CYCLOPHORIDAE

- *Japonia lanyuensis*, *Platyrphe lanyuensis*, *P. sunggsangensis*, *Cyclotus taivanus dilatus*, *C. t. diminutus* and *Leptopoma tigris*. Described by Y.-C. Lee & W.-L. Wu (2001, Bulletin of Malacology, 25: 45-78) from Taiwan.
- *Cyathopoma camerunense*. Described by A.J. de Winter (2002, Collectanea Malacologica: 237-251) from Cameroon, Sud Province.

### DIPLOMMATINIDAE

- *Diplommatina abundans*, *D. carinaspinoosa*, *D. gaduensis*, *D. karoensis*, *D. vanderblommi*, *D. wilhelminae*, *Palaina reederi*, *Plectostoma kitteli*, *Opisthostoma banki*, *O. clerxi* and *O. secretum*. Described by W.J.M. Maassen (2002, Basteria, 66 (4-6): 163-182) from Sumatra (Indonesia).



*Opisthostoma hemistreptum* - Van Benthem Jutting, 1961, Bull. Raffles Mus., 26: pl. 14 fig. 12.

## HELICARIONIDAE

- Papuarioninae, Papuarionini, Laocaini, Atoxoniini and Buetneriini. New subfamily and new tribes described by A.A. Schileyko (2002, Treatise on Recent terrestrial molluscs, 9: 1167-1710).

## HELICIDAE

- *Klika orbiculata* and *Helicigona planata*. Described by H. Binder (2002, Beiträge zur Paläontologie, 27: 161-203) from the Karpatian (lower Miocene) deposits of the Korneburg basin, Austria.

## HYDROBIIDAE

- *Hydrobia djerbaensis*. Described by T. Wilke, M. Pfenninger & G.M. Davis (2002, Proceedings of the Academy of Natural Sciences of Philadelphia, 152: 45-66) from the Tunisian island Djerba.
- *Heraultiella*. Nomen novum introduced by M. Bodon, G. Manganelli & F. Giusti (2002, Journal of Conchology, 37 (6): 681) for *Heraultia*.
- *Spathogyna*. New genus described by B. Arconada & M.A. Ramos (2002, Journal of Molluscan Studies, 68 (4): 319-327) for the Spanish taxon *fezi* Altimira, 1960.
- *Bythinella wawrzinekii*. Described by R. Bernasconi (2002, Documents Malacologiques, 3: 9-12) from Saint-Bris-des-Bois, dépt. Charente-Maritime, France.
- *Bythinella bouloti* and *Bythinella galerae*. Described by H. Girardi, J.-M. Bichain & M. Wienin (2002, Documents Malacologiques, 3: 43-54) from Castelbouc and Montbrun, dépt. Lozère, France.
- *Bythiospeum diaphanum michaellensis*. Described by H. Girardi (2002, Documents Malacologiques, 3: 57-64) from Caromb, dépt. Vaucluse, France.

## HYGROMIIDAE

- *Leucochroopsis phaseolina papillata*. Described by H. Binder (2002, Beiträge zur Paläontologie, 27: 161-203) from the Karpatian (lower Miocene) deposits of the Korneburg basin, Austria.
- *Canariella (Canariella) falkneri*. Described by M.R. Alonso, M. Ibáñez & C.E. Ponte-Lira (2002, Collectanea Malacologica: 79-87) from La Gomera (Canary Islands).
- *Leptaxis (Katostoma) nivosa calensis*. New name introduced by R.A. Bank, K. Groh & Th.E.J. Ripken (2002, Collectanea Malacologica: 89-235) for *Helix [Katostoma] phlebophora* var. *planata* R.T. Lowe, 1855 (Madeira: Ilhéu de Baixo ou da Cal).

## HYPSELOSTOMATIDAE


- *Boysidia (Bensonella) tongguanensis*. Described by D.-N. Chen & W.-H. Zhang (2002, Acta Zootaxonomica Sinica, 27 (4): 690-698) from Lijia Town, Tongguan County, Shaanxi Province, China.

## LAURIIDAE

- *Leiostyla (Leiostyla) falknerorum*. New name introduced by R.A. Bank, K. Groh & Th.E.J. Ripken (2002, Collectanea Malacologica: 89-235) for *Pupa [Leiostyla] laevigata* R.T. Lowe, 1852 (Madeira).

## MAIZANIIDAE

- *Ankoravaratra* new genus and the new species *A. ambalaniranae*, *A. ambrensis*, *A. analamerae* and *A. capdambrae*. Described from K.C. Emberton (2002, The Veliger, 45 (4): 278-289) from northern Madagascar.
- *Maizaniella (Spirulozania) falkneri*. Described by A.J. de Winter (2002, Collectanea Malacologica: 237-251) from Cameroon, Sud Province.



*Maizaniella falkneri* - De Winter, 2002, Collectanea Malacologica: fig. 2

## MELANOPSIDAE


- *Melanopsis tchernovi*. Described by J. Heller & N. Sivan (2002, Journal of Conchology, 37 (6): 607-625) from the late Pliocene of 'Erq el-Ahmar, Jordan Valley, Israel.

## NERITIDAE

- *Nerita (Amphinerita) bartonensis*. Described by M.F. Symonds (2002, Tertiary Research, 21 (1-4): 1-10) from the Barton Group (Middle Eocene), Highcliff, Hampshire, southern England.

## PACHYCHILIDAE

- *Brotia praetermissa*. Described by F. Köhler & M. Glaubrecht (2002, Journal of Molluscan Studies, 68 (4): 353-357) from Borneo.
- *Jagora*. New genus described by F. Köhler & M. Glaubrecht (2003, Zoologica Scripta, 32 (1): 35-59) for *Melania asperata* Lamarck, 1822 and *Melania dactylus* I. Lea & H.C. Lea, 1850.



*Brotia praetermissa* - Köhler & Glaubrecht, 2002, J. Moll. Stud., 68: fig. 1

## PUPINIDAE

- *Pupina acehensis* and *Pupina falkneri*. Described by W.J.M. Maassen (2002, Collectanea Malacologica: 277-290) from Leupung (prov. Aceh, Sumatra) and near the entrance of the Cave Liangdehar (Kuta Buluh, Sumatra).

## RHYSOTINIDAE

- Rhysotinidae. New family described by A.A. Schileyko (2002, Treatise on Recent terrestrial molluscs, 9: 1167-1710).

## STREPTAXIDAE

- *Gulella* (8 new species) and *Parvedentulina* (83 new species). Described by K.C. Emberton (2002, Archiv für Molluskenkunde, 131 (1-2): 67-165) from Madagascar.
- *Gibbulinella dewinteri*. New name introduced by R.A. Bank, K. Groh & Th.E.J. Ripken (2002, Collectanea Malacologica: 89-235) for *Pupa [Gibbulinella] dealbata* var. *minor* Mousson, 1872 (Canary Islands: La Palma, La Gomera, Tenerife).

## SUBULINIDAE

- *Allopeas myrmekophilos*. Described by R. Janssen (2002, Archiv für Molluskenkunde, 131 (1-2): 211-215) from ant colonies 30 km NE. of Kuala Lumpur, Malaysia.

## TRIPTYCHIDAE

- *Triptychia obliqueplicata austriaca* and *T. suturalis gracilis*. Described by H. Binder (2002, Beiträge zur Paläontologie, 27: 161-203) from the Karpatian (lower Miocene) deposits of the Korneburg basin, Austria.

## VIVIPARIDAE

- *Albianopalin* new genus and the new species *A. benkeari* and *A. lizsmithae*. Described by R.J. Hamilton-Bruce, B.J. Smith & K.L. Gowlett-Holmes (2002, Records of the South Australian Museum, 35 (2): 193-203) from the Early Cretaceous (middle-late Albian) Griman Creek Formation of Lightning Ridge, northern New South Wales, Australia.