

**Notes on Bulimulidae (Gastropoda, Euthyneura), 3¹
Bulimulus trinidadensis, sp.n. from
Ilha da Trindade, Brazil**

Abraham S.H. BREURE

Department of Systematic Zoology/of the University,
c/o Rijksmuseum van Natuurlijke Historie, Leiden

& Arnaldo C. DOS SANTOS COELHO

Museu Nacional, Rio de Janeiro

During a recent visit² of the first author to the Museu Nacional, Rio de Janeiro, some material turned up of an apparently undescribed species of *Bulimulus* s.str. We take the opportunity to describe this material here as:


***Bulimulus trinidadensis* sp.n.**

Diagnosis — A small and slender species of the genus *Bulimulus* s.str. with rather convex sides. The columellar margin is fold-like inside the aperture.

Description — Shell up to 9.0 mm long, 2.2 times as long as wide, narrowly perforate, acute with rather convex sides, rather solid, more or less elongate. Colour light yellowish brown, turning light brown towards apex. Surface slightly shining, with traces of spiral striation in some specimens. Growth lines may be slightly stronger below the

¹ Notes on Bulimulidae, 2, see *Basteria* 39: 97-113, 1975.

² This visit was made possible by a grant of the Foundation for the Advancement of Tropical Research (WOTRO).



Figs. 1-4. *Bulimulus trinidadensis* sp. n. 1. Paratype M.N. 3712(A). 2. Upper whorls of same specimen. 3. Penultimate whorl of same specimen. 4. Holotype M.N. 3711. Scale 1 mm.

suture. Protoconch sculptured with axial costulae forming fine wrinkles, more or less anastomosing on lower part, while also very fine spiral striae may be observed. Whorls about seven, hardly convex. Last whorl about 6/10 of the height of the shell. Suture hardly impressed. Aperture subovate, about 1.6 times as long as wide, 4/10 of the height of the shell. Peristome thin, hardly expanded. Some specimens with slight, knoblike swelling on parietal region inside the aperture. Columellar margin sinuous, dilated over umbilicus, inside the aperture knob- or fold-like. Margins joined by a callus.

Type locality — Brazil, Ilha da Trindade, Praia dos Cabritos (about 20° 30' S 29° 20' W). "The material was found imbedded in the loose sediment on the terrace near the beach, away from the tide line" [A.S.C.].

Material — Holotype Museu Nacional (M.N.) 3711; (many) paratypes M.N. 3712; 72 paratypes Rijksmuseum van Natuurlijke Historie, RMNH 55081; 1 paratype Zoologische Staatssammlung, München; 1 paratype Naturhistorisches Museum, Wien. All material J. Becker leg., 28-XII-1958.

Remarks — This species does not resemble any of the other species of *Bulimulus*. The distribution of *Bulimulus* s.s. being predominantly limited to eastern South America, makes the locality of the present novelty fall well outside the range as the distance to the nearest locality of a *Bulimulus* s.s. species (*B. tenuissimus* d'Orbigny, 1835) is ca. 1500 km in a straight line.

Some overall resemblance may be noted when the new species is compared to some species of the genus *Itaborahia* Maury, 1935, from the fossil beds near Rio de Janeiro. However, that genus is sufficiently characterized by the sculpture of the protoconch (axial riblets) and the different position of the teeth in the aperture. Perhaps *B. trinidadensis* might be regarded as a conservative species in morphological respect within the genus.

Table 1. Measurements of *Bulimulus trinidadensis* sp.n. (in mm).

shell height	diameter	aperture height	width	height of last whorl	number of whorls	
8.0	3.5	3.4	1.9	4.8	7.0	holotype
7.8	3.6	3.3	2.3	4.7	6.9	RMNH 55081
9.0	3.8	3.5	2.3	5.0	7.3	idem
8.0	3.8	3.3	2.1	4.8	5.9	idem
8.3	3.4	3.2	2.1	4.5	6.2	idem
7.8	3.7	3.1	2.0	4.5	6.8	idem
8.4	3.6	3.3	2.1	4.9	6.9	idem
7.8	3.6	3.1	2.1	4.4	6.9	idem
8.5	3.8	3.3	2.2	4.8	7.3	idem
8.2	3.7	3.3	2.1	4.9	7.0	idem
8.2	3.7	3.3	2.0	4.8	7.1	idem
8.3	3.8	3.7	2.3	5.0	5.9	idem
7.5	3.5	3.1	2.1	4.3	6.8	idem
7.8	3.6	3.1	2.0	4.6	6.7	idem