

Two overlooked host fish species of *Margaritifera auricularia*
(Bivalvia, Unionoidea, Margaritiferidae)

Rafael ARAUJO

Museo Nacional de Ciencias Naturales, José Gutiérrez Abascal 2, E 28006 Madrid, Spain

Key words: Bivalvia, Unionoidea, Margaritiferidae, *Margaritifera auricularia*, ecology, host fish.

Nienhuis (2003) recently published an article on French populations of *Pseudunio auricularius* (Spengler, 1793) (= *Margaritifera auricularia*). He apparently overlooked that *Salaria fluviatilis* Asso, 1801, the river blenny, and *Acipenser baeri* Brandt, 1869, the Siberian sturgeon, were already identified by Araujo, Bragado & Ramos (2001) as additional host species for the glochidia of *M. auricularia*, next to the Atlantic sturgeon, *Acipenser sturio* L., 1758.

More recently it has been proved that the Adriatic sturgeon (*Acipenser naccarii* Bonaparte, 1836), is yet another suitable host for successful metamorphosis of the glochidium of *M. auricularia* (Araujo, Quirós & Ramos, 2003).

REFERENCES

- ARAUJO, R., D. BRAGADO & M.A. RAMOS, 2001. Identification of the river blenny, *Salaria fluviatilis*, as a host to the glochidia of *Margaritifera auricularia*. — Journal of Molluscan Studies 67: 128-129.
- ARAUJO, R., M. QUIROS & M.A. RAMOS, 2003. Laboratory propagation and culturing of juveniles of the endangered freshwater mussel *Margaritifera auricularia* (Spengler, 1793). — Journal of Conchology 38: 53-60.
- NIENHUIS, J.A.J.H., 2003. The rediscovery of Spengler's freshwater pearl mussel *Pseudunio auricularius* (Spengler, 1793) (Bivalvia, Unionoidea, Margaritiferidae) in two river systems in France, with an analysis of some factors causing its decline. — Basteria 67: 67-86.