

Literatuuropgaven uit niet-malacologische tijdschriften

Biggs, H.E., 1969. A contribution to the census of the non-marine mollusca of Suffolk. Trans. Suffolk Nat. Soc. 14: 182-188.

Burchell, J.P.T., 1969. Incidence and distribution of the molluscan fauna in certain lateral valleys of the Thames and Medway from the beginning of late glacial times until the end of the sub-boreal period. Zool.J. Linn. Soc. 48: 113-115.

Cameron, R.A.D., 1969. The distribution and variation of three species of land snail near Rickmansworth, Hertfordshire. Zool.J. Linn. Soc. 48: 83-111. (Arianta arbustorum, Cepaea nemoralis, C.hortensis).

- " -, 1969. Predation by song thrushes Turdus ericetorum (Turton) on the snails Cepaea hortensis (Müll.) and Arianta arbustorum (L.) near Rickmansworth. J. Anim. Ecol. 38: 547-555.

Dance, S.P., 1969. Re-discovery of the wall whorl snail in South Wales. Nature Wales 11: 161-163. (Vertigo pusilla)

Fogan, M., 1969. Vitrina (Semilimax) pyrenaica (Férussac) in Kerry North. Ir. Nat. J. 16: 175.

- Gibson, J.S., 1970. The function of the operculum of Thais lapillus (L.) in resisting desiccation and predation. *J. Anim. Ecol.* 39: 159-168.
- Hughes, R.N., 1970. Population dynamics of the bivalve Scrobicularia plana (Da Costa) on an intertidal mud-flat in North Wales. *J. Anim. Ecol.* 39: 333-356.
- " -, 1970. An energy budget for a tidal-flat population of the bivalve Scrobicularia plana (Da Costa). *J. Anim. Ecol.* 39: 357-382.
- James, B.L., 1968. The distribution and keys of species in the family Littorinidae and of their digenean parasites, in the region of Dale, Pembrokeshire. *Fld. Stud.* 2: 615-650. (L.littorea, L.littoralis, L.neritoides + subspecies en varieteiten, L.saxatilis. Beschrijving van juveniele Littorina soorten en subspecies),
- Ladle, M. & Baron, F., 1969. Studies on three species of Pisidium (Mollusca: Bivalva) from a stalk stream. *J. Anim. Ecol.* 38: 407-415. (P.subtruncatum, P.nitidum, P.milium).
- McMillan, N.F., 1969. The slug Agriolimax caruanae Pollonera in Ireland. *Ir. Nat. J.* 16: 178.
- Norton, P.E.P., 1967. Marine molluscan assemblages in the early Pleistocene of Sidestrand, Bramerton and the Royal Society Borehole at Ludham, Norfolk. *Phil. Trans. R. Soc. B*, 253: 161-200.
- O'Riordan, C.E., 1968. Lutraria angustior Philippi: A lamellibranch mollusc new to Ireland. *Ir. Nat. J.* 16: 47-48.
- Pallant, D., 1969. The food of the grey field slug (Agriolimax reticulatus (Müller)) in woodland. *J. Anim. Ecol.* 38: 391-399.
- Phillips, B.F. & Campbell, N.A., 1970. Comparison of methods of estimating population size using data on the whelk Dicathais aegrota (Reeve). *J. Anim. Ecol.* 39: 753-761.
- Prebble, L.C., 1966. Conchological notes on the land and freshwater mollusca of the Isle of Wight. *Proc. Isle Wight Nat. Hist. Archaeol. Soc.* 5: 442-443.
- Seed, R., 1968. Factors influencing shell shape in the mussel Mytilus edulis. *J. Mar. Biol. Ass. U.K.* 48: 561-584.
- Wolda, H., 1969. Stability of a steep cline in morph frequencies of the snail Cepaea nemoralis. *J. Anim. Ecol.* 38: 623-637.
- " -, 1969. Fine distribution of morph frequencies in the snail Cepaea nemoralis near Groningen. *J. Anim. Ecol.* 38: 305-329.

E.X. Maier & J. van Raam